



**Health
Information
and Quality
Authority**

An tÚdarás Um Fhaisnéis
agus Cáilíocht Sláinte

Health technology assessment (HTA) of smoking cessation interventions - APPENDICES

05 January 2017

Table of Contents

Table of Contents	2
List of abbreviations used in this report	3
Appendices	4
Appendix 1 Search details for review of smoking cessation interventions among users of secondary mental health services	4
Appendix 2 Included studies for those with a diagnosis of schizophrenia or schizoaffective disorder	6
Appendix 3 Included studies for those with a diagnosis of bipolar disorder	10
Appendix 4 Studies from the UK Centre for Tobacco Control Studies (UKCTCS) report that were excluded from this review.....	11
Appendix 5 Studies from the Cochrane review of smoking cessation in schizophrenia that were excluded from this review.....	14
Appendix 6 Studies from the Cochrane review of smoking cessation in current and past depression that were excluded from this review.....	16
Appendix 7 Search details for review of smoking cessation interventions among unselected adults	19
Appendix 8 Included studies for unselected adults	21
Appendix 9 Search details for review of smoking cessation interventions in pregnancy	112
Appendix 10 Included studies smoking cessation interventions in pregnancy	114
Appendix 11 Studies smoking cessation interventions in pregnancy that were excluded from this review.....	136
Appendix 12 Cost-effectiveness plane showing the ICER for each strategy when all strategies that are more clinically effective are excluded (with e-cigarettes included as a treatment option)	138
Appendix 13 References to included studies.....	150

List of abbreviations used in this report

BIOCHEM	biochemically validated abstinence
CAR	continuous abstinence rate
COPD	chronic obstructive pulmonary disease
EAG	Expert Advisory Group
EMA	European Medicines Agency
ENDS	electronic nicotine delivery system
ENNDS	electronic non-nicotine delivery system
FDA	Food and Drugs Authority
HIQA	Health Information and Quality Authority
HPRA	Health Products Regulatory Authority
HR	hazard ratio
HSE	Health Service Executive
HTA	health technology assessment
ICER	incremental cost-effectiveness ratio
NICE	National Institute for Health and Care Excellence
NRPA	nicotine receptor partial agonist
NRT	nicotine replacement therapy
OR	odds ratio
PCRS	Primary Care Reimbursement Scheme
PP	point prevalence abstinence
QALY	quality-adjusted life year
RCT	randomised controlled trial
RR	relative risk
SR	self reported
UK	United Kingdom
WHO	World Health Organization

Appendices

Appendix 1 Search details for review of smoking cessation interventions among users of secondary mental health services

Figure 1.1 Flow diagram of included studies from the updated search




Table 1.1 Search string used in PubMed

Search string	Results
<pre> ((((((((bipolar[Title/Abstract]) OR depress*[Title/Abstract]) OR schizophren*[Title/Abstract]) OR mental health[Title/Abstract]) OR mental illness*) OR mental disorder*[Title/Abstract]) OR "Mental Health Services"[Mesh]) OR "Mental Disorders"[Mesh])) AND (("Tobacco Use Cessation products"[Mesh]) OR "Tobacco Use Cessation"[Mesh]) OR "Smoking"[Mesh]) OR ((((abstinence[Title/Abstract]) OR cessat*[Title/Abstract]) OR quit*[Title/Abstract]) OR stop*[Title/Abstract])) AND (((cigarette*[Title/Abstract]) OR tobacco[Title/Abstract]) OR smok*[Title/Abstract])) AND ((Clinical Trial[ptyp] OR Randomized Controlled Trial[ptyp] OR Clinical Trial, Phase I[ptyp] OR Clinical Trial, Phase II[ptyp] OR Clinical Trial, Phase III[ptyp] OR Clinical Trial, Phase IV[ptyp] OR systematic[sb] OR Review[ptyp]) AND ("2012/01/01"[PDat] : "2016/12/31"[PDat]) AND Humans[Mesh]) </pre>	1,424

Appendix 2 Included studies for those with a diagnosis of schizophrenia or schizoaffective disorder

Table 2.1 Bupropion + behavioural intervention with/without NRT versus placebo + behavioural intervention with/without NRT

Study	Country	Setting	Participants	Mean age	Female	Population	Intervention	Control	Support	Outcome	Risk of bias
Evins 2001 ⁽¹⁾	USA	Community volunteers	19	44.1	42%	DSM-IV diagnosis of schizophrenia	Bupropion 150 mg/day for 12 wks	Placebo for 12 wks	Both groups received brief advice and 9 wkly 1-hour sessions of group CBT	6m (biochem, PP)	High
Evins 2005 ⁽²⁾	USA	Community volunteers	57	45.7	31.6%	DSM-IV diagnosis of schizophrenia or schizoaffective disorder, depressed type.	Bupropion 300 mg/day for 12 wks (150 mg/day for first wk)	Placebo for 12 wks	Both groups received 12 wkly 1-hour sessions of group CBT.	6m (biochem and self-reported, CAR, PP)	High
Evins 2007 ⁽³⁾	USA	Community volunteers	51	44.2	NR	DSM-IV diagnosis of schizophrenia	Bupropion SR 300 mg/day for 12 wks. (150 mg/day for first 7 days)	Placebo for 12 wks	Both received 1) 12 wkly 1-hour sessions of group CBT; (2) TNP (from wk 4) 21 mg/day for 4 wks, then 14 mg/day for 2 wks, 7 mg/day for 2 wks + up to 18 mg per day NG as required	6m, 12m (biochem, PP)	Low
George 2002 ⁽⁴⁾	USA	Community volunteers	32	43.2	43.8%	DSM-IV diagnosis of schizophrenia or schizoaffective disorder. 20 had	Bupropion 300 mg/day for 10 wks (150 mg/day for first 3	Placebo for 10 wks	Both groups received 10 wkly 1-hour sessions of group therapy for motivational	6m (biochem, CA)	Unclear

						a diagnosis of schizophrenia.	days)		enhancement, psychoeducation and relapse prevention		
George 2008 ⁽⁵⁾	USA	Community volunteers	59	40.3	40.7%	DSM-IV diagnosis of schizophrenia or schizoaffective disorder. 32 had a diagnosis of schizophrenia.	Bupropion 300 mg/day for 10 wks (150 mg/day for first 3 days)	Placebo for 10 wks	Both groups received 10 wky 50-minute sessions of GBT and TNP (21 mg per 24 hours) from day 15 to day 70	6m (PP, biochem CA)	Unclear

Table 2.2 Varenicline + behavioural intervention versus placebo + behavioural intervention

Study	Country	Setting	Participants	Mean age	Female	Population	Intervention	Control	Support	Outcome	Risk of bias
Williams 2012 ⁽⁶⁾	USA, Canada	Community volunteers	128	41.1	23.4%	DSM-IV TR diagnosis of schizophrenia or schizoaffective disorder. 91 had schizophrenia. 109 pts on atypical antipsychotic	Varenicline (0.5mg daily for 3 days then twice daily for 4 days then 1mg twice daily) for 12 wks	Placebo for 12 wks	Both groups received wky counselling (less than 30 minutes)	6m (biochem, PP)	Unclear

Table 2.3 Individual counselling + NRT versus routine care

Study	Country	Setting	Participants	Mean age	Female	Population	Intervention	Control	Additional Support	Outcome	Risk of bias
Baker 2006 ⁽⁷⁾	Australia	Community volunteers	298	37.2	48%	ICD-10 diagnosis of	Individually administered SC	Routine care	Both groups received	6m, 12m, 4yrs (CA,	High

						psychotic disorder. 126 had a diagnosis of schizophrenia and 43 a diagnosis of schizoaffective disorder	intervention (6 wkly sessions & 2 boosters at wks 8 and 10, 1 hour each): based on MI and CBT + TNP (21 mg from wk 3 (quit date set) to 8; 14 mg from week 9 to 10; 7 mg from wk 11 to 12)		booklets on smoking cessation	biochem, PP)	
--	--	--	--	--	--	---	--	--	-------------------------------	--------------	--

Table 2.4 High intensity individual counselling + NRT versus lower intensity individual counselling + NRT

Study	Country	Setting	Participants	Mean age	Female	Population	Intervention	Control	Additional Support	Outcome	Risk of bias
Williams 2010 ⁽⁸⁾	USA	Community volunteers	100	45.2	45%	DSM-IV diagnosis of schizophrenia or schizoaffective disorder.	High intensity: TANS (treatment of nicotine addiction in schiz): 24 individual sessions of 45 minutes psychological intervention over 26 wks (MI, social skills training, use of NRT, relapse prevention technique)	Medication Management: 9 individual sessions of 20 minutes psychological intervention over 26 wks (medication compliance, education about NRT)	Both groups also received TNP for 16 wks after quit date (21 mg for 12 wks then 14 mg for remaining 4 wks)	6, 12m (PP, biochem)	High

Table 2.5 Group behavioural intervention (generic) + NRT versus group behavioural intervention (tailored to schizophrenia) + NRT

Study	Country	Setting	Participants	Mean age	Female	Population	Intervention	Control	Additional Support	Outcome	Risk of bias
George 2000 ⁽⁹⁾	USA	Unclear	45	39.7	33.3%	DSM-IV diagnosis of schizophrenia (n=19) or schizoaffective disorder.	Specialised SC group therapy designed for pts with schizophrenia, wkly for 10 wks (60 minutes each session): first 3 wks motivational enhancement therapy + last 7 wks psychoeducation, social skills training and relapse prevention strategy.	American Lung Asso. (ALA) programme wkly for 10 wks (60 minutes each session): first 7wks behavioural group therapy + final 3 wks supportive group counselling	Both groups received TNP (21 mg/day for first 6 wks then 14 mg/day for another 2 wks and 7 mg/day for final 2 wks)	6m (CA, biochem + SR)	High

Appendix 3 Included studies for those with a diagnosis of bipolar disorder

Table 3.1 Varenicline + brief advice versus placebo + brief advice

Study	Country	Setting	Participants	Mean age	Female	Population	Intervention	Control	Additional Support	Outcome	Risk of bias
Chengappa 2014 ⁽¹⁰⁾	USA	Outpatient clinics	60	46	37%	DSM-IV bipolar disorder	Varenicline	Placebo	Smoking cessation counselling provided to all patients	6m (PP, biochem at 12 weeks only)	Low

Appendix 4 Studies from the UK Centre for Tobacco Control Studies (UKCTCS) report that were excluded from this review

	Study (year)		Reason for exclusion
1	Akbarpour 2010	Outcome	Outcome not applicable (reduction in smoking - abstinence not defined or measured).
2	Axtmayer 2011	Outcome	Outcome not applicable ('number of cigarettes smoked per day' - not smoking cessation).
3	Baker 2009	Study design	Study design not applicable (uncontrolled before and after study).
4	Barnett 2008	Intervention	Intervention not applicable (participants received \$25 payment for each research assessment).
5	Bloch 2010	Outcome	Outcome not applicable (reduction in smoking - abstinence not defined or measured).
6	Batra 2010	Population	Current depression population - not clear if all attending secondary mental health services.
7	Brown 2003	Population	Population not applicable, (people aged <18 years old).
8	Brown 2007	Population	Current depression population - not clear if all attending secondary mental health services.
9	Catley 2005	Population	Current depression population - not clear if all attending secondary mental health services.
10	Chen 2002	Short-term	Short-term only (7 day point prevalence quit rates at 4 and 8 weeks). No 6-month cessation data.
11	Chou 2004	Short-term	Short-term only (cessation measured at 8 weeks only).
12	Cornelius 1997	Intervention	Intervention not applicable (fluoxetine).
13	Culhane 2008	Short-term	Short-term only (12 weeks only).
14	Currie 2008	Study design	Study design not applicable (not random assignment).
15	Dalack 1999	Outcome	Outcome not applicable (reduction in smoking, abstinence not defined or measured).
16	de Leon 2005b	Intervention	Intervention not applicable (clozapine).
17	Dixon 2009	Study design	Patients not randomised. For each clinic pair, one clinic was randomly assigned to the “immediate” implementation condition and the other to the “delayed” implementation condition
18	Dutra 2012	Comparator	No comparison group. Uncontrolled before-and-after study
19	Fatemi 2005	Outcome	Outcome not applicable (reduction in smoking, abstinence not defined or measured).
20	Gallagher 2007	Intervention	Intervention not applicable (includes contingency payments).
21	Hall 2006	Population	Current depression population - not clear if all attending secondary mental health services.
22	Hartman 1991	Outcome	Outcome not applicable (reduction in smoking, abstinence not defined or measured).
23	Hertzberg 2001	Population	Population not applicable (PTSD in male combat veterans).

24	Hill 2007	Short-term	Non-randomised pilot study, outcomes measured at 3 months only.
25	Kelly 2008	Intervention	Intervention not applicable (galantamine).
26	Kinnunen 2008	Population	Current depression population - not clear if all attending secondary mental health services.
27	Kisely 2003	Study design	Study design not applicable, non-randomised cross-over study.
28	Levine 2010	Population	Current depression, but not clear if all attending secondary mental health services.
29	Li 2009	Short-term	Short-term only, cessation measured at 1, 2, 4 and 8 weeks.
30	McEvoy 2005	Intervention	Intervention not applicable (clozapine).
31	McFall 2005	Population	Population not applicable, (PTSD in veterans).
32	McFall 2010	Population	Population not applicable, (PTSD).
33	Morris 2011	Outcome	Outcomes not applicable (50% reduction and not cessation).
34	Panchas 2012	Study design	Study design not applicable - before-and-after study without randomisation
35	Rabius 2007a	Population	Current depression population - not clear if all attending secondary mental health services.
36	Rabius 2007b	Population	Current depression population - not clear if all attending secondary mental health services.
37	Roll 1998	Intervention	Intervention not applicable (contingency payments).
38	Saxon 2003	Population	Population and intervention not applicable: veterans in treatment for alcohol or drug dependency.
39	Smith 2009	Short-term	Short-term follow-up (9 weeks only).
40	Steinberg 2003	Outcome	Outcome not applicable (reduction in smoking, abstinence not defined or measured).
41	Szombathyne 2010	Outcome	Outcome (reduction in smoking, abstinence not defined or measured) and intervention (altrexone) not applicable.
42	Thorsteinsson 2001	Short-term	Short-term follow-up period (29 days).
43	Tidey 2002	Intervention	Intervention not applicable (contingency payments).
44	Tidey 2011	Outcome	Outcome not applicable (reduction in smoking, abstinence not defined or measured).
45	Van der Meer 2010	Population	Current depression population - not clear if all attending secondary mental health services.
46	Weinberger 2008a	Study design	Study design not applicable (pilot study of 5 people).
47	Weiner 2001	Comparator	No comparison group.
48	Weiner 2011 a	Short-term	Short-term only (cessation measured at 4 and 12 weeks only).
49	Weiner 2011 b	Short-term	Short-term only (cessation measured at 12 weeks only).
50	Williams 2007	Short-term	Short-term only (cessation measured at 8 weeks only).

51	Wotjna 2009	Study design	Study design (lack of randomisation).
----	-------------	--------------	---------------------------------------

Appendix 5 Studies from the Cochrane review of smoking cessation in schizophrenia that were excluded from this review

	Study (year)		Reason for exclusion
1	Akbarpour 2010	Outcome	Outcome is reduction in smoking; abstinence not defined or measured.
2	Bloch 2010	Outcome	Outcome is reduction in smoking; abstinence not defined or measured.
3	Brown 2003	Population	<18 years old
4	Chen 2012	Short-term	Abstinence reported at 8 weeks only.
5	Dalack 1999	Outcome	Outcome is reduction in smoking; abstinence not defined or measured.
6	de Leon 2005b	Intervention	Clozapine
7	Dutra 2012	Comparator	No comparison group. Uncontrolled before-and-after study
8	Fatemi 2005	Outcome	Outcome is reduction in smoking, abstinence not defined or measured.
9	Gallagher 2007	Intervention	Includes contingency payments
10	Gelkopf 2012	Outcome	Outcome is reduction in smoking; abstinence not defined or measured.
11	Hartman 1991	Outcome	Outcome is reduction in smoking; abstinence not defined or measured.
12	Hong 2011	Outcome	Effect of varenicline on neurobiological and cognitive biomarkers in schizophrenia. Abstinence not defined or measured.
13	Horst 2005	Outcome	Relapse prevention after smoking cessation. Abstinence not defined or measured.
14	Kelly 2008	Intervention	Galantamine
15	Li 2009	Short-term	Cessation measured at 1, 2, 4 and 8 weeks only.
16	McEvoy 1995	Intervention	Clozapine
17	McEvoy 1999	Intervention	Clozapine
18	Meszaros 2012	Outcome	Effect of varenicline on alcohol dependence (as primary outcome according to protocol). Abstinence not defined or measured.
19	Pachas 2012	Study design	Before and after study without randomisation
20	Roll 1998	Intervention	Contingency payments
21	Sacco 2009	Intervention	Atomoxetine

22	Shim 2012	Outcome	Outcome is reduction in smoking; abstinence not defined or measured.
23	Steinberg 2003	Outcome	Outcome is reduction in smoking; abstinence not defined or measured.
24	Szombathyne 2010	Outcome Intervention	Outcome is reduction in smoking; abstinence not defined or measured. Naltrexone
25	Tidey 2002	Intervention	Contingency payments
26	Tidey 2011	Outcome	Outcome is reduction in smoking; abstinence not defined or measured.
27	Weinberger 2008	Intervention	Topiramate
28	Weiner 2001	Comparator	No comparison group
29	Weiner 2011 a (var)	Short-term	Cessation measured at 4 and 12 weeks only.
30	Weiner 2012	Short-term	Cessation measured at 12 weeks only.
31	Williams 2007	Short-term	Cessation measured at 8 weeks only.
32	Wing 2012	Intervention	Active Repetitive Transcranial Magnetic Stimulation (rTMS) - bilateral to dorsolateral prefrontal cortex v Sham rTMS

Appendix 6 Studies from the Cochrane review of smoking cessation in current and past depression that were excluded from this review

	Study (year)		Reason for exclusion
1	Batra 2010	Population	Current depression population; not clear if all attending secondary mental health services.
2	Blondal 1999	Intervention	Fluoxetine
3	Brown 2001	Population	Past depression
4	Brown 2007	Population	Current depression population; not clear if all attending secondary mental health services.
5	Carmody 2008	Intervention	Includes hypnosis
6	Catley 2003	Intervention	Culturally-sensitive self-help materials
7	Catley 2005	Population	Current depression population; not clear if all attending secondary mental health services.
8	Cinciripini 2010	Population	Pregnant smokers
9	Covey 1999	Intervention	Naltrexone
10	Covey 2002	Intervention	Sertraline
11	Duffy 2006	Population	Smokers with head and neck cancer
12	Hall 1994	Population	Past depression
13	Hall 1996	Population	Subgroup of past depression only
14	Hall 1998	Population	Past depression
15	Hall 2002	Population	Past depression. Only information available which compared nortriptyline versus bupropion (CR)
16	Hall 2006	Population	Current depression population; not clear if all attending secondary mental health services.
17	Hall 2009	Population	Subgroup of past depression only
18	Hayes 2010	Population	Medically ill smokers using home healthcare nursing services with current depressive symptoms
19	Hayford 1999	Population	Past depression
20	Kahler 2008	Population	Heavy drinking and smoking adults
21	Killen 2000	Intervention	Paroxetine
22	Killen 2008	Population	Current depression population; not clear if all attending secondary mental health services.
23	Kinnunen 2008	Population	Current depression population; not clear if all attending secondary mental health services.

24	Kodl 2008	Population	Adult alcohol dependent smokers
25	Levine 2003	Population	Current depression population; not clear if all attending secondary mental health services.
26	Levine 2010	Population	Current depression population; not clear if all attending secondary mental health services.
27	MacPherson 2010	Population	Mildly elevated depressive symptoms (BDIII \geq 10). Recruited through community and University, so not clear if attending secondary health. Intervention = BATS
28	McAlister 2004	Population	Mild depression: depression according to a single self-report question: "Have you felt sad or blue every day for the last two weeks?" Recruited through community and quitline so not clear if attending secondary mental health.
29	McFall 2010	Population	PTSD related to military service
30	Muñoz 1997	Intervention	Quit smoking guide (Spanish: Guía) + mood management
31	Muñoz 2006a	Intervention	Guía + Individually Timed Educational Messages + mood management course versus Guía + Individually Timed Educational Messages
32	Muñoz 2006b	Intervention	Guía + Individually Timed Educational Messages + mood management course versus Guía + Individually Timed Educational Messages
33	Muñoz 2009	Intervention	Guía + Individually Timed Educational Messages + mood management course versus Guía + Individually Timed Educational Messages versus Guía + Individually Timed Educational Messages + mood management course + virtual group versus Guía alone
34	Patten 1998	Population	Abstinent alcoholics
35	Piper 2009	Population	Not clear if attending secondary mental health services.
36	Rabius 2007a	Population	Current depression population; not clear if all attending secondary mental health services.
37	Rabius 2007b	Population	Current depression population; not clear if all attending secondary mental health services.
38	Rabius 2008	Intervention	Tailored interactive site versus control (minimally interactive) site
39	Sauls 2004	Intervention	Fluoxetine
40	Schnoll 2010	Population	Cancer and depressive symptoms
41	Smith 2003	Population	Subgroup of past depression only
42	Spring 2007	Intervention	Fluoxetine
43	Swan 2010	Population	Not clear if attending secondary mental health services.
44	Thorndike 2008	Population	Acute cardiovascular disease
45	Van der Meer	Population	Current depression population; not clear if all attending secondary mental health services.

	2010		
46	Vickers 2009	Intervention	Exercise counselling
47	Walsh 2008	Intervention	Naltrexone
48	Weinberger 2010b	Intervention	Selegiline hydrochloride (antidepressant)
49	Wewers 2009	Population	PC and women's health clinics that served a socioeconomically diverse population and reported >200 Pap smears per month were invited to participate in the study. Not clear if attending secondary mental health.

Appendix 7 Search details for review of smoking cessation interventions among unselected adults

Figure 7.1 Flow diagram of included studies from the updated search




Table 7.1 Sample search strategy: Updated review for NRT (Medline)

Pubmed 18/07/2016	Search strings	Results
Searches #1	((("Smoking Cessation"[Mesh]) OR ("Tobacco Use Cessation"[Mesh]) OR (Stop* smoking) OR (Quit* smoking)))	31023
#2	((((((((((NRT) OR (nicotine AND patch*)) OR (nicotine AND gum)) OR (nicotine AND nasal AND spray)) OR (nicotine AND lozenge*)) OR (nicotine AND tablet*)) OR (nicotine AND sublingual)) OR (nicotine AND inhal*)) OR (nicotine AND replacement)) OR (nicotine AND therap*))	10421
#3	#1 AND #2	4353
#4	#3 AND Filters: <ul style="list-style-type: none"> • <u>From 2012/05/01 to 2016/07/18</u> • Clinical Trial; Randomized Controlled Trial; Review; Systematic Reviews; Meta-Analysis • Humans 	451

Search string: (((((((((((NRT) OR (nicotine AND patch*)) OR (nicotine AND gum)) OR (nicotine AND nasal AND spray)) OR (nicotine AND lozenge*)) OR (nicotine AND tablet*)) OR (nicotine AND sublingual)) OR (nicotine AND inhal*)) OR (nicotine AND replacement)) OR (nicotine AND therap*)) AND (((("Smoking Cessation"[Mesh]) OR ("Tobacco Use Cessation"[Mesh]) OR (Stop* smoking) OR (Quit* smoking))))

Appendix 8 Included studies: Unselected adults

Table 8.1 Included studies from systematic review: nicotine gum versus placebo

Study	Country	Setting	Participants	Mean Age	Female	Intervention	Control	Support	Outcome	Risk of bias
Ahluwalia 2006⁽¹¹⁾	USA	Community volunteers	755	45	67%	Nicotine gum (2 mg), recommended use tailored to cpd. Highest 10/day for 4 wks, tapering for 4 wks	Placebo gum, 8 wks	3 in-person visits at randomization, wk 1, wk 8, and phone contact at wk 3, wk 6, wk 16, content based on either motivational interviewing or health education principles	6m (biochem, pp)	Low
Areechon 1988⁽¹²⁾	Thailand	Community volunteers	200	39	6%	Gum (2 mg) x 8 boxes	Placebo gum x 8 boxes	Weekly visits with physician	6m (biochem, pp)	High
Blondal 1989⁽¹³⁾	Iceland	Community volunteers	182	42	57%	Gum (4 mg) for at least 1m	Placebo gum (containing pepper) for 1m or more	Group therapy, 5x1hr sessions, TQD at session 1	12m (biochem, car)	Unclear
Campbell 1987⁽¹⁴⁾	UK	Primary care	836	39	61%	Nicotine gum (2 mg) x 6 boxes	Placebo gum x 6 boxes	No further face-to-face contact, 2/3rds received a letter after 1m	12m (biochem, car)	Unclear
Clavel 1985⁽¹⁵⁾	France	Community volunteers	427	34	51%	Nicotine gum (2 mg) x 1 box	Control group (time lock controlled cigarette case)	3x1hr group therapy sessions in first month	13m (biochem, car)	High
Clavel-Chapelon	France	Community volunteers	996	34	45%	Nicotine gum (2 mg) for up	Placebo gum (contained 1	3 acupuncture session at 0, 7, 28	13m (self, NR)	High

1992⁽¹⁶⁾						to 6m, max 30/day	mg unbuffered nicotine)	days		
Cooper 2005⁽¹⁷⁾	USA	Community volunteers	439	38	100%	Nicotine gum (2mg), 10-12 pieces/day recommended, for 9 wks, weaning last 3 wks.	Placebo gum	13x1hr weekly cognitive behavioural group sessions. Reduction prior to TQD wk 5	12m (biochem, pp)	Unclear
Fagerstrom 1982⁽¹⁸⁾	Sweden	Smoking cessation clinic	100	NS	59%	Nicotine gum (2 mg) for at least 4 wks	Placebo gum for at least 4 wks	Individual counselling, average 7.7 sessions	6m (biochem, pp)	Unclear
Fagerstrom 1984⁽¹⁹⁾	Sweden	Primary care	145	40	56%	Nicotine gum (2 or 4 mg) + long or short follow-up	Advice + long/short follow-up	Low	12m (self, car)	High
Fee 1982⁽²⁰⁾	UK	Smoking cessation clinic	352	NS	NS	Gum (2 mg) given for 5 wks	Placebo gum given for 5 wks	10 group therapy sessions	12m (biochem, pp)	Unclear
Fortmann 1995⁽²¹⁾	USA	Community volunteers	1044	40	42%	Nicotine gum (2 mg, 1 per hr, at least 10/day and not more than 30/day)	Incentive alone	Low	12m (biochem, pp)	Unclear
Garcia 1989⁽²²⁾	Spain	Primary care	106	36	65%	Gum (2 mg) for 3-4m	Placebo gum for 3-4m		6m (biochem, car)	Unclear
Garvey 2000⁽²³⁾	USA	Community volunteers	608	NS	51%	4 mg nicotine gum (recommended 9-15 pieces), weaning from 2m; 2 mg nicotine gum,	Placebo gum	Brief counselling (5-10 mins) at each study visit (1, 7, 14, 30 days, 2, 3, 6, 9, 12m)	12m (biochem, car)	High

						use as 1.				
Gilbert 1989⁽²⁴⁾	Canada	Primary care	223	NR	NS	Support from physician plus offer of nicotine gum prescription (2 mg)	Support from physician (no placebo)	Enrolment, quit day, offer of 4 support visits, 2 in wk 1, 1m, 2m	12m (biochem, car)	High
Gross 1995⁽²⁵⁾	USA	Community volunteers	177	42	51%	Nicotine gum (2 mg), tapered from wk 12. Active gum groups further randomized to chew 7, 15 or 30 pieces of gum.	No gum	1 pre-quit group counselling session, 14 clinic visits in 10 wks	6m (biochem, car)	High
Hall 1985⁽²⁶⁾	USA	Community volunteers + physician referrals	120	38	47%	Combined - 2mg nicotine gum (period of use not specified) and intensive behavioural treatment	Intensive behavioural treatment (14 group sessions over an 8 wk period)	High	12m (biochem, NR)	High
Hall 1987⁽²⁷⁾	USA	Community volunteers	139	39	47%	Nicotine gum (2 mg) up to 12m	Placebo gum up to 12m	Both group-based, 14x75 min sessions, or 5x60min sessions	12m (biochem, pp)	Unclear
Hall 1996⁽²⁸⁾	USA	Community volunteers	207	40	52%	Nicotine gum (2 mg) for 8 wks, 1 piece/hr for 12 hrs/day recommended	Placebo gum, same schedule	Group-based, 10 sessions over 8 wks, TQD session 3	12m (biochem, car)	Unclear
Harackiewicz 1988⁽²⁹⁾	USA	Primary care	197	36	63%	Nicotine gum (2 mg, 6 wks)	Self help manual;	Single appointment with doctor or nurse,	12m (biochem,	High

						initial supply, suggested tapering after 3m, available for 6m) plus self help manual	Control (booklet)	length not specified	car)	
Herrera 1995⁽³⁰⁾	Venezuela	Community volunteers	322	38	43%	2 mg nicotine gum (low dependence)	Placebo gum (low dependence)	12 group sessions over 6 wks + 6 weekly maintenance sessions	24m (biochem, car)	Unclear
Hjalmarson 1984⁽³¹⁾	Sweden	Smoking cessation clinic	206	42	56%	Nicotine gum (2 mg) (no restrictions on amount or duration of use)	Placebo gum	6 group sessions in 6 wks	12m (biochem, car)	Unclear
Huber 1988⁽³²⁾	Germany	Community volunteers	225	NS	NS	Nicotine gum (no details of dose) and behaviour therapy	Behaviour therapy, 5 weekly group meetings	High	12m (self, NR)	Unclear
Hughes 1989⁽³³⁾	USA	Primary care	315	37	56%	Nicotine gum (2 mg for 3-4m)	Placebo gum	29-35 min at 1st visit including nurse & physician advice, & materials, follow-up appointment 1-2 wks later	12m (biochem, car)	Low
Hughes 1990⁽³⁴⁾	USA	Community volunteers	78	40	54%	Nicotine gum - 1mg/2mg/4mg	Placebo gum	29-35 min at 1st visit including nurse & physician advice, & materials, follow-up appointment 1-2 wks later	6m (self, car)	Unclear
Jamrozik 1984⁽³⁵⁾	UK	Primary care	200	NS	NS	Nicotine gum (2 mg) for	Placebo gum	Follow-up visits at 2, 4, 12 wks for data	6m (biochem,	Unclear

						3m+		collection	pp)	
Jarvis 1982⁽³⁶⁾	UK	Smoking cessation clinic	116	40	55%	Nicotine gum (2 mg) unrestricted amount for at least 3m	Placebo gum (1 mg unbuffered nicotine)	Group therapy 6x1hr weekly	12m (biochem, car)	High
Jensen 1991⁽³⁷⁾	Denmark	Smoking cessation clinic	293	42	54%	Nicotine gum (2 mg for 3m)	Standard chewing gum	9 group meetings over a year, weekly to wk 4	12m (biochem, car)	High
Killen 1984⁽³⁸⁾	USA	Community volunteers	64	44	72%	Nicotine gum (2 mg) for 7 wks; Skills training plus nicotine gum	Skills training	Group therapy	10.5m (biochem, car)	High
Killen 1990⁽³⁹⁾	USA	Community volunteers	1218	43	52%	Nicotine gum (2 mg, 8 wks) ad lib dosing; Nicotine gum on a fixed dose	Placebo gum; No gum	Randomized to 1 of 3 psychological interventions	12m (biochem, pp)	Unclear
Llivina 1988⁽⁴⁰⁾	Spain	Smoking cessation clinic	216	NR	NR	Nicotine gum (dose not stated) for 1m	Placebo gum	Group support	12m (biochem, car)	Unclear
Malcolm 1980⁽⁴¹⁾	UK	Community volunteers	194	45	41%	Nicotine gum (2 mg) at least 10/day for at least 3m	Placebo gum; Control	Weekly individual counselling for 1m	6m (biochem, car)	Unclear
McGovern 1992⁽⁴²⁾	USA	Community volunteers	293	NS	NS	ALA Freedom from Smoking clinic program plus nicotine gum (2 mg for 3m)	ALA Freedom from Smoking clinic program alone (no placebo gum)	Group support	12m (biochem, pp)	Unclear

Nakamura 1990⁽⁴³⁾	Japan	Community volunteers	60	NS	NS	Nicotine gum (2 mg, 2m or longer)	Non-placebo control group received counselling	High	6m (biochem, car)	High
Nebot 1992⁽⁴⁴⁾	Spain	Primary care	425	NS	NS	Physician counselling plus nicotine gum	Brief counselling from physician; health education from nurse	Low	12m (biochem, pp)	High
Niaura 1994⁽⁴⁵⁾	USA	Outpatients	173	42	50%	Nicotine gum 2 mg, ad lib for up to 4m (participants given prescription for gum, not free)	No gum	4 individual counselling sessions and ALA self help treatment manuals	12m (biochem, car)	Unclear
Niaura 1999⁽⁴⁶⁾	USA	Community volunteers	62	28	50%	Intensive cognitive behavioral relapse prevention (CBRP) with cue exposure + nicotine gum	Intensive CBRP with cue exposure	5 group sessions within 3 wks of TQD	12m (biochem, car)	Unclear
Ockene 1991⁽⁴⁷⁾	USA	Primary care	1223	35	57%	Patient-centred counselling and offer of nicotine gum(2mg) plus minimal or intensive follow-up by	Patient-centred counselling	Mixed	12m (self, car)	Unclear

						telephone				
Page 1986 ⁽⁴⁸⁾	Canada	Primary care	275	NS	NS	Advice to quit plus offer of nicotine chewing gum prescription (2 mg)	No advice; Advice to quit	Low	6m (self, car)	High
Pirie 1992 ⁽⁴⁹⁾	USA	Community volunteers	417	NS	100%	Group therapy plus nicotine gum; Group therapy plus weight control programme and nicotine gum.	Group therapy; Group therapy plus weight control programme	High	12m (biochem, car)	Unclear
Puska 1979 ⁽⁵⁰⁾	Finland	Community volunteers	229	NS	NS	Nicotine gum (4 mg) for 3 wks	Placebo gum for 3 wks	Group therapy	6m (self, pp)	Unclear
Richmond 1993 ⁽⁵¹⁾	Australia	Primary care	450	NS	NS	Smokescreen programme plus nicotine gum, dose and duration not stated	Smokescreen programme alone	5 visits during first 3m	12m (biochem, car)	High
Roto 1987 ⁽⁵²⁾	Finland	Primary care	121	NS	43%	Nicotine gum (2 mg and 4 mg), + advice	Advice only (no placebo)	Low	6m (NR, NR)	Unclear
Russell 1983 ⁽⁵³⁾	UK	Primary care	2106	NS	NS	As group 2, plus offer of nicotine gum prescription, Individual therapy, Single visit 1 minimal content, 1 more intensive	No intervention; Advised to stop smoking plus provided with a "give up smoking" booklet	Low	12m (biochem, car)	High

						content, untrained therapist				
Schneider 1985A⁽⁵⁴⁾	USA	Community volunteers	60	38	60%	Nicotine gum, (2 mg duration not stated)	Placebo gum	Individual support at multiple clinic assessment visits, daily during week 1, weekly to wk 5	12m (biochem, car)	Unclear
Schneider 1985B⁽⁵⁴⁾	USA	Community volunteers	36	NS	NS	Nicotine gum, (2 mg duration not stated)	Placebo gum	Weekly laboratory visits for 5 wks but no support provided	12m (biochem, car)	Unclear
Segnan 1991⁽⁵⁵⁾	Italy	Primary care	923	NS	NS	Repeated counselling plus prescription for nicotine gum unless contraindicate d, dose not stated, up to 3m	Advice and leaflet; Repeated counselling (follow-up at 1, 3, 6, 9m); Repeated counselling plus spirometry	High	12m (biochem, car)	Low
Shiffman 2009 (2mg)⁽⁵⁶⁾	USA	Community volunteers	1636	42	64%	Nicotine gum 2mg. Instructed to gradually reduce smoking while increasing gum use for up to 8 wks. Postquit instructed to use 1 piece every 1-2hrs for first 6 wks; 1 every 2-4hrs for next 3 wks;	Placebo gum, same schedule as above	Designed to simulate OTC setting	6m (biochem, car)	Unclear

						1 every 4-8hrs for final 3 wks				
Shiffman 2009 (4mg)⁽⁵⁶⁾	USA	Community volunteers	1661	46	50%	Nicotine gum 4mg. Instructed to gradually reduce smoking while increasing gum use for up to 8 wks. Postquit instructed to use 1 piece every 1-2hrs for first 6 wks; 1 every 2-4hrs for next 3 wks; 1 every 4-8hrs for final 3 wks	Placebo gum, same schedule as above	Designed to simulate OTC setting	6m (biochem, car)	Unclear
Tonnesen 1988⁽⁵⁷⁾	Denmark	Primary care	113	45	56%/58%	Nicotine Gum (2 mg) for 16 wks	Placebo	Informal group support, 6 sessions	12m (biochem, car)	Unclear
Villa 1999⁽⁵⁸⁾	Spain	Worksite volunteers	47	36	72%	Nicotine gum (2 mg)	No gum	8 weekly group sessions, 5 before TQD. Reduction prior to quitting	12m (self, NR)	Unclear
Zelman 1992⁽⁵⁹⁾	USA	Community volunteers	116	32	54%	Nicotine gum 2 mg, average 10 pieces/day, duration not stated + skills training; Nicotine gum + support counselling	Rapid smoking + support counselling; Rapid smoking + skills training	6x60-75 min group sessions over 2 wks, starting on quit day	12m (self, car)	High

Table 8.2 Included studies from systematic review: NRT patches versus placebo

Study	Country	Setting	Participants	Mean Age	Female	Intervention	Control	Support	Outcome	Risk of bias
Abelin 1989⁽⁶⁰⁾	Switzerland	Primary care	199	41	40%	Nicotine patch, 24hr, 12 wks with weaning; 21 mg smokers of >20 cpd, 14 mg for <20 cpd	Placebo patch	Unclear	12m (biochem, car)	High
Buchkremer 1988⁽⁶¹⁾	Germany	Community volunteers	131	35	50%	Nicotine patch (24hr/day, 8 wks, 15cm with weaning) + behavioural therapy	Placebo patch + behavioural therapy; Behavioural therapy alone	9 weekly group sessions	12m (self, NR)	Unclear
CEASE 1999⁽⁶²⁾	Europe	Community volunteers	3575	41	48%	Nicotine patch 15/25 mg (16hr), duration of active treatment 12/28 wks (incl 4 wk fading)	Placebo	Brief advice & self help brochure, visits at enrolment, TQD, wk 1, 2, 4, 8, 12, 22, 26	12m (biochem, car)	Low
Daughton 1991⁽⁶³⁾	USA	Community volunteers	158	42	53%	Nicotine patch (15cm ² , 4 wks) worn for 16 or 24hr/day	Placebo patch, 4 wks	Unclear & differed between sites	6m (self, car)	Unclear
Daughton 1998⁽⁶⁴⁾	USA	Community volunteers	158	42	53%	Nicotine patch (21 mg, 16hr, 10 wks with weaning)	Placebo patch	Nicoderm Committed Quitters Programme support booklet + follow-up visit 1 wk after quit day	12m (biochem, car)	Low

Davidson 1998⁽⁶⁵⁾	USA	Community volunteers	802	39	54%	Nicotine patch (22 mg, 24hr, for up to 6 wks)	Placebo patch	Self help book provided. Participants visited mall weekly to obtain patches. CO levels were monitored	6m (self, car)	Low
Ehrsam 1991⁽⁶⁶⁾	Switzerland	Primary care	112	26	NS	Nicotine patch (21 or 14 mg/24hr, 9 wks, tapered)	Placebo patch	High (no counselling)?	12m (biochem, car)	Unclear
Fiore 1994A⁽⁶⁷⁾	USA	Community volunteers	88	NS	NS	Nicotine patch (22 mg/24hr, 8 wks, no weaning)	Placebo patch	Intensive group counselling	6m (biochem, pp)	Low
Fiore 1994B⁽⁶⁷⁾	USA	Community volunteers	112	NS	NS	Nicotine patch (22 mg/24hr, 6 wks incl weaning)	Placebo patch	8 weekly 10-20 min individual counselling	6m (biochem, pp)	Unclear
Glavas 2003a⁽⁶⁸⁾	Croatia	NS	112	34	66%	Nicotine patch, 24hr, 25 mg/15 mg/8mg starting dose depending on baseline cpd. 3 wks	Placebo patch	Visits to pick up patch at 7, 14, 21 days, no details about advice given	12m (biochem, car)	Unclear
Glavas 2003b⁽⁶⁹⁾	Croatia	Community volunteers	160	NS	NS	Nicotine patch, 24hr, 25 mg/15 mg/8mg starting dose depending on baseline cpd. 6 wks; Nicotine patch, 24hr, 25 mg/15 mg starting dose depending on baseline cpd. 3 wks	Placebo patch. 6 wks; Placebo patch 3 wks	Low	6m (biochem, NR)	Unclear
Hays 1999⁽⁷⁰⁾	USA	Community volunteers	958	44	50%	Nicotine patches (22 mg, 24hr for 6 wks) purchased by participants, open	Placebo patches provided	No advice, counselling or interaction with medical	6m (biochem, pp)	Low

						label; Nicotine patches (22 mg, 24hr for 6 wks) provided, double blind		personnel		
Hughes 1999⁽⁷¹⁾	USA & Australia	Community volunteers	1039	43	50%	42 mg nicotine patch (24hr, 6 wks + 10 wks tapering); 35 mg nicotine patch; 21 mg nicotine patch	Placebo patch	Group behaviour therapy for 7 wks, brief individual counselling at 5 dose tapering meetings. Self help booklet	6m (biochem, car)	High
Hurt 1990⁽⁷²⁾	USA	Community volunteers	62	39	53%	Nicotine patch (30mg 24hrs, 6 wks + option of further 12 wks +/- tapering)	Placebo patch (continuing smokers at 6 wks were offered active patch)	Brief advice from nurse co-ordinator at 6 weekly visits	12m (biochem, car)	Unclear
Hurt 1994⁽⁷³⁾	USA	Community volunteers	240	43	53%	Nicotine patch (22 mg/24hr, 8 wks, no tapering)	Placebo patch	Nurse counselling at 8 weekly visits, weekly phone calls to wk 12	12m (biochem, car)	Unclear
ICRF 1994⁽⁷⁴⁾	UK	Primary care	1686	43	55%	Nicotine patch (21 mg/24hr, 12 wks incl tapering)	Placebo patch	Brief advice from nurse at 4 study visits	12m (biochem, car)	Low
Jorenby 1999⁽⁷⁵⁾	USA	Community volunteers	893	43	52%	Nicotine patch and placebo tablets	Placebo patch and placebo tablets	<15min individual counselling session at each weekly assessment.	12m (biochem, car)	Unclear

								One phone call 3 days after quit day		
Killen 1997⁽⁷⁶⁾	USA	Community volunteers	424	45	50%	Nicotine patch (21 mg/24hr) for 8 wks, 14 mg for 4 wks, 7mg for 4 wks; Nicotine patch and video	Placebo patch; Placebo patch and video	All treatment groups received a self help treatment manual designed to develop self regulatory skills	12m (biochem, car)	Unclear
Kornitzer 1995⁽⁷⁷⁾	Belgium	Worksite volunteers	374	40	39%	Nicotine patch (12 wks 15 mg/16hr, 6 wks 10mg, 6 wks 5 mg) and placebo gum	Placebo patch and placebo gum	Nurse counselling	12m (biochem, car)	Low
Oncken 2007⁽⁷⁸⁾	USA	Community volunteers	152	55	100%	Nicotine patch (21 mg for 13 wks incl 4 wks tapering)	Placebo patch	7 visits incl 4 x 2hr group counselling, 1 pre-TQD	16m (biochem, pp)	Unclear
Paoletti 1996⁽⁷⁹⁾	Italy	Community volunteers	297	43	40%	Nicotine patch (15 mg/16hr, 18 wks incl taper); Nicotine patch 25 mg	Placebo patch	Low	12m (biochem, pp)	Unclear
Piper 2009⁽⁸⁰⁾	USA	Community volunteers	1504	45	58%	Nicotine patch (24hr, 21, 14, and 7mg titrated down over 8 wk period postquit)	Placebo (5 groups matched to above 5 interventi ons)	7 one-to-one 10-20min counselling sessions	6m (biochem, pp)	Unclear
Richmond 1994⁽⁸¹⁾	Australia	Community volunteers	315	NS	NS	Nicotine patch (24 hr, 22 mg/24 hr, 10 wks incl tapering)	Placebo patch	Group therapy	12m (biochem, car)	Low
Sachs 1993⁽⁸²⁾	USA	Community volunteers	220	NS	NS	Nicotine patch (15 mg/16hr, 12 wks +	Placebo patch	Physician advice, 8 visits	12m (biochem,	Unclear

						6 wks tapering)		during treatment period	car)	
Sonderskov 1997⁽⁸³⁾	Denmark	Pharmacy customers	522	39	50%	Nicotine patch (24 hr). >20/day smokers used 21 mg for 4 wks, 14 mg for 4 wks, 7mg for 4 wks. Smokers of <20/day used 14 mg for first 8 wks, 7mg for 4 wks	Placebo patches	Brief instructions on patch use at baseline, visit to collect further patches at 4 & 8 wks, no behavioural support	6m (self, pp)	Low
Stapleton 1995⁽⁸⁴⁾	UK	Primary care	1200	NS	NS	Nicotine patch standard dose (15 mg/16hr for 18 wks); Nicotine patch with dose increase to 25 mg at 1 wk if required	Placebo patch group	Physician advice & brief support at 1, 3, 6, 12 wks	12m (biochem, car)	Low
TNSG 1991⁽⁸⁵⁾	USA	Community volunteers	808	43	60%	Nicotine patch (21 mg /24 hr, 6 wks+); Nicotine patch 14 mg	Placebo patch	Group therapy, 6+ sessions	6m (biochem, car)	Unclear
Tonnesen 1991⁽⁸⁶⁾	Denmark	Community volunteers	289	45	70%	Nicotine patch (15 mg/16hr for 12 wks with tapering)	Placebo patch	7 clinic visits including a few minutes of advice	12m (biochem, car)	Low
Westman 1993⁽⁸⁷⁾	USA	Community volunteers	158	41	57%	Nicotine patch (25 mg/24hr, 6 wks incl weaning)	Placebo patches	Brief counsellor support at 3 clinic visits, 4 telephone counselling sessions, self help materials	6m (biochem, car)	Unclear
Otero 2006⁽⁸⁸⁾	Brazil	Community volunteers	1199	42	63%	Nicotine patch (21 mg, 14 mg for FTND<5) 8 wks incl	Cognitive behavioural support	Mixed (Low=single 20 min session.	12m (self, pp)	High

						tapering + behavioural support	only	High= 1, 2, 3 or 4 weekly 1hr sessions. Maintenance or recycling sessions provided at 3, 6, 12m)		
Cinciripini 1996⁽⁸⁹⁾	USA	Community volunteers	64	NR	70%	Nicotine patch (21 mg, 12 wks incl weaning)	Behaviour therapy only (no placebo)	Group therapy weekly for 9 wks	12m (biochem, car)	Unclear
Prapavessis 2007⁽⁹⁰⁾	New Zealand	Community volunteers	121	NR	100%	Nicotine patch (21 mg/24hr for 10 wks, no weaning)	No patch	36x45 min session over 12 wks of group CBT or supervised vigorous exercise, starting 6 wks before TQD	12m (biochem, car)	Unclear
Wong 1999⁽⁹¹⁾	USA	Community volunteers	100	42	53%	Nicotine patch: 21 mg (24hr) for 8 wks, tapering to 14 mg for 4 wks	Naltrexone: 50mg/day for 12 wks	Individual counselling, 15-20 mins at 8 study visits	6m (biochem, car)	High
EAGLES 2016⁽⁹²⁾	International	Community volunteers	4028	46	66%	Patch, 21 mg x 7 weeks, 14 mg x 2 wks, 7 mg x 2 weeks (11 weeks)	Placebo	All participants received counselling (up to 10 mins) at all contacts. Participants were monitored at weeks 1 - 6, 8, 12, 13, 16, 20, 24; contacts	Abstinence at 6m (biochem, car)	Low

								were up to 15 face-to-face visits and 11 telephone visits		
--	--	--	--	--	--	--	--	---	--	--

Table 8.3 Included studies from systematic review: NRT inhaler versus placebo

Study	Country	Setting	Participants	Mean Age	Female	Intervention	Control	Support	Outcome	Risk of bias
Hjalmarson 1997⁽⁹³⁾	Sweden	Smoking cessation clinic	247	48	64%	Nicotine Inhaler (recommended minimum 4/day, tapering after 3m, use permitted to 6m)	Placebo inhaler	8 group meetings over 6 wks	12m (biochem, car)	Low
Leischow 1996⁽⁹⁴⁾	USA	Community volunteers	222	44	55%	Nicotine Inhaler (10mg). Advised to use 4-20 cartridges/day for 3m. After this tapering was encouraged until 6m	Placebo inhaler	Brief individual smoking cessation support at each study visit, 10 in all	12m (biochem, car)	Unclear
Schneider 1996⁽⁹⁵⁾	USA	Community volunteers	223	44	37%	Nicotine inhaler (4-20 inhalers per day) for up to 6m, with weaning from 3m	Placebo inhaler	Repeated clinic visits for assessment	12m (biochem, car)	Low
Tonnesen 1993⁽⁹⁶⁾	Denmark	Community volunteers	286	39	60%	Nicotine inhaler (2-10/day) up to 6m	Placebo inhaler	Brief advice at 8 clinic visits, 0, 1, 2, 3, 6,12, 24, 52 wks	12m (biochem, car)	Low

Table 8.4 Included studies from systematic review: NRT intranasal spray versus placebo

Study	Country	Setting	Participants	Mean Age	Female	Intervention	Control	Support	Outcome	Risk of bias
Blondal 1997⁽⁹⁷⁾	Iceland	Community volunteers	159	42	44%	Nicotine nasal spray (NNS) ad lib use. Each dose (2 squirts) delivered 1 mg nicotine. Maximum dose 5 mg/hr and 40 mg/day. Recommended duration of use 3m	Placebo nasal spray containing piperine to mimic sensory effect of nicotine.	Group therapy 6x1hr sessions	12m (biochem, car)	Unclear
Hjalmarson 1994⁽⁹⁸⁾	Sweden	Smoking cessation clinic	248	45	57%	Nicotine nasal spray (0.5 mg/spray) used as required up to 40mg/day for up to 1yr.	Placebo spray	8x45-60min group sessions over 6 wks with clinical psychologist	12m (biochem, car)	Unclear
Schneider 1995⁽⁹⁹⁾	USA	Community volunteers	255	NS	NS	Nicotine nasal spray (0.5 mg of nicotine per spray)	Placebo spray	Repeated clinic visits for assessment	12m (biochem, car)	Unclear
Sutherland 1992⁽¹⁰⁰⁾	UK	Smoking cessation clinic	227	NS	NS	Nicotine nasal spray, maximum 40mg/day	Placebo spray	4 wks group support	12m (biochem, car)	Unclear

Table 8.5 Included studies from systematic review: NRT lozenge or tablet versus placebo

Study	Country	Setting	Participants	Mean Age	Female	Intervention	Control	Support	Outcome	Risk of bias
-------	---------	---------	--------------	----------	--------	--------------	---------	---------	---------	--------------

Dautzenberg 2001 ⁽¹⁰¹⁾	France	Community volunteers	433	39	52%	Nicotine lozenge (1 mg, 8-24/day, 6 wks + 6 wks weaning for quitters)	Placebo lozenge	Not stated	6m (biochem, pp)	Unclear
Piper 2009 ⁽⁸⁰⁾	USA	Community volunteers	1504	45	58%	Nicotine lozenge 2 or 4 mg for 12 wks (based on dose-for-dependence level as per instructions)	Placebo; 5 groups matched to above 5 interventions	7 one-to-one 10-20min counselling sessions	6m (biochem, pp)	Unclear
Shiffman 2002 (2mg) ⁽¹⁰²⁾	USA & UK	Community volunteers	917	41	58%	Nicotine lozenge, 2 mg. Recommended dose 1 every 1-2hrs, min 9, max 20/day for 6 wks, decreasing 7-12 wks, available as needed 13-24 wks	Placebo lozenge, same schedule	Brief advice at 4 visits in 4 wks from enrolment	12m (biochem, car)	Unclear
Shiffman 2002 (4mg) ⁽¹⁰²⁾	USA & UK	Community volunteers	901	44	55%	Nicotine lozenge, 4 mg. Recommended dose 1 every 1-2hrs, min 9, max 20/day for 6 wks, decreasing 7-12 wks, available as needed 13-24 wks	Placebo lozenge, same schedule	Brief advice at 4 visits in 4 wks from enrolment	12m (biochem, car)	Unclear
Glover 2002 ⁽¹⁰³⁾	USA	Community volunteers	241	42	54%	Nicotine sublingual tablet (2 mg). Recommended dosage 1 tab/hr for smokers with FTND<7, 2 tabs/hr for scores 7. After 3mtreatment, tapering period of 3mif necessary	Placebo tablet	Brief counselling at all visits 1, 2, 3, 6 wks, 3, 6,12m	12m (biochem, car)	Low
Wallstrom 2000 ⁽¹⁰⁴⁾	Sweden	Community volunteers	247	45	59%	Nicotine sublingual tablet. Recommended dosage 1 tab/hr for smokers with FTND<7,	Placebo tablet	Brief 5 mins counselling at study visits (0, 1,	12m (biochem, car)	Unclear

						2 tabs/hr for scores 7. After 3m treatment, tapering period of 3m if necessary		2, 3, 6 wks, 3, 6m)		
--	--	--	--	--	--	--	--	---------------------	--	--

Table 8.6 Included studies from systematic review: NRT intranasal spray versus placebo

Study	Country	Setting	Participants	Mean Age	Female	Intervention	Control	Support	Outcome	Risk of bias
Tonnesen 2012⁽¹⁰⁵⁾	Germany/Denmark	Community volunteers	479	47	56%	1 mg/spray oral nicotine spray (in development, name not provided) Active: wks 1-6: 1-2 sprays when participants would normally have smoked a cigarette or experienced a craving, up to 4 sprays/hr and 64 sprays/day. Tapered down wks 7-12 (end of wk 9 instructed to be using half as much as in wks 1-6, reducing to max 4 sprays/ day by wk 12). Occasional use (max 4 sprays/day) permitted wks 13-24	Placebo on same schedule	General written and oral advice (less than 10min) at study start and less than 3mins at subsequent visits up to and including wk 24 (9 visits total)	12m (biochem, car)	Low

Table 8.7 Included studies from systematic review: choice of NRT versus placebo

Study	Country	Setting	Participants	Mean Age	Female	Intervention	Control	Support	Outcome	Risk of bias
-------	---------	---------	--------------	----------	--------	--------------	---------	---------	---------	--------------

Kraliko va 2002 ^(106, 107)	Czech Republic	Community volunteers	314	46	58%	Choice of 4 mg nicotine gum (up to 24/day) or 10mg inhaler (6-12 daily) for up to 6m with further 3m tapering	Placebo gum or inhaler	Brief behavioural support at clinic visits (9 scheduled)	12m (biochem, car)	Unclear
Wittchen 2011 ⁽¹⁰⁸⁾	Germany	Primary care	467	43	52%	CBT + NRT for 9-12 wks, patient's choice of patch (7mg-52.5 mg), gum (2 or 4 mg) or spray (10mg/ml)	No NRT	CBT, 4-5 one on one counselling sessions for 20-30min	12m (self, NR)	High

Table 8.8 Included studies from systematic review: Patch plus lozenge versus placebo

Study	Country	Setting	Participants	Mean Age	Female	Intervention	Control	Support	Outcome	Risk of bias
Piper 2009 ⁽⁸⁰⁾	USA	Community volunteers	1504	45	58%	Lozenge + patch (duration and dosage as above)	Placebo; 5 groups matched to above 5 interventions	7 one-to-one 10-20min counselling sessions	6m (biochem, pp)	Unclear

Table 8.9 Included studies from systematic review: NRT plus bupropion versus placebo

Study	Country	Setting	Participants	Mean Age	Female	Intervention	Control	Support	Outcome	Risk of bias
Jorenby 1999 ⁽⁷⁵⁾	USA	Community volunteers	893	43	52%	Nicotine patch (21 mg/24hr for 6 wks, tapered for 2 wks) and sustained release bupropion 300mg for 9 wks from 1 wk before quit day	Placebo patch and placebo tablets	<15min individual counselling session at each weekly assessment. One phone call 3 days after	12m (biochem, car)	Unclear

								quit day		
Piper 2009⁽⁸⁰⁾	USA	Community volunteers	1504	45	58%	Bupropion + lozenge (duration and dosage as above)	Placebo; 5 groups matched to above 5 interventions	7 one-to-one 10-20min counselling sessions	6m (biochem, pp)	Unclear

Table 8.10 Included studies from systematic review: NRT versus bupropion

Study	Country	Setting	Participants	Mean Age	Female	Intervention	Control	Support	Outcome	Risk of bias
Patch										
Gariti 2009⁽¹⁰⁹⁾	USA	Community volunteers	260	54	57%	Patch (for participants >10 cpd: 21 mg/day for wks 1-4, 14 mg/day wks 5-6, 7mg/day wks 7-8; participants <10cpd: 14 mg/day for 6 wks, 7mg/day for wks 7-8) + 9 wks placebo bupropion + 10 wks individualized counselling sessions; Patch (dose as above) for 8 wks + 9 wks placebo bupropion + four 5-10min counselling sessions	Placebo patch for 8 wks + 9 wks bupropion SR + 10 wks individualized counselling sessions; Placebo patch for 8 wks + 9 wks bupropion SR + 4x5-10min counselling sessions	High level of support via individual counselling sessions	12m (biochem, pp)	Low
Jorenby 1999⁽⁷⁵⁾	USA	Community volunteers	893	43	52%	Nicotine patch and placebo tablets	Bupropion 300mg and placebo patch	<15min individual counselling session at each weekly assessment	12m (biochem, car)	Unclear

								nt. One phone call 3 days after quit day		
Piper 2009⁽⁸⁰⁾	USA	Community volunteers	1504	45	58%	Nicotine patch (24hr, 21, 14, and 7mg titrated down over 8 wk period postquit)	Bupropion SR (150mg bid, 1 wk pre-quit, 8 wks postquit)	7 one-to-one 10-20min counselling sessions	6m (biochem, pp)	Unclear
Smith 2009⁽¹¹⁰⁾	USA	Primary care	1346	44	56%	Nicotine patch only (21 mg post-quit wk 1-4; 14 mg wk 5-6; 7mg wk 7-8)	Bupropion only (up-titrated during wk pre-quit, 150mg bid for 8 wks postquit)	Behavioural support optional	6m (self, pp)	High
EAGLES 2016⁽⁹²⁾	International	Community volunteers	4028	46	66%	Nicotine patch, 21 mg x 7 weeks, 14 mg x 2 wks, 7 mg x 2 weeks (11 weeks)	Bupropion SR, 150 mg x 2/day (titrated for 3 days, then full dose for 11 weeks)	All participants received counselling (up to 10 mins) at all contacts. Participants were monitored at weeks 1 - 6, 8, 12, 13, 16, 20, 24; contacts were up to 15	6m (biochem, car)	Low

									face-to-face visits and 11 telephone visits		
Lozenge											
Piper 2009⁽⁸⁰⁾	USA	Community volunteers	1504	45	58%	Nicotine lozenge 2 or 4 mg for 12 wks (based on dose-for-dependence level as per instructions)	Bupropion SR (150mg bid, 1 wk pre-quit, 8 wks postquit)	7 one-to-one 10-20min counselling sessions	6m (biochem, pp)	Unclear	
Smith 2009⁽¹¹⁰⁾	USA	Primary care	1346	44	56%	Nicotine lozenge only (4 mg lozenge if first cig of day smoked >30 min after waking, 2 mg otherwise. 1 lozenge every 1-2hrs postquit wk 1-6; 1 lozenge every 2-4hrs wk 7-9; 1 lozenge every 4-8hrs wk 10-12)	Bupropion only (up-titrated during wk pre-quit, 150mg bid for 8 wks postquit)	Behavioral support optional	6m (self, pp)	High	
Choice of NRT											
Wittchen 2011⁽¹⁰⁸⁾	Germany	Primary care	467	43	52%	CBT + NRT for 9-12 wks, patient's choice of patch (7mg-52.5 mg), gum (2 or 4 mg) or spray (10mg/ml)	CBT + bupropion SR (9-12 wks, 150mg;1/d for first 6d; 2/d thereafter)	4-5 one on one counselling sessions for 20-30min	12m (self, NR)	High	

Table 8.11 Included studies from systematic review: NRT versus NRT

Study	Country	Setting	Participants	Mean Age	Female	Intervention	Control	Support	Outcome	Risk of bias
-------	---------	---------	--------------	----------	--------	--------------	---------	---------	---------	--------------

Komitzer 1995⁽⁷⁷⁾	Belgium	Worksite volunteers	374	40	39%	Patch+gum	Patch	Nurse counselling	12m (biochem, car)	Low
Puska 1995⁽¹¹¹⁾	Finland	Community volunteers	300	20-65	NS	Patch+gum	Gum	Advice from nurses	12m (biochem, car)	Unclear
Blondal 1999⁽¹¹²⁾	Iceland	Community volunteers	237	42	67%	Nasal spray+patch	Patch	4 supportive group meetings	12m (biochem, car)	Low
Croghan 2003⁽¹¹³⁾	USA	Community volunteers	1384	42	58%	Nasal spray+patch	Spray or patch	Advice at each visit, 30-45 mins total	6m (biochem, pp)	High
Bohadana 2000⁽¹¹⁴⁾	France	Community volunteers	400	NS	51%	Patch+inhaler	Inhaler	Brief counselling and support from investigator at each visit	12m (biochem, car)	Unclear
Piper 2009⁽⁸⁰⁾	USA	Community volunteers	1504	45	58%	Patch+lozenge	Patch or lozenge	7 one-to-one 10-20min counselling sessions	6m (biochem, pp)	Unclear
Smith 2009⁽¹¹⁰⁾	USA	Primary care	1346	44	56%	Patch+lozenge	Patch or lozenge	Behavioural support optional	6m (self, pp)	High
Croghan 2003⁽¹¹³⁾	USA	Community volunteers	1384	42	58%	Nasal spray	Patch	Advice at each visit, 30-45 mins total	6m (biochem, pp)	High
Lerman 2004⁽¹¹⁵⁾	USA	Community volunteers	350	46	54%	Nasal spray	Patch	7x90 min behavioural group counselling sessions. TQD in wk 3	6m (biochem, pp)	Low
Piper 2009⁽⁸⁰⁾	USA	Community volunteers	1504	45	58%	Lozenge	Patch	7 one-to-one 10-20min counselling sessions	6m (biochem, pp)	Unclear

Schnoll 2010b ⁽¹¹⁶⁾	USA	Community volunteers	642	45	57%	Lozenge	Patch	5 individual counselling sessions	6m (biochem, car)	High
Smith 2009 ⁽¹¹⁰⁾	USA	Primary care	1346	44	56%	Lozenge	Ppatch	Behavioural support optional	6m (self, pp)	High

Table 8.12 Included studies from systematic review: e-cigarette versus placebo

Study	Setting	Mean Age	Ethnicity	Female	Participants	Intervention	Control	Behavioural Support	Outcome	Risk of bias
Bullen 2013 ⁽¹¹⁷⁾	New Zealand	42	33% Maori	62%	657	E-cig 16mg for 13 weeks	Placebo	Referred to Quitline and received an invitation to access phone- or text-based support.	Abstinence at 6m (biochem, car)	Low
Caponnetto 2013 ⁽¹¹⁸⁾	Italy	44	NS	36%	300	E-cig 7.2mg for 12 wks or 7.2 for 6 wks + 5.4mg for 6 wks	Placebo	Baseline visit and up to 7 follow-up visits to receive more cartridges, hand in diaries, measure CO and vital signs	Abstinence at 12m (self, car)	Low

Table 8.13 Included studies from systematic review: e-cigarette versus nicotine patch

Study	Setting	Mean Age	Ethnicity	Female	Participants	Intervention	Control	Behavioural Support	Outcome	Risk of bias
Bullen 2013 ⁽¹¹⁷⁾	New Zealand	42	33% Maori	62%	657	E-cig 16mg for 13 weeks	Patch 21 mg/24-hour, for 13 weeks	Referred to Quitline and received an invitation to access phone- or text-based support.	Abstinence at 6m (biochem, car)	Low

Table 8.14 Included studies from systematic review: bupropion versus placebo

Study	Setting	Mean Age	Ethnicity	Female	Participants	Intervention	Control	Behavioural Support	Outcome	Risk of bias
Ahluwalia 2002⁽¹¹⁹⁾	USA	44	African American	70%	600	Bupropion 300mg for 7 weeks	Placebo	8 sessions of in-person or telephone counselling & self help guide	Abstinence at 6m (biochem, car)	Low
Aubin 2004⁽¹²⁰⁾	France	41	NS	56%	504	Bupropion 300mg for 7 weeks	Placebo	Motivational support at clinic visits at baseline, w3, w7, w12 & 3 phone calls TQD, 2-3 days later, w5, w18	Abstinence at 6m (biochem, car)	Low
Brown 2007⁽¹²¹⁾	USA	44	NS	48%	524	Bupropion 300 mg/day for 12 weeks	Placebo	12 x 90 min groups twice weekly/ weekly/ monthly for 12w	Abstinence at 12m (biochem, car)	Unclear
Cinciripini 2013⁽¹²²⁾	USA	44	NS	39%	294	Bupropion 150mg/d days 1-3, 300mg/d thereafter for 12 weeks	Placebo	10 individual counselling sessions (6 in person, 4 via phone, 240 mins total)	Abstinence at 6m (biochem, car)	Unclear
Collins 2004⁽¹²³⁾	USA	46	NS	57%	555	Bupropion 300mg/day for 10 weeks begun 2 weeks before TQD	Placebo	7 sessions of group behavioural therapy	Abstinence at 6m (biochem, car)	Low
Cox 2012⁽¹²⁴⁾	USA	47	African American	66%	540	Bupropion 300mg for 7 weeks	Placebo	Up to 6 one-to-one 15-20 minute individual counselling sessions, self-help guide at start	Abstinence at 6m (biochem, pp)	Unclear
Dalsgard 2004⁽¹²⁵⁾	Denmark	43	NS	75%	335	Bupropion 300mg for 7 weeks	Placebo	Motivational support around TQD, at w3 & 7, and at 12w follow up	Abstinence at 6m (biochem, car)	Low
Fossati	Italy	49	NS	40%	593	Bupropion	Placebo	GP visits at enrolment & 4, 7,	Abstinence	Unclear

2007⁽¹²⁶⁾						300mg for 7 weeks		26 & 52w, phone calls 1 day pre TQD, 3 days post TQD, 10w post enrolment. Classified as low intensity	at 12m (biochem, car)	
Gonzales 2001⁽¹²⁷⁾	USA	45	NS	52%	450	Bupropion 300mg for 12 weeks, begun 7 days pre-TQD	Placebo	Brief individual counselling at visits w1-7, 9, 12, + telephone counselling at 4 and 5m	Abstinence at 12m (biochem, car)	Unclear
Gonzales 2006⁽¹²⁸⁾	USA	42	NS	46%	1025	Bupropion 300mg for 12 weeks, begun 7 days pre-TQD	Placebo	Brief (<10 min) standardized individual counselling at 12 weekly visits during drug phase and 11 clinic/phone visits during follow up, problem solving and relapse prevention	Abstinence at 12m (biochem, car)	Low
Haggstrom 2006⁽¹²⁹⁾	Brazil	44	NS	59%	156	Bupropion 300mg for 60 days, TQD during week 2	Placebo	6x 15-min individual CBT, weekly then bi-weekly	Abstinence at 6m (biochem, car)	Unclear
Hall 2002⁽¹³⁰⁾	USA	40	NS	44%	220	Bupropion 300mg/day for 12 weeks	Placebo	Medical management or psychological intervention	Abstinence at 12m (biochem, car)	High
Holt 2005⁽¹³¹⁾	New Zealand	40	Maori	72%	134	Bupropion 300mg for 7 weeks	Placebo	Counselling at 3 clinic visits during medication & 3 monthly follow ups, motivational phone call 1 day before & 2 days after TQD	Abstinence at 12m (biochem, car)	High
Hurt 1997⁽¹³²⁾	USA	44	NS	55%	615	Bupropion 300mg for 7 weeks	Placebo	Physician advice, S-H materials, and brief individual counselling by study assistant at each visit	Abstinence at 12m (biochem, car)	Unclear
Jorenby 1999⁽⁷⁵⁾	USA	43	NS	52%	893	Bupropion 300 mg for 9w from 1w	Placebo	Brief (< 15 min) individual counselling session at each weekly assessment. One	Abstinence at 12m (biochem,	Unclear

						before quit day		telephone call 3 days after quit day	car)	
Jorenby 2006 ⁽¹³³⁾	USA	42	NS	41%	1027	Bupropion 300mg for 12 weeks	Placebo	Brief (< 10 min) individual counselling at each weekly assessment for 12w & 5 follow-up visits. One telephone call 3 days after quit day	Abstinence at 12m (biochem, car)	Low
Levine 2010 ⁽¹³⁴⁾	USA	42	NS	100%	349	Bupropion SR for 26 weeks. 150mg/d for first 2 days and 300mg/d for remainder of treatment.	Placebo	12, 90 minute group counselling sessions delivered over 3 months	Abstinence at 12m (biochem, car)	High
McCarthy 2008 ⁽¹³⁵⁾	USA	39	NS	50%	463	Bupropion SR 300mg for 8 weeks	Placebo	Counselling; 8 x10min session, 2 prequit, TQD, 5 over 4 wks OR psychoeducation about medication, support & encouragement. Same no. of sessions, 80mins less contact time	Abstinence at 12m (biochem, pp)	Low
Nides 2006 ⁽¹³⁶⁾	USA	41	NS	51%	638	Bupropion 300mg for 7 weeks	Placebo	Up to 10 mins counselling at 7 weekly clinic visits, 12 & 24w	Abstinence at 12m (biochem, car)	Low
Piper 2007 ⁽¹³⁷⁾	USA	42	NS	58%	608	Bupropion 300mg for 9 weeks, starting 1wk before TQD	Placebo	3x 10 min counselling over 3 weeks	Abstinence at 12m (biochem, pp)	Unclear
Piper 2009 ⁽⁸⁰⁾	USA	45	NS	58%	1504	Bupropion SR (150 mg bid, 1 wk pre-quit, 8 wks postquit)	Placebo	7 one-to-one 10-20min counselling sessions	Abstinence at 6m (biochem, pp)	Unclear

Rovina 2009 ⁽¹³⁸⁾	Greece	45	NS	40%	205	Bupriopion 300mg for 19 weeks	No treatment	Cognitive behavioral group therapy (CBGT), 1 hour weekly for 1 m, then every 3 wks until 19 wks	Abstinence at 12m (biochem, car)	High
Schmitz 2007 ⁽¹³⁹⁾	USA	48	NS	100%	154	Bupropion 300mg for 7 weeks	Placebo	Either CBT based on relapse prevention model, or group support therapy, both 7 weekly 60 min meetings, TQD morning of 1st session, 10 days after start of medications	Abstinence at 12m (biochem, pp)	Low
Selby 2003 ⁽¹⁴⁰⁾	Canada	NR	NS	NS	284	Bupropion 300mg for 12 weeks	Placebo	Not described	Abstinence at 12m (biochem, pp)	Unclear
SMK2000 1 ⁽¹⁴¹⁾	USA	42	NS	48%	286	Bupropion 300mg for 7 weeks	Placebo	Not described	Abstinence at 12m (biochem, car)	Unclear
Tonnesen 2003 ⁽¹⁴²⁾	Europe, Australia, New Zealand	42	NS	51%	710	Bupropion SR 300mg for 7 weeks	Placebo	Brief motivational support at weekly clinic visits and telephone support during follow up. 11 clinic visits and 10 phone calls scheduled.	Abstinence at 12m (biochem, car)	Low
Uyar 2007 ⁽¹⁴³⁾	Turkey	36	NS	19%	131	Bupropion 300mg for 7 weeks	No treatment	Brief counselling on consequences of smoking with follow up for 24 weeks-more than low intensity	Abstinence at 6m (biochem, pp)	High
Zellweger 2005 ⁽¹⁴⁴⁾	Europe	40	NS	64%	667	Bupropion SR 300mg for 7 weeks	Placebo	Brief (10-15 min) motivational support at weekly clinic visits and telephone support one day before TQD, 3 days after TQD, monthly during follow up	Abstinence at 12m (biochem, car)	Unclear
Wittchen	Germany	43	NS	48%	467	Bupriopion	No	CBT of 4-5 1on1 counselling	Abstinence	High

2011⁽¹⁰⁸⁾						300mg for 12 weeks	treatment	sessions for 20-30mins	at 12m (self, car)	
EAGLES 2016⁽⁹²⁾	International	46	NS	66%	4028	Bupropion SR, 150 mg x 2/day (titrated for 3 days, then full dose for 11 weeks)	Varenicline 2mg/day for 12 weeks	All participants received counselling (up to 10 mins) at all contacts. Participants were monitored at weeks 1 - 6, 8, 12, 13, 16, 20, 24; contacts were up to 15 face-to-face visits and 11 telephone visits	Abstinence at 6m (biochem, car)	Low

Table 8.15 Included studies from systematic review: bupropion versus varenicline

Study	Setting	Mean Age	Ethnicity	Female	Participants	Bupropion	Varenicline	Behavioural Support	Outcome	Risk of bias
Cinciripini 2013⁽¹²²⁾	USA	44	NS	39%	294	Bupropion 150mg/d days 1-3, 300mg/d thereafter for 12 weeks	Varenicline 0.5mg days 1-3, 1mg days 4-7, 2mg thereafter for 12wks	10 individual counselling sessions (6 in person, 4 via phone, 240 mins total)	Abstinence at 6m (biochem, car)	Unclear
Gonzales 2006⁽¹²⁸⁾	USA	42	NS	46%	1025	Bupropion 300mg for 12 weeks, begun 7 days pre-TQD	Varenicline 2mg/day	Brief (<10 min) standardized individual counselling at 12 weekly visits during drug phase and 11 clinic/phone visits during follow up, problem solving and relapse prevention	Abstinence at 12m (biochem, car)	Low
Jorenby 2006⁽¹³³⁾	USA	42	NS	41%	1027	Bupropion 300mg for 12 weeks	Varenicline 2mg for 12 weeks	Brief (< 10 min) individual counselling at each weekly assessment for 12w & 5	Abstinence at 12m (biochem, car)	Low

								follow-up visits. One telephone call 3 days after quit day	car)	
Nides 2006⁽¹³⁶⁾	USA	41	NS	51%	638	Bupropion 300mg for 7 weeks	Varenicline 2mg for 7 weeks	Up to 10 mins counselling at 7 weekly clinic visits, 12 & 24w	Abstinence at 12m (biochem, car)	Low
EAGLES 2016⁽⁹²⁾	International	46	NS	66%	4028	Bupropion SR, 150 mg x 2/day (titrated for 3 days, then full dose for 11 weeks)	Varenicline 2mg/day for 12 weeks	All participants received counselling (up to 10 mins) at all contacts. Participants were monitored at weeks 1 - 6, 8, 12, 13, 16, 20, 24; contacts were up to 15 face-to-face visits and 11 telephone visits	Abstinence at 6m (biochem, car)	Low

Table 8.16 Included studies from systematic review: bupropion versus nicotine replacement therapy (NRT)

Study	Setting	Mean Age	Ethnicity	Female	Participants	Bupropion	NRT	Behavioural Support	Outcome	Risk of bias
NRT Patches										
Gariti 2009⁽¹⁰⁹⁾	USA	54	NS	57%	260	Bupropion SR for 9 weeks	Nicotine patch for 8 weeks	10 weeks of individual counselling sessions OR Four 5-10 minute counselling sessions	Abstinence at 12m (biochem, pp)	Low
Jorenby 1999⁽⁷⁵⁾	USA	43	NS	52%	893	Bupropion 300 mg for 9w from 1w before quit day	Patch 24 hr, 21mg for 6 weeks, tapered for 2 weeks	Brief (< 15 min) individual counselling session at each weekly assessment. One telephone call 3 days after quit day	Abstinence at 12m (biochem, car)	Unclear
Piper 2009⁽⁸⁰⁾	USA	45	NS	58%	1504	Bupropion SR (150 mg bid, 1 wk pre-quit, 8	Patch 24 hr, 21, 14 and 7mg titrated down over 8	7 one-to-one 10-20min counselling sessions	Abstinence at 6m (biochem, pp)	Unclear

						wks postquit)	weeks			
Smith 2009⁽¹¹⁰⁾	USA	44	NS	56%	1346	Bupropion SR (150 mg bid, 1 wk pre-quit, 8 wks postquit)	Patch, 21, 14 and 7mg titrated down over 8 weeks	Quitline counselling - up to 4 calls + additional support if required	Abstinence at 6m (self, pp)	High
Uyar 2007⁽¹⁴³⁾	Turkey	36	NS	19%	131	Bupropion 300mg for 7 weeks	Patch 21mg for 6 weeks	Brief counselling on consequences of smoking with follow up for 24 weeks- more than low intensity	Abstinence at 6m (biochem, pp)	High
EAGLES 2016⁽⁹²⁾	International	46	NS	66%	4028	Bupropion SR, 150 mg x 2/day (titrated for 3 days, then full dose for 11 weeks)	Nicotine patch, 21 mg x 7 weeks, 14 mg x 2 wks, 7 mg x 2 weeks (11 weeks)	All participants received counselling (up to 10 mins) at all contacts. Participants were monitored at weeks 1 - 6, 8, 12, 13, 16, 20, 24; contacts were up to 15 face-to-face visits and 11 telephone visits	Abstinence at 6m (biochem, car)	Low
NRT Lozenges										
Study	Setting	Mean Age	Ethnicity	Female	Participants	Bupropion Dose	NRT Dose	Behavioural Support	Outcome	Risk of bias
Piper 2009⁽⁸⁰⁾	USA	45	NS	58%	1504	Bupropion SR (150 mg bid, 1 wk pre-quit, 8 wks postquit)	Lozenge 2 or 4mg for 12 weeks	7 one-to-one 10-20min counselling sessions	Abstinence at 6m (biochem, pp)	Unclear
Smith 2009⁽¹¹⁰⁾	USA	44	NS	56%	1346	Bupropion SR (150 mg bid, 1 wk pre-quit, 8 wks postquit)	Lozenge 2 or 4mg for 12 weeks	Quitline counselling - up to 4 calls + additional support if required	Abstinence at 6m (self, pp)	High
NRT Patch plus Lozenge										

Study	Setting	Mean Age	Ethnicity	Female	Participants	Bupropion Dose	NRT Dose	Behavioural Support	Outcome	Risk of bias
Piper 2009 ⁽⁸⁰⁾	USA	45	NS	58%	1504	Bupropion SR (150 mg bid, 1 wk pre-quit, 8 wks postquit)	Lozenge 2 or 4mg for 12 weeks + 24 hr Patch, 21, 14 and 7mg titrated down over 8 weeks	7 one-to-one 10-20min counselling sessions	Abstinence at 6m (biochem, pp)	Unclear
Smith 2009 ⁽¹¹⁰⁾	USA	44	NS	56%	1346	Bupropion SR (150 mg bid, 1 wk pre-quit, 8 wks postquit)	Lozenge 2 or 4mg for 12 weeks +Patch, 21, 14 and 7mg titrated down over 8 weeks	Quitline counselling - up to 4 calls + additional support if required	Abstinence at 6m (self, pp)	High
Any NRT										
Study	Setting	Mean Age	Ethnicity	Female	Participants	Bupropion Dose	NRT Dose	Behavioural Support	Outcome	Risk of bias
Stapleton 2013 ⁽¹⁴⁵⁾	UK	41	NS	53%	1071	Bupropion 300mg for 8 weeks	Choice of patches, gums, lozenges, inhalator, nasal spray or microtab, for 12 weeks	7 weekly behavioural support session, mainly in groups, lasting 60-90 mins each	Abstinence at 6m (biochem, car)	High
Wittchen 2011 ⁽¹⁰⁸⁾	Germany	43	NS	48%	467	Bupropion 300mg for 12 weeks	Choice of patch (7-52.5mg), gum (2-	CBT of 4-5 1on1 counselling sessions for 20-30mins	Abstinence at 12m (self, car)	High

							4mg) or spray (10mg/ml)			
--	--	--	--	--	--	--	-------------------------	--	--	--

Table 8.17 Included studies from systematic review: bupropion versus bupropion plus nicotine replacement therapy (NRT)

GUM										
Study	Setting	Mean Age	Ethnicity	Female	Participants	Bupropion	Bupropion+ NRT	Behavioural Support	Outcome	Risk of bias
Piper 2007 ⁽¹³⁷⁾	USA	42	NS	58%	608	Bupropion 300mg for 9 weeks, starting 1wk before TQD	4mg nicotine gum, 300mg bupropion	3x 10 min counselling over 3 weeks	Abstinence at 12m (biochem, pp)	Unclear
LOZENGE										
Study	Setting	Mean Age	Ethnicity	Female	Participants	Bupropion	Bupropion+ NRT	Behavioural Support	Outcome	Risk of bias
Piper 2009 ⁽⁸⁰⁾	USA	45	NS	58%	1504	Bupropion SR (150 mg bid, 1 wk pre-quit, 8 wks postquit)	Bupropion SR (150 mg bid, 1 wk pre-quit, 8 wks postquit)+Lozenge 2 or 4mg for 12 weeks	7 one-to-one 10-20min counselling sessions	Abstinence at 6m (biochem, pp)	Unclear
Smith 2009 ⁽¹¹⁰⁾	USA	44	NS	56%	1346	Bupropion SR (150 mg bid, 1 wk pre-quit, 8 wks postquit)	Bupropion SR (150 mg bid, 1 wk pre-quit, 8 wks postquit)+Lozenge 2 or 4mg for 12 weeks	Quitline counselling - up to 4 calls + additional support if required	Abstinence at 6m (self, pp)	High
ANY NRT										

Study	Setting	Mean Age	Ethnicity	Female	Participants	Bupropion	Bupropion+NRT	Behavioural Support	Outcome	Risk of bias
Stapleton 2013⁽¹⁴⁵⁾	UK	41	NS	53%	1071	Bupropion 300mg for 8 weeks	Bupropion 300mg for 8 weeks+ choice of patches, gums, lozenges, inhalator, nasal spray or microtab, for 12 weeks	7 weekly behavioural support session, mainly in groups, lasting 60-90 mins each	Abstinence at 6m (biochem, car)	High
PATCH										
Study	Setting	Mean Age	Ethnicity	Female	Participants	Bupropion	Bupropion+NRT	Behavioural Support	Outcome	Risk of bias
Jorenby 1999⁽⁷⁵⁾	USA	43	NS	52%	893	Bupropion 300 mg for 9w from 1w before quit day	Bupropion 300 mg for 9w from 1w before quit day + Patch 24 hr, 21mg for 6 weeks, tapered for 2 weeks	Brief (< 15 min) individual counselling session at each weekly assessment. One telephone call 3 days after quit day	Abstinence at 12m (biochem, car)	Unclear

Table 8.19 Included studies from systematic review: bupropion 300mg versus bupropion 150mg

Study	Setting	Mean Age	Ethnicity	Female	Participants	Bupropion Dose	Bupropion Dose	Behavioural Support	Outcome	Risk of bias
Swan 2003⁽¹⁴⁶⁾	USA	45	NS	57%	1524	Bupropion 300mg	Bupropion 150mg	4 telephone calls+quitline+self-help material OR self-help+support line+1 telephone call	Abstinence at 12m (biochem, pp)	High
Hurt 1997⁽¹³²⁾	USA	44	NS	55%	615	Bupropion 300mg for 7 weeks	Bupropion 150mg for 7 weeks	Physician advice, S-H materials, and brief individual counselling by	Abstinence at 12m (biochem, car)	Unclear

								study assistant at each visit		
--	--	--	--	--	--	--	--	-------------------------------	--	--

Table 8.20 Included studies from systematic review: varenicline versus placebo

Study	Setting	Mean Age	Ethnicity	Female	Participants	Varenicline	Control	Behavioural Support	Outcome	Risk of bias
Cinciripini 2013⁽¹²²⁾	USA	44	66% White	39%	294	12 weeks varenicline 0.5mg/day days 1-3, 1.0mg/day days 4-7, 2.0mg/day thereafter	Placebo	10 individual counselling sessions (6 in person, 4 via phone, 240 mins total)	Abstinence at 6m (biochem, car)	Unclear
Gonzales 2006⁽¹²⁸⁾	USA	42	79% White	46%	1025	Varenicline 2mg/day for 12 weeks	Placebo	Brief (<10 min) standardized individual counselling at 12 weekly visits during drug phase and 11 clinic/phone visits during follow up, problem solving and relapse prevention	Abstinence at 12m (biochem, car)	Low
Jorenby 2006⁽¹³³⁾	USA	43	84% White	42%	1027	Varenicline 2mg/day for 12 weeks	Placebo	Brief (< 10 min) individual counselling at each weekly assessment for 12w & 5 follow-up visits. One telephone call 3 days after quit day	Abstinence at 12m (biochem, car)	Low
Nides 2006⁽¹³⁶⁾	USA	42	87% White	52%	638	Varenicline 2mg for 7 weeks	Placebo	Up to 10 mins counselling at 7 weekly clinic visits, 12 & 24w	Abstinence at 12m (biochem, car)	Low
Bolliger	Internatio	44	NS	34%	593	Varenicline	Placebo	All participants received	Abstinence at	Low

2011⁽¹⁴⁷⁾	nal					2mg/day for 12 weeks		You can quit smoking self-help booklet at baseline, and brief counselling (≤ 10 mins) at each clinic or telephone contact, plus a phone call 3 days post-TQD	6m (biochem, car)	
Nakamura 2007⁽¹⁴⁸⁾	Japan	40	NS	25%	619	Varenicline 2mg/day for 12 weeks	Placebo	All participants received a smoking cessation booklet Clearing the Air at baseline, + brief counselling (≤ 10 mins) at each clinic visit. Weekly visits throughout treatment phase, plus a 5 min phone call at TQD and +3 days post-TQD	Abstinence at 12m (biochem, car)	Low
Niaura 2008⁽¹⁴⁹⁾	USA	42	91% White	48%	320	Varenicline 0.5 to 2mg/day for 12 weeks	Placebo	All participants received a smoking cessation booklet Clearing the Air at baseline, + brief counselling (≤ 10 mins) at each clinic visit. Weekly visits throughout treatment phase.	Abstinence at 12m (biochem, car)	Low
Oncken 2006⁽¹⁵⁰⁾	USA	43	80% White	50%	647	Varenicline 2mg/day for 12 weeks	Placebo	All groups received self-help booklet at baseline, + brief (≤ 10 mins) counselling at weekly clinic visits throughout treatment phase, and phone call 3d post-TQD. At each visit smoking status reported and CO-verified; vital signs, weight and adverse	Abstinence at 12m (biochem, car)	Low

								events. Urine, blood tests and ECGs at screening, baseline, wks 1, 2, 4, 7 and 12		
Rennard 2012⁽¹⁵¹⁾	International	43	68% White	40%	659	Varenicline 2mg/day for 12 weeks	Placebo	All participants received Clearing the Air: Quit smoking today booklet at baseline, + brief counselling (≤ 10 mins) at each clinic visit. Weekly visits throughout treatment phase, and in follow-up phase clinic visits at wks 13, 16, 20 and 24. Phone calls at wks 14, 18 and 22	Abstinence at 6m (biochem, car)	Low
Tsai 2007⁽¹⁵²⁾	Taiwan and Korean	40	NS	11%	250	Varenicline 2mg/day for 12 weeks	Placebo	All participants received a smoking cessation booklet <i>Clearing the Air</i> at baseline, + brief counselling (≤ 10 mins) at each clinic visit. Clinic visits at baseline and at wks 1, 2, 3, 4, 6, 8, 10, 12, plus a 5 min phone call at +3 days post-TQD, and at wks 5, 7, 9, 11	Abstinence at 6m (biochem, car)	Low
Wang 2009⁽¹⁵³⁾	China, Singapore Thailand	39	NS	3%	333	Varenicline 2mg/day for 12 weeks	Placebo	All participants received a smoking cessation booklet at baseline, + brief counselling (≤ 10 mins) at each clinic visit, except for wks 5 and 7, when counselling was conducted by phone	Abstinence at 6m (biochem, car)	Unclear
DeDios	USA	42	Latino	53%	32	Varenicline	Placebo	30-minute face-to-face	Abstinence at	Unclear

2012⁽¹⁵⁴⁾						for 12 weeks		"culturally informed" smoking cessation behavioural intervention, + a non-tailored self-help brochure, all available in both English and Spanish	6m (biochem, pp)	
EAGLES 2016⁽⁹²⁾	International	46	NS	66%	4028	Varenicline 2mg/day for 12 weeks	Placebo	All participants received counselling (up to 10 mins) at all contacts. Participants were monitored at weeks 1 - 6, 8, 12, 13, 16, 20, 24; contacts were up to 15 face-to-face visits and 11 telephone visits	Abstinence at 6m (biochem, car)	Low
Gonzales 2014⁽¹⁵⁵⁾	International	48	93% White	50%	498	Varenicline 12 wks, titrated in 1st wk, 1 mg x 2/day	Placebo	Brief (< 10 mins) counselling at each contact. TQD set for wk 1 visit. Clinic visits at wks 1, 2, 3, 4, 6, 8, 9, 10, 11, 12; 13, 16, 24, 32, 40, 48, 52. Brief phone calls at wks 5, 7, 14, 20, 36, 44	Abstinence at 12m (biochem, car)	Low
Heydari 2012⁽¹⁵⁶⁾	Iran	43	NS	41%	272	Varenicline 2mg/day for 8 weeks	No treatment	All received brief (5 mins) education and counselling at 4 x weekly sessions	Abstinence at 12m (biochem, pp)	High
LONG TERM USAGE										
Study	Setting	Mean Age	Ethnicity	Female	Participants	Varenicline	Control	Behavioural Support	Outcome	Risk of bias
Williams 2007⁽¹⁵⁷⁾	USA, Australia	47	89% White	50%	377	Varenicline 2mg/day for 52 weeks	Placebo	Weekly visits throughout wks 1-8, then every 4 wks to wk 52, + wk 53 assessment.	Abstinence at 12m (biochem, pp)	Unclear

Ebbert 2015⁽¹⁵⁸⁾	International	44	NS	44%	1510	Varenicline for 24 weeks	Placebo	All participants asked to reduce their smoking rate by 50% by wk 4, by 75%+ by wk 8, and 100% by wk 12. Individual 10-minute counselling at each visit (18 face-to-face and 10 phone calls), + a self help book	Abstinence at 12m (biochem, car)	Low
NCT008 28113	USA	43	NS	43%	101	52-week varenicline therapy (2mg/day)	13 weeks of varenicline therapy (2mg/day)	Individual smoking cessation counselling. Brief (<10 minutes) smoking cessation counselling delivered at clinic visits	Abstinence at 12m (biochem, pp)	High

Table 8.21 Included studies from systematic review: varenicline versus NRT

Study	Setting	Mean Age	Ethnicity	Female	Participants	Varenicline Dose	Patch Dose	Behavioural Support	Outcome	Risk of bias
Aubin 2008⁽¹⁵⁹⁾	Europe & USA	43	93% White	51%	757	Varenicline 2mg for 12 weeks	Patch 21mg week 2-6, 14mg weeks 7-9, 7mg weeks 10-11	Self help booklet at baseline, and brief counselling (≤ 10 mins) at each clinic visit or by phone. TQD was at wk 1 visit. Weekly visits throughout treatment phase, plus a phone call 3 days post-TQD	Abstinence at 6 months (biochem, car)	High
Baker 2016⁽¹⁶⁰⁾	USA	48	67% White	52%	1086	Varenicline 2mg for 12 weeks	Patch 21mg week 1-8, 14mg weeks 9-10, 7mg	All participants received counselling (20 mins at visits 1, 2 and 3, and 10 mins by phone and at	Abstinence at 6 months (biochem, car)	High

							weeks 11-12 (11+ CPD) : Patch 14mg week 1-10, 7mg weeks 11-12 (5-10 CPD)	visits 4, 5) at 1 week pre-TQD and at TQD, wks 1, 4, 12 post-TQD, plus phone call at wk 8. In follow-up phase, participants were contacted at wks 26 and 52 by phone		
DeDios 2012⁽¹⁵⁴⁾	USA	42	Latino	53%	32	Varenicline for 12 weeks	Patch 12 wks: 4 wks @ 14 mg, 8 wks @ 7 mg	30-minute face-to-face "culturally informed" smoking cessation behavioural intervention, + a non-tailored self-help brochure, all available in both English and Spanish	Abstinence at 6m (biochem, pp)	Unclear
EAGLES 2016⁽⁹²⁾	International	46	NS	66%	4028	Varenicline 2mg/day for 12 weeks	Patch, 21 mg x 7 weeks, 14 mg x 2 wks, 7 mg x 2 weeks (11 weeks)	All participants received counselling (up to 10 mins) at all contacts. Participants were monitored at weeks 1 - 6, 8, 12, 13, 16, 20, 24; contacts were up to 15 face-to-face visits and 11 telephone visits	Abstinence at 6m (biochem, car)	Low
Heydari 2012⁽¹⁵⁶⁾	Iran	43	NS	41%	272	Varenicline 2mg/day for 8 weeks	Patch: 8 wks of 15 mg	All received brief (5 mins) education and counselling at 4 x weekly sessions.	Abstinence at 6m (biochem, pp)	High
Tsukahara 2010⁽¹⁶¹⁾	Japan	46	NS	25%	32	Varenicline 1.0 mg x 2/day for 12 wks, following 1 wk titration	Patch for 8 wks (52.5 mg/day for 4 wks, 35 mg/day for 2 wks, 17.5 mg/day for 2 wks)	Varenicline group received 8 clinic visits and nicotine group 5 visits over 12 wks, with 5 brief counselling sessions (\leq 10 mins)	Abstinence at 6m (self car)	High

NRT Patch and Lozenge

Study	Setting	Mean age	Ethnicity	Female	Participants	Varenicline	Patch + Lozenge	Behavioural Support	Outcome	Risk of bias
Baker 2016 ⁽¹⁶⁰⁾	USA	48	67% White	52%	1086	Varenicline 2mg for 12 weeks	Patch (21mg week 1-8, 14mg weeks 9-10, 7mg weeks 11-12 (11+ CPD) : Patch 14mg week 1-10, 7mg weeks 11-12 (5-10 CPD)) + Lozenge (2mg or 4mg 5 id for 12 weeks)	All participants received counselling (20 mins at visits 1, 2 and 3, and 10 mins by phone and at visits 4, 5) at 1 week pre-TQD and at TQD, wks 1, 4, 12 post-TQD, plus phone call at wk 8. In follow-up phase, participants were contacted at wks 26 and 52 by phone	Abstinence at 6 months (biochem, car)	High

Table 8.22 Included studies from systematic review: varenicline versus bupropion

Study	Setting	Mean Age	Ethnicity	Female	Participants	Varenicline	Bupropion	Behavioural Support	Outcome	Risk of bias
EAGLES 2016 ⁽⁹²⁾	International	46	NS	66%	4028	Varenicline 2mg/day for 12 weeks	Bupropion SR, 150 mg x 2/day (titrated for 3 days, then full dose for 11 weeks)	All participants received counselling (up to 10 mins) at all contacts. Participants were monitored at weeks 1 - 6, 8, 12, 13, 16, 20, 24; contacts were up to 15 face-to-face visits and 11 telephone visits	Abstinence at 6m (biochem, car)	Low
Gonzales 2006 ⁽¹²⁸⁾	USA	42	79% White	46%	1025	Varenicline 2mg/day for 12 weeks	Bupropion 150 mg x 2/day for 12 weeks	Brief (<10 min) standardized individual counselling at 12 weekly visits during drug phase	Abstinence at 12m (biochem, car)	Low

								and 11 clinic/phone visits during follow up, problem solving and relapse prevention		
Jorenby 2006⁽¹³³⁾	USA	42	NS	41%	1027	Varenicline 2mg for 12 weeks	Bupropion 300mg for 12 weeks	Brief (< 10 min) individual counselling at each weekly assessment for 12w & 5 follow-up visits. One telephone call 3 days after quit day	Abstinence at 12m (biochem, car)	Low
Nides 2006⁽¹³⁶⁾	USA	41	NS	51%	638	Varenicline 2mg for 7 weeks	Bupropion 300mg for 7 weeks	Up to 10 mins counselling at 7 weekly clinic visits, 12 & 24w	Abstinence at 12m (biochem, car)	Unclear
Cinciripini 2013⁽¹²²⁾	USA	44	66% White	39%	294	Varenicline 0.5mg days 1-3, 1mg days 4-7, 2mg thereafter for 12 weeks	Bupropion 150mg/d days 1-3, 300mg/d thereafter for 12 weeks	10 individual counselling sessions (6 in person, 4 via phone, 240 mins total)	Abstinence at 6m (biochem, car)	Unclear

Table 8.23 Included studies from systematic review: cytisine versus placebo

Study	Setting	Mean Age	Ethnicity	Female	Participants	Cytisine	Control	Behavioural Support	Outcome	Risk of bias
Scharfenberg 1971⁽¹⁶²⁾	East Germany	NS	NS	12%	1214	20-day course. 1.5mg tabs: Days 1-3 6/day; days 4-12 5/day; days 13-16 4/day; days 17-20 3/day	Placebo	None	Abstinence at 24m (self reported pp)	Unclear
Vinnikov 2008⁽¹⁶³⁾	Kyrgyzstan	39	NS	3%	197	25-day course. 1.5mg tabs: First 3	Placebo	Behaviour counselling	Abstinence at 6m (biochem)	Low

						days: 6 tabs per day; reduce smoking by half. Days 4-12: 5 tabs per day; stop smoking completely. Days 13-16: 4 tabs per day. Days 17-20: 3 tabs per day. Days 21-22: 2 tabs per day. Days 23-25: 1 tab per day			car)	
West 2011 ⁽¹⁶⁴⁾	Poland	48	NS	53%	740	25-day course. 1.5mg tabs: First 3 days: 6 tabs per day. Days 4-12: 5 tabs per day. Days 13-16: 4 tabs per day. Days 17-20: 3 tabs per day. Days 21-22: 2 tabs per day. Days 23-25: 1 tab per day	Placebo	Quitting advice, randomization and drugs dispensed at baseline visit; phone calls at TQD + 1 wk later (+ optional clinic visit). Clinic visit 4wks post-TQD, then phone calls at 6m and 12m, with visit to confirm abstinence if claimed. Behavioural support was minimal	Abstinence at 12m (biochem car)	Low

Table 8.24 Included studies from systematic review: cytisine versus NRT

Study	Setting	Mean Age	Ethnicity	Female	Participants	Cytisine	NRT	Behavioural Support	Outcome	Risk of bias
Walker 2014⁽¹⁶⁵⁾	New Zealand	38	Maori	57%	1310	Cytisine for 25 days + NRT vouchers if they needed them after 25 days	Choice of NRT (patch/gum/lozenge) for 8 weeks	Quitline support (3 10-15 minute calls over 8 weeks)	Abstinence at 6m (self reported, car)	High

Table 8.25 Included studies from systematic review: Acupuncture versus sham acupuncture/no treatment

Study	Setting	Participants	Mean Age	Ethnicity	Female	Intervention Dose	Control Dose	Other Support	Outcome	Risk of bias
Clavel 1992⁽¹⁶⁾	France	515	>18	NS	NS	Facial acupuncture to two points (GB8 and Bitong)	Sham acupuncture (wrong points 2cm from the active sites)	Both groups received placebo nicotine gum	Abstinence at 13m (self, car)	Unclear
Clavel 1992 +NG⁽¹⁶⁾	France	481	>18	NS	NS	Facial acupuncture to two points (GB8 and Bitong)	Sham acupuncture (wrong points 2cm from the active sites)	Both groups received active nicotine gum (2mg dose, up to 30 pieces/day, during first 6 months)	Abstinence at 13m (self, car)	Unclear
Cotraux 1983⁽¹⁶⁶⁾	France	558	18-50	NS	NS	Facial acupuncture, 3 weekly sessions	Waiting-list control	NR	Abstinence at 12m (self,	High

							(assessed at 12 months only)		car)	
Gillams 1984⁽¹⁶⁷⁾	UK	81	NS	NS	NS	Indwelling needle in active auricular point ('Lung') for 4 weeks	Indwelling needle in inactive auricular point (as far from 'Lung' as possible) for 4 weeks	NR	Abstinence at 6m (self, car)	Low
He 1997⁽¹⁶⁸⁾	Norway	46	NS	NS	NS	Combined body & auricular acup. Genuine points described for smoking cessation: Body: LU6, LU7; Ear: Shenmen, mouth, lung; Ear sustained acupressure: Shenmen, mouth, lung, trachea, hunger, endocrine	Sham points described for treating musculoskeletal conditions: Body: LI10, TE8; Ear: knees, lumbar vertebra, neck; Ear sustained acupressure: knees, lumbar vertebra, neck, shoulder, shoulder joint, buttock.	Both groups received combination of body electroacupuncture, ear acupuncture and ear acupressure. 6 treatments over 3 weeks	Abstinence at 8m (biochem, pp)	Low

Lamontagne 1980⁽¹⁶⁹⁾	Canada	75	20-50	NS	NS	Acupuncture to auricular points ('Zero' and 'Lung')	Self-monitor and report back	All subjects given 2 appointments 1 week apart. All smokers also given written advice on smoking cessation	Abstinence at 6m (self, pp)	Unclear
Leung 1991⁽¹⁷⁰⁾	Hong Kong	95	NS	NS, Asian	NS	Indwelling needles in auricular points ('Shenmen' and 'Lung') checked every 7 days; Introductory information sessions followed by 8 attendances in total, for supervision of needles	Waiting-list control	NR	Abstinence at 6m (self, pp)	High
Martin 1981a⁽¹⁷¹⁾	NZ	126	NS	NS	NS	Indwelling needles to 'effective' auricular points (lung and hunger) for 3 weeks	Indwelling needles to 'ineffective' auricular points (elbow and eye)	NR	Abstinence at 6m (self, pp)	Unclear
Martin 1981b⁽¹⁷¹⁾	NZ	126	NS	NS	NS	Combined body & auricular acup. indwelling needles to 'effective' auricular points (lung and hunger) for 3 weeks plus electroacupuncture for 20 minutes to LI4 in the hand and tongue point in the	Indwelling needles to 'ineffective' auricular points (elbow and eye) plus electroacupuncture for 20 minutes to LI4 in	NR	Abstinence at 6m (self, pp)	Unclear

						ear at the second attendance	the hand and tongue point in the ear at the second attendance			
Vandevienne 1985⁽¹⁷²⁾	France	200	NR	NS	NS	Combined body & auricular acup. acupuncture to 3 auricular and 2 body points	Sham acupuncture to nearby areas	both interventions given on days 1, 4, 10 and 20	Abstinence at 12m (self, pp)	Low
Waite 1998⁽¹⁷³⁾	UK	78	>18	NS	NS	Active group, lung point in ear	Control group, medial aspect of the patella, not on recognised acupuncture point	All smokers received structured counselling and written information before randomisation	Abstinence at 6m (biochem, pp)	Low
Wu 2007⁽¹⁷⁴⁾	Taiwan	131	≥18	NS, Asian	NS	'Real' points Shenmen, Lung, Mouth, Sympathetic (auricular and active states on pg 9)	'Irrelevant' points Eye, Elbow, Shoulder, Knee	NR	Abstinence at 6m (self, pp)	Low

Table 8.26 Included studies from systematic review: Acupuncture versus NRT

Study	Setting	Participants	Mean Age	Ethnicity	Female	Intervention Dose	NRT Dose	Other Support	Outcome	Risk of bias
Clavel 1992⁽¹⁶⁾	France	515	>18	NS	NS	Facial acupuncture to two points (GB8)	2mg nicotine gum	NR	Abstinence at 13m (self, car)	Unclear

						and Bitong)				
Clavel 1985⁽¹⁵⁾	France	651	NS	NS	NS	Facial acupuncture using 2 points bilaterally, single session	2mg nicotine gum	All groups also received 3 one-hour sessions of group therapy in first month	Abstinence at 13m (self, car)	Unclear

Table 8.27 Included studies from systematic review: Acupuncture versus counselling/psychological intervention

Study	Setting	Participants	Mean Age	Ethnicity	Female	Intervention	Control	Other Support	Outcome	Risk of bias
Cotraux 1983⁽¹⁶⁶⁾	France	558	18-50	NS	NS	Facial acupuncture, 3 weekly sessions	Behavioural therapy, 3 weekly sessions	NS	Abstinence at 12m (self, car)	High
Gillams 1984⁽¹⁶⁷⁾	UK	81	NS	NS	NS	Indwelling needle in active auricular point ('Lung') for 4 weeks	Group therapy sessions, 1 hr/wk for 4 weeks	NS	Abstinence at 6m (self, car)	Low
Leung 1991⁽¹⁷⁰⁾	Hong Kong	95	NS	Asian	NS	Indwelling needles in auricular points ('Shenmen' and 'Lung')	10 daily sessions of behavioural therapy lasting 1.5 hours	NS	Abstinence at 6m (self, pp)	High

Table 8.28 Included studies from systematic review: Group Behavioural Therapy (n=34)

Study	Setting	Participants	Age	Ethnicity	Female	Intervention Dose	Control Dose	Other Support	Outcome	Risk of bias
-------	---------	--------------	-----	-----------	--------	-------------------	--------------	---------------	---------	--------------

Bakkevig 2000 ⁽¹⁷⁵⁾	Norway	139	avg. 44	NS	67%	Group therapy; participants asked to attend 'Smokenders'. 7 weekly sessions + 1 follow up 4w later. Quit day after 5w. Multifaceted including cognitive therapies	Physician (GP) advice; participants instructed to visit their GP for support. GP told to offer NRT as appropriate and provide 1 follow-up visit		Abstinence 1 yr post-quit date Biochem	Unclear
Batra 1994 ⁽¹⁷⁶⁾	Germany	232	avg. 41	NS	53%	Group therapy, 9 weekly 90 min sessions	Self-help materials	Both conditions received nicotine patch	Continuous abstinence at 12m SR	Unclear
Camarellas 2002 ⁽¹⁷⁷⁾	Spain	106	avg. 47	NS	54%	Group therapy, 7 x2 hrs over 3w, TQD after w3	Individual counselling, not matched for intensity, 2 sessions over 2w, with S-H materials	72 participants eligible for nicotine patch, 53 used.	Sustained abstinence at 6m SR	Unclear
Cottraux 1983 ⁽¹⁶⁶⁾	France	558 (418 in arms of interest)	NS	NS	24%	Behaviour therapy. Includes discussion, training in relaxation. 3 x 3 hr sessions over 2 weeks. Relaxation and stress-desensitization audiotape for daily use	Placebo - lactose capsules for 2w. Met 2 x10min with a doctor		Abstinence at 12m SR	Unclear

Cottraux 1983⁽¹⁶⁶⁾	France	558 (418 in arms of interest)	NS	NS	24%	Behaviour therapy. Includes discussion, training in relaxation. 3 x 3 hr sessions over 2 weeks. Relaxation and stress-desensitization audiotape for daily use	1 yr waiting list control		Abstinence at 12m SR	Unclear
Curry 1988⁽¹⁷⁸⁾	USA	139	avg. 41	NS	51%	1. Relapse Prevention Group. Focused on smoking as learned behaviour. Quit day at 3rd session. Additional elements included identifying high risk situations, cognitive restructuring and role playing. AND 3. 'Absolute Abstinence' (AA)Group. Focused on addictive component of smoking. Quit day at 5th session. Additional elements included focused smoking, health education and contingency contract	2. Relapse Prevention Self help. 8 workbook units. AND 4. Absolute Abstinence Self help. 8 workbook units		Abstinence from months 9 to 12 of follow up. Biochem	Unclear

DePaul 1987⁽¹⁷⁹⁾	USA	233	avg. 43	NS	72%	Twice weekly 45 min group meetings for 3w	Self-help alone		Abstinence at 12m (multiple PP) SR	Unclear
DePaul 1989⁽¹⁸⁰⁾	USA	419	avg. 38	NS	63% in groups, 54% in S-H.	6 x twice-weekly group meetings to coincide with the 3w television series, then monthly meetings for 1 yr. Abstinent smokers and 5 of their family and 5 co-workers entered for a lottery at the final group meeting and 12m follow up	Self-help manuals only		Abstinence from end of programme to 12m Biochem	Unclear
DePaul 1994⁽¹⁸¹⁾	USA	564	avg. 38	NS	58%	6 group meetings over 3w followed by 14 booster meetings over 6m. Incentive payments. Handouts from same S-H manual. Maintenance manual (ALA A Lifetime of Freedom from Smoking)	S-H manual and incentive payment of US\$1 for each day abstinent up to US\$175	Worksite interventions were timed to coincide with a mass media intervention consisting of a week-long smoking cessation series on TV, and a complementary	Sustained abstinence at 12m Biochem	Unclear

								newspaper supplement		
Garcia 1989 ⁽²²⁾	Spain	68	avg. 34	NS	41%	Group therapy, 7 sessions over 3m	Individual counselling in clinic, same schedule as groups	Both received NG 2 mg	Sustained abstinence (quit at previous follow ups) at 6m Biochem	Unclear
Garcia 2000 ⁽¹⁸²⁾	Spain	162	avg. 32	NS	52%	1. Multicomponent programme, 10x 1hr sessions over 5w AND 2. Multicomponent, 5x 1hr over 5w AND 3. As 2 plus S-H manual	4. S-H manual, 1 orientation session		PP (7 day) abstinence at 12m Biochem	Unclear
Ginsberg 1992 ⁽¹⁸³⁾	USA	99	avg. 38	NS	54%	2. NG and behavioural programme including skill training, 5 sessions over 4w AND 3. NG and behavioural programme and partner support programme, 8 sessions over 5w	1. NG and educational materials, 2 sessions over 2w		Abstinence at 52w (not clear if abstinence required at prior assessment at weeks 4, 12, 26) 2&3v1 Biochem	Unclear

Glasgow 1981 ⁽¹⁸⁴⁾	USA	88 (85 included in analysis)	NS	NS	NS	Therapist administered programme based on either Danaher & Lichtenstein manual, Pomerleau & Pomerleau manual or I Quit Kit . 8 sessions over 8w	Self- administered using same 3 manuals		Abstinence at 6m Biochem	Unclear
Gruder 1993 ⁽¹⁸⁵⁾	USA	1440	NS	NS	NS	1. Social Support. 3 x90 min group meetings and copy of Quitters Guide for smokers, and 1 group meeting + Buddy Guide for buddies. Participants were instructed on how to get help from their buddies and others. Telephone calls to subjects and buddies at 1 and 2m AND 2. Discussion. Same schedule of meetings and phone calls as 1, but general information and review of self-help manual	3. No-contact control	All participants sent ALA Freedom from Smoking in 20 days manual and instructed to watch TV programme	Multiple PP abstinence (post- intervention , 6m and 12m). 24m rates also given but substantial loss to follow up by this time so 12m rates used here. SR	Unclear

Hill 1993⁽¹⁸⁶⁾	USA	82	50+	NS	NS	1. Behavioural Training (BT) adapted from Lung Health Study programme. Included quit date setting, RP training with role play of coping responses. 12 x 90 min session over 3m	2. Exercise and S-H pamphlet. This was a placebo control matched for contact time to 3. Therapist, who was blind to study hypothesis, encouraged smokers to quit at the exercise meetings		5 day abstinence at 12m. (Abstinence at previous follow ups not required) Biochem	Unclear
Hilleman 1993⁽¹⁸⁷⁾	USA	150	avg. ~50	NS	67%	1. Behaviour modification training, 12 x 1hr classes over 3m + transdermal clonidine 2. Same behaviour modification as 1, plus placebo patches	3. S-H printed material (I Quit Kit), transdermal clonidine AND 4. S-H printed material, placebo patches		Cessation at 1 yr SR	Unclear
Hollis 1993⁽¹⁸⁸⁾	USA	2707	NS	NS	NS	3. Group referral. Cessation advice, CO assessment. Video encouraged use of intensive (9 meetings over 2m) group programme, and waiver of fee. Effort made to schedule attendance	2. Self quit - Cessation advice, CO assessment, 10 min video, stop smoking kit, and choice of S-H manuals. Encouraged to set quit date. 1 follow-up telephone call and series of mailings	Smokers who received provider (physician, physician assistant or nurse practitioner) advice to quit	1 yr 2-PP abstinence (7 days at 3 and 12m) Biochem	High

<p>Hollis 1993⁽¹⁸⁸⁾</p>						<p>3. Group referral. Cessation advice, CO assessment. Video encouraged use of intensive (9 meetings over 2m) group programme, and waiver of fee. Effort made to schedule attendance</p>	<p>1. Advice - In addition to provider advice, given brief pamphlet by health counsellor</p>	<p>Smokers who received provider (physician, physician assistant or nurse practitioner) advice to quit</p>		
<p>Jorenby 1995⁽¹⁸⁹⁾</p>	<p>USA</p>	<p>504</p>	<p>avg. 44</p>	<p>NS</p>	<p>53%</p>	<p>Group: Given S-H pamphlet at screening visit along with motivational message. Received 8x 1hr weekly group sessions. Skills training, problem-solving skills</p>	<p>Individual: Given S-H pamphlet at screening visit along with motivational message. Also met nurse counsellor x3 following quit date. Nurse helped generate problem-solving strategies and provided praise and encouragement</p>	<p>Compared 22 mg vs 44 mg nicotine patch and 3 types of adjuvant treatment. Patch groups collapsed. All participants had 8 weekly assessments by research staff</p>	<p>7 day PP abstinence at 26w Biochem</p>	<p>Unclear</p>

Leung 1991⁽¹⁷⁰⁾	Hong Kong	95 (65 in relevant arms)	avg. ~37	NS, Asian	26%	Behavioural programme including self monitoring, management techniques, coping skills. 10 x 1½ hr sessions over 2w	Waiting list control		Abstinence (not defined) at 6m SR	Unclear
McDowell 1985⁽¹⁹⁰⁾	Canada	366	avg. 36	NS	60%	Operation Kick-It programme. 9 sessions. Therapists: public health nurse or health educator AND 3. Cognitive Behavior Modification programme. 9 sessions. Therapists: 1 of 2 M. Educational psychologists	Physician advice by one of 12 family physicians. 15 min counselling session with U.S. 'NCI Helping Smokers Quit Kit' and one postal follow up		Abstinence (over 1w diary period) at 12m SR	Unclear
McDowell 1985⁽¹⁹⁰⁾						Operation Kick-It programme. 9 sessions. Therapists: public health nurse or health educator AND 3. Cognitive Behavior Modification programme. 9 sessions. Therapists: 1 of 2 M. Educational psychologists	Self-monitoring control followed up at 2, 6 and 12m			

Minthorn-Biggs 2000⁽¹⁹¹⁾	Canada	75	avg. 41	NS	68%	1. Canadian Lung Association Countdown programme. 7 weekly sessions AND 2. Social interaction programme. 12 sessions over 6w + 4 weekly. Skills training	3. No-treatment control		Abstinence at 6m (12m rates only available for groups 1 and 2) SR	Unclear
Nevid 1997⁽¹⁹²⁾	USA	93	avg. 44	Hispanic	48%	Group therapy. 8 x 2 hrs. Included videos using culturally specific components. Motivation, nicotine fading, quitting techniques, RP, 'buddy' support. TQD 5th week	S-H with 1 group session for motivation and instructions and telephone contact. ALA Freedom from Smoking in 20 days in English & Spanish, also Guia para Dejar de Fumar	Both conditions received same maintenance programme ; ALA S-H manual A Lifetime of Freedom from Smoking and 2 telephone calls a month for 6m	Abstinence at 12m (sustained from post-treatment). PP rates also reported. (excludes 56 people, 35 Gr, 21 S-H who were randomized but did not attend any session and were not included in further analysis); Biochem valid.	Unclear

Omenn 1988⁽¹⁹³⁾	USA	159	avg. 43	NS	44%	<p>1. Multiple Component programme. 3 sessions over 3w. Didactic format AND 2. RP programme. 8 sessions over 8w. Interactive format, choice of immediate or phased quit</p>	<p>3. Minimal Treatment programme. S-H materials only. ACS 22-page Quitter's Guide 7-day plan</p>		<p>Abstinence at 12m (single PP) Biochem</p>	Unclear
Otero 2006⁽⁸⁸⁾	Brazil	1199 (includes 254 non-attenders)	avg. 42	NS	63%	<p>2. Cognitive behavioural, 1 or 2 weekly x1 hr sessions AND 3. As 2, with 3 or 4 weekly sessions</p>	<p>1. Single 20 min session - classified as brief intervention control in meta-analysis</p>	<p>No patch in comparison . 5 levels of behavioural support collapsed into 3 for analysis. Maintenance or recycling sessions provided to all groups at 3, 6, 12m</p>	<p>Abstinence at 12m (7 day PP) SR</p>	Low

<p>Otero 2006⁽⁸⁸⁾</p>	<p>Brazil</p>	<p>1199 (includes 254 non-attenders)</p>	<p>avg. 42</p>	<p>NS</p>	<p>63%</p>	<p>2. Cognitive behavioural, 1 or 2 weekly x1 hr sessions AND 3. As 2, with 3 or 4 weekly sessions</p>	<p>1. Single 20 min session - classified as brief intervention control in meta-analysis</p>	<p>Patch in comparison . Factorial design with NRT 21mg or 14mg patch for 8w incl tapering and 5 levels of behavioural support collapsed into 3 for analysis. Maintenance or recycling sessions provided to all groups at 3, 6, 12m</p>	<p>Abstinence at 12m (7 day PP) SR</p>	<p>Low</p>
<p>Pederson 1981⁽¹⁹⁴⁾</p>	<p>USA</p>	<p>40</p>	<p>avg. 39</p>	<p>NS</p>	<p>60%</p>	<p>1. Pomerleau & Pomerleau manual, an introductory session, followed by 1 hr group meetings at 2 and 6w AND 2. Danaher & Lichtenstein manual and same schedule of meetings as 1</p>	<p>3. Waiting list control</p>		<p>Abstinence at 6m for at least 3m SR</p>	<p>Unclear</p>

Pisinger 2005⁽¹⁹⁵⁾	Denmark	2408	avg. 46	NS	40%	1. 'Low intensity': single 15-45 min session of individual lifestyle counselling using motivational interviewing	2. 'High intensity': as 1 plus offer of participation in 6 session group course over 5m. Option to consider and be invited again in 3m		PP abstinence at 5 yrs (follow up at 1 & 3 yrs also) Biochem	Unclear
Rabkin 1984⁽¹⁹⁶⁾	Canada	168 (67 in relevant arms)	avg. 40	NS	NS	Behaviour modification. Multicomponent, 5 x 45-90 min meetings over 3w	Health Education. Single group meeting with didactic lectures by a health professional, film, discussion. Individual session with a therapist 1w later including a counselling element		Self-reported abstinence via questionnaire at 6m follow up SR	
Romand 2005⁽¹⁹⁷⁾	France	228	avg. 42	NS	54%	Five Day Plan (FDP); 5 sessions on consecutive nights, & supplementary sessions 1-2w later	Control; 1 hr of general information on tobacco-related health problems		Abstinence at 12m, lapse-free (PP also reported) Biochem	Unclear

Slovinc 2005⁽¹⁹⁸⁾	Canada	332	avg. 40	NS	100%	As control, plus Stress Management Training. 8 x 2 hr, 2 & 1w before TQD, 1,2,3,4,5,7w after. CBT targeted smoking-specific and life stressors	'Usual Care' 3 x 15 min physician visits, 2w before & 4 & 8w after TQD. Nicotine patch, S-H materials		Abstinence at 12m (7 day PP) Biochem	Low
Zheng 2007⁽¹⁹⁹⁾	China	232	avg. 56 (int), 53 (control)	NS, Asian	6%	Social cognitive group intervention, 5 x 2 hr twice weekly sessions	Waiting list control		Sustained abstinence at 6m Biochem	High

Table 8.29 Included studies from systematic review: Individual Behavioural Counselling

Study	Setting	Participants	Age	Ethnicity	Female	Intervention	Control	Other Support	Outcome	Risk of bias
Ahluwalia 2006⁽¹¹⁾	USA	755	avg. 45	African American	67%	Counselling using Motivational Interviewing (MI) approach. 3 in-person visits at randomization, wk1, wk8, and phone contact at wk3, wk6, wk16, S-H materials	Counselling using Health education (HE) approach. Same schedule & materials as intervention		PP abstinence at 6m (7 day PP) Biochem	Low
Alexandre 1998⁽²⁰⁰⁾	Spain	48	avg. 36	NS	65%	'Advanced', 4 x30 min over 4 wks, video, cognitive therapy, social influences, relapse prevention	'Minimal' 3 min advice immediately after randomization		Abstinence at 12m, SR	Unclear
Aveyard 2007⁽²⁰¹⁾	UK	925	ag. 43	NS	51%	Weekly support; as control, plus additional call at 10 days & visits at 14 & 21 days	Basic support; 1 visit (20-40 mins) before quit attempt, phone call on TQD, visits/ phone calls at 7-14 days & at 21-28 days (10-20 mins)	Both include 8 wks 16mg nicotine patch	Abstinence at 12m (sustained at 1, 4, 12, 26 wks) Biochem	Low

Bronson 1989⁽²⁰²⁾	USA	155	avg. 42	NS	62%	Two 20 min counselling sessions during a periodic health examination (benefits of quitting, assessment of motivation, quit plan, high risk/problem solving)	Control (completed smoking behaviour questionnaire)		Abstinence at 18m (sustained from 6-18m) Biochem	Unclear
Glasgow 2000⁽²⁰³⁾	USA	1154	avg. 24	NS	100%	Video (9 min) targeted at young women. 12-15 min counselling session, personalized strategies, stage-targeted S-H materials. Offered telephone support call	Generic S-H materials	Both groups received 20 sec provider advice	Abstinence at 6m (for 30 days) Biochem	Unclear
Jorenby 1995⁽¹⁸⁹⁾	USA	504	avg. 44	NS	NS	Individual - S-H at screening visit + motivational message. Met nurse counsellor x3 after TQD. Counsellor helped generate problem-solving strategies and provided praise and encouragement	Minimal - S-H materials from physician at screening visit for trial entry, instructed not to smoke whilst wearing patch. No further contact with counsellors		7 day PP abstinence at 26 wks Biochem	Unclear

Table 8.30 Included studies from systematic review: Brief Advice

Study	Setting	Participants	Age	Ethnicity	Female	Intervention	Control	Outcome	Biochem	Risk of bias
Betson 1997 ⁽²⁰⁴⁾	Hong Kong	865	>65	NS	8%	Arm 3. Physician advice (1min, based on 4As) AND Arm 4. Physician advice and booklet	Arm 1. No intervention AND Arm 2. Written materials (Chinese translation of American Cancer Society booklet)	Abstinence at 1 yr (sustained from 3m)	N	Unclear
Butler 1999 ⁽²⁰⁵⁾	UK	536	NS	NS	70%	Standardised brief advice (estimated time 2 minutes)	Structured motivational counselling (mean length 10mins) (based on stage of readiness to change)	PP at 6m (self-reported abstinence in the previous month)	N	Unclear
Demers 1990 ⁽²⁰⁶⁾	USA	519	NS	NS	NS	3 - 5 min smoking cessation counselling (and written materials) plus routine care.	Routine care	Sustained at 12m (& 6m)	N	
Fagerström 1984 ⁽¹⁹⁾	Sweden	145 (49 in relevant arms)	NS	NS	NS	Advice plus 2 appointments, phone call + letter (long follow-up)	Advice plus 1 appointment (short follow-up)	Sustained abstinence at 1, 6 and 12m	Y on random subset only	High

Haug 1994 ⁽²⁰⁷⁾	Norway	274	18-34	NS	100%	Advice + leaflet + invitation to attend 4 follow-up visits	Normal care controls	Abstinence at 18m	Y at 12m only not at 18m	High
Higashi 1995 ⁽²⁰⁸⁾	Japan	957	NS	NS	NS	Brief advice plus leaflet, encouragement card at 1m and telephone card (incentivised) at 6m	No intervention	PP abstinence at 12m	N	Unclear
Jamrozik 1984 ⁽³⁵⁾	UK	2110	NS	NS	NS	Brief advice to quit plus smoking cessation pamphlet	Normal care control group	PP abstinence at 12m	Y, random sample	High
Janz 1987 ⁽²⁰⁹⁾	USA	250	NS	NS	NS	Brief advice from physician and brief consultation from nurse	Normal care	PP abstinence at 6m	N	High
Lang 2000 ⁽²¹⁰⁾	France	1095	NS	NS	17%	Intensive intervention; contract with quit date, phone call 7 days post quit date, follow-up visit	Minimal advice; 5-10 min from occupational physician	Sustained abstinence (\geq 6m) at 12m, assessed at annual check up	Y for subsample	Low
Marshall 1985 ⁽²¹¹⁾	UK	200	NS	NS	NS	As control, plus offer of 4 follow-up visits over 3m	Advice plus nicotine gum	Sustained at 12m (from 6m)	Y	Low

McDowell 1985⁽¹⁹⁰⁾	Canada	366 (153 relevant to review)	NS	NS	NS	Brief physician advice	Control: self-monitoring of smoking	PP abstinence at 12m	N (but threatened to test them)	High
Meyer 2008⁽²¹²⁾	Germany	1499 (1011 in relevant conditions?)	18-70	NS	48%	Brief advice from trained physician and selected self-help manuals	Control group (assessment only - 22-sided questionnaire administered in waiting room)	Abstinence at 24m (sustained for 6m)	N	High
Morgan 1996⁽²¹³⁾	USA	659	50-74	NS	NS	Physician advice, stage-based, tailored self-help guide. Follow-up letter from physician and call from project staff. Smokers in contemplation given prescription and free 1 wk supply of gum	Usual care (delayed intervention)	Abstinence at 6m (assume PP)	N	Unclear
Nebot 1989⁽²¹⁴⁾	Spain	424	NS	NS	NS	Brief physician advice, 3 - 5 min, and self-help leaflet	Usual care	Abstinence at 12m (states definition unclear)	N incomplete	High

Ockene 1991⁽⁴⁷⁾	USA	1286	NS	NS	NS	Patient-centred counselling, written materials, asked to schedule follow-up visit, follow-up letter	Advice only	PP abstinence at 6m (self-reported)	N	Unclear
Page 1986⁽⁴⁸⁾	Canada	289	NS	NS	NS	Advice to quit	No advice	PP abstinence at 6m	N	High
Pieterse 2001⁽²¹⁵⁾	Netherlands	530	NS	NS	NS	Advice/counselling tailored to stage of change, self-help manual, follow-up visit if quit date set. Approx 10 mins	Usual care	Sustained at 12m (from 6m)	N	Unclear
Porter 1972⁽²¹⁶⁾	UK	191	NS	NS	NS	5 mins of advice delivered 'with conviction and vigour' plus antismoking leaflet	No advice	PP abstinence at 6m	N, validation: by family report/neighbour report	High
Richmond 1986⁽²¹⁷⁾	Australia	200	NS	NS	NS	Six visits to the GP over 6m, including advice, spirometry demonstration and serum cotinine and written materials	Control group completed questionnaire and gave blood sample at single visit	Sustained abstinence 3 yrs (assessed at 6m & 3 yrs)	Y but not on all, some confirmation by relatives and/or friends	High

Russell 1979⁽²¹⁸⁾	UK	2138	NS	NS	NS	Advice to stop smoking. AND 4. Advice to stop smoking plus leaflet plus a warning that the patient would be followed up	1. No intervention AND 2. Questionnaire only	Sustained abstinence at 12m (& 1m)	Y, sample only(n=23)	High
Russell 1983⁽⁵³⁾	UK	2106 (1377 in relevant arm)	NS	NS	NS	Advised to stop smoking plus provided with a 'give up smoking' booklet	No intervention.	Sustained abstinence 12m (& 4m)	Y, 66% of quitters biochem val	High
Segnan 1991⁽⁵⁵⁾	Italy	923	20-60	NS	NS	Repeated counselling (follow-up at 1,3,6,9m). OR Repeated counselling plus spirometry (conducted by specialist centre)	Advice and leaflet	Sustained abstinence at 12m (sustained for 3m by self report)	Y	Low
Severson 1997⁽²¹⁹⁾	USA	1478	New mothers	NS	100%	As control, and extended support (counselling plus follow-up at 2, 4, and 5m visits) and materials (incl video tape, written materials, signs, magnets, bib)	Information pack including a letter from paediatrician on risks of passive smoking, provided by birth hospital.	PP abstinence at 12m	N	Low
Slama 1990⁽²²⁰⁾	Australia	311	18-64, mean age NR	NS	NS	Advice plus leaflets (minimal intervention)	Control	Sustained abstinence at 12m (& 1m & 6m)	Y	Unclear

Thompson 1988 ⁽²²¹⁾	USA	1039	NR	NS	NS	A. Structured physician advice (3 - 5 min talk); C. Referral to group therapy	A. Structured physician advice (3 - 5 min talk)	PP abstinence at 9m	N	Unclear
Unrod 2007 ⁽²²²⁾	USA	518	NR	NS	NS	Physician & patient given 1-page tailored report based on computer-based assessment in waiting room. Physician trained to provide 5As-based brief counselling. Follow-up appointment could be arranged	No intervention (usual care)	PP abstinence at 6m	Y	High
Wilson 1982 ⁽²²³⁾	Canada	211	NR	NS	NS	Brief advice plus follow-up appointments at 1, 3 and 6m	Brief advice (5 min counselling)	PP abstinence at 6m	N	Unclear
Wilson 1990 ⁽²²⁴⁾	Australia	1238	NR	NS	NS	Personalised advice plus leaflets and visual aids at single visit	Normal care control group (advice given if clinically indicated)	Sustained abstinence at 12m (& 6m)	N	Low

Table 8.31 Included studies from systematic review: Nursing interventions

Study	Setting	Participants	Age	Ethnicity	Female	Intervention	Control	Adjunct	Outcome	Biochem valid	Risk of bias
-------	---------	--------------	-----	-----------	--------	--------------	---------	---------	---------	---------------	--------------

Aveyard 2003 ⁽²²⁵⁾	UK	831	NR	NS	NS	In addition to tailored self help, asked to make appointment to see practice nurse. Single postal reminder if no response. Up to 3 visits, at time of letters. Reinforced use of manual	Self-help manual based on Transtheoretical model, maximum of 3 letters generated by expert system. No face-to-face contact		Abstinence at 12m, self-reported sustained for 6m	Y	Low
Aveyard 2007 ⁽²⁰¹⁾	UK	925	avg. 43	NS	NS	Basic support; 1 visit (20 - 40 mins) before quit attempt, phone call on TQD, visits/phone calls at 7 - 14 days and at 21 - 28 days (10 - 20 mins)	Weekly support; as 1. plus additional call at 10 days and visits at 14 and 21 days	Both interventions included 8 wks 16 mg nicotine patch. Intensity high for both groups.	Abstinence at 12m (sustained at 1, 4, 12, 26 wks)	Y	Low
Borrelli 2005 ⁽²²⁶⁾	USA	278	avg. 57	NS	54%	Motivational enhancement. 3 x 20 - 30-min sessions during nursing visits. 5-min follow-up call	Standard care control based on 5Amodel, single 5 - 15-min session with brief support at subsequent nursing visits, consistent with		Abstinence at 12m (no smoking since 6m assessment)	Y but obtained for 60% only	Unclear

							guidelines				
Janz 1987 ⁽²⁰⁹⁾	USA	NR?	NS	NS	NS	Physician discussed personal susceptibility, self efficacy and concern, trained nurse counselled on problems and strategies. 2. As 1, and self-help manual 'Step-by-Step Quit Kit' + 1 telephone call	Usual Care control (from physicians not involved in study)		Abstinence at 6m (self report by telephone)	N	High
Lancaster 1999 ⁽²²⁷⁾	UK	497	NS	NS	NS	As control, plus invitation to contact a trained practice nurse for more intensive tailored counselling. Up to 5 follow-up visits offered	Physician advice (face-to-face or in a letter) and a leaflet	Physician advice (face-to-face or in a letter) and a leaflet to both groups	Abstinence at 12m (sustained at 3m and 12m)	Y	Low
Sanz-Pozo 2006 ⁽²²⁸⁾	Spain	125	avg. ~40	NS	52% (intervention) 62% (control)	1. Brief advice from doctor at recruitment, appointment with clinic nurse 7 days before TQD, on TQD, 1wk, 1m, 2m, 3m	2. Brief advice only		Sustained abstinence at 24m (from 12m)	Y	Unclear

Table 8.32 Included studies from systematic review: Internet interventions

Study	Setting	Mean Age	Ethnicity	Female	Participants	Intervention	Control	Adjunct	Outcome	Risk of bias
An 2008 ⁽²²⁹⁾	USA	20	NS	75%	517	Interactive, tailored	Non-active		30w Abstinence (biochem, pp)	Low
Brendryen 2008a ⁽²³⁰⁾	Norway	29.5	NS	50%	290	Interactive, tailored, phone contact	Self-help book		Abstinence at 6m (SR, PP), 12m (SR, car)	Low
Brendryen 2008b ⁽²³¹⁾	Norway	36	NS	50%	396	Interactive, tailored, phone contact	Self-help book	NRT (both groups)	Abstinence at 6m (SR, PP), 12m (SR, car)	Low
Elfeddali 2012 ^(232, 233)	Netherlands	42	NS	62%	2031	Interactive, tailored (2 different types, combined for analysis)	Usual care		Abstinence at 12m (SR, car)	High
McDonnell 2011 ⁽²³⁴⁾	USA	35	Korean-American	12%	1409	Interactive, not tailored	Booklet of same content		Abstinence at 6m (SR, car)	Low
Smit 2012 ⁽²³⁵⁾	Netherlands	49.5	NS	52%	1123	Interactive, tailored	No Intervention		Abstinence at 6m (SR, car)	High

Table 8.33 Included studies from systematic review: Motivational Interviewing versus routine care

Study	Setting	Mean Age	Ethnicity	Female	Participants	Intervention	Control	Adjunct	Outcome	Risk of bias
Bastian 2013 ⁽²³⁶⁾	USA	47	NS	58%	496	6 weekly MI telephone calls & control group	Self-directed Materials	NRT (patches)	Abstinence at 6m (SR).	Unclear

						Interventions		to both groups	PPA)	
Bock 2008 ⁽²³⁷⁾	USA	47.7	NS	47%	543	30 min MI counselling session, follow-up telephone calls	Usual care - referral sheet to SC resources	NRT (patches) to both groups	Abstinence at 6m (biochem. car)	Unclear
Bock 2014 ⁽²³⁸⁾	USA	39.6	NS	69%	846	45 min MI counselling, follow-up telephone call	Usual care - self-help guides	NRT (patches) to both groups	Abstinence at 6m, 12m (biochem (eCO2. PP)	Unclear
Borrelli 2005 ⁽²²⁶⁾	USA	57.2	NS	54%	278	3X20-30 min MI counselling by nurse	5-15min brief counselling (1 visit - nurse)	All received manual (clear horizons)	Abstinence at 6m, 12m (biochem (eCO2), car)	Unclear
Butler 1999 ⁽²⁰⁵⁾	UK	41	NS	71%	536	MI counselling (mean 10 mins) by GP	Brief advice (2 mins) by GP		Abstinence at 6m (SR. PPA)	Low
Davis 2011 ⁽²³⁹⁾	USA	37.6	NS	45%	218	MI counselling (nurse, mean 15min)	Prescriptive 15 min interview, nurse	No pharmacotherapy	Abstinence at 6m (SR. PPA)	Unclear
De Azevedo 2010 ⁽²⁴⁰⁾	Brazil	47	NS	36%	273	MI counselling (counsellor) 30min Follow up telephone call (X&, 10mins)	Low-intensity counselling (15min) by counsellor	No pharmacotherapy	Abstinence at 6m (SR. PPA)	High
Dornelas 2000 ⁽²⁴¹⁾	USA	54	NS	22%	100	MI counselling (psychologist, 20 mins, follow up phone calls)	Brief advice (10mins, psychologist)		Abstinence at 6m (SR. car)	High
Ellerbeck 2009 ⁽²³²⁾	USA	47.2	NS	59%	726	MI counselling calls (both Medium and High intensity groups), counsellors	Smoking cessation materials mailed	Both groups offered NRT +/- bupropion	Abstinence at 12m (biochem. PP)	Low
Glasgow 2000 ⁽²⁰³⁾	USA	24	NS	100%	1154	MI counselling (12-15min, counsellor)	Brief advice + brochure		Abstinence at 12m (biochem.	Low

									PP)	
Hennrikus 2005⁽²⁴²⁾	USA	47	NS	53%	2095	MI counselling (nurse, 20 mins) & usual care. Follow-up telephone calls	Usual care	NRT / Bupropion encouraged but not provided	Abstinence at 12m (biochem. PP)	Unclear
Hollis 2007⁽²⁴³⁾	USA	41	NS	60%	4614	MI counselling (40 mins MI counselling, follow-up call)	Usual care (brief counselling, 15 min call)	NRT in groups	Abstinence at 6m, 12m (SR. PPA)	Low
Lindqvist 2013⁽²⁴⁴⁾	Sweden	48.6	NS	81%	772	MI counselling	Standard Swedish usual care	Pharmacotherapy received both groups	Abstinence at 12m (SR. PP)	High
McClure 2005⁽²⁴⁵⁾	USA	33	NS	100%	275	MI counselling (15 min telephone calls, up to 4). Counsellor	Usual care	Pharmacotherapy to both groups	Abstinence at 12m (biochem. car)	Unclear
Rigotti 1997⁽²⁴⁶⁾	USA	48	NS	46%	650	MI counselling (15 mins) + brief advice	Usual care		Abstinence at 6m (biochem. PP)	High
Soria 2006⁽²⁴⁷⁾	Spain	38	NS	53%	200	MI counselling (3x20 mins, GP)	Brief advice (GP)	Bupropion offered to highly dependent - both groups	Abstinence at 6m, 12m (biochem. PPA)	Low
Tevyaw 2009⁽²⁴⁸⁾	USA	19.8	NS	NR	110	MET (with MI principles) x3 sessions (60min, 30min, 30min)	Muscle relaxation (Progressive Muscle Relaxation) to match intervention contact time	No pharmacotherapy	Abstinence at 6m (biochem. PP)	Unclear

Table 8.34 Included studies from systematic review: Motivational Interviewing versus nothing

Study	Setting	Mean Age	Ethnicity	Female	Participants	Intervention	Control	Adjunct	Outcome	Risk of bias
Harris 2010 ⁽²⁴⁹⁾	USA	19.5	NS	46%	452	MI counselling (smoking cessation) (20-30mins, 4 sessions)	MI counselling (healthy eating) (20-30mins, 4 sessions)	Pharmacotherapy to both (heavy smokers)	Abstinence at 6m (biochem. PP)	Unclear

Table 8.35 Included studies from systematic review: Mobile Phone interventions

Study	Setting	Mean Age	Ethnicity	Female	Participants	Intervention	Control	Adjunct	Outcome	Risk of bias
Borland 2013 ⁽²⁵⁰⁾	Australia	42.1	NS	60%	3530	Internet intervention: personalised tailored internet-delivered (QuitCoach)	Minimal advice	No pharmacological adjunct	7 months (6-month CAR), SR	Low
Borland 2013	Australia	42.1	NS	60%	3530	SMS intervention: interactive automated SMS program (onQ)	Minimal advice	No Pharmacological adjunct	7 months (6-month CAR), SR	Low
Borland 2013	Australia	42.1	NS	60%	3530	Combined internet and SMS intervention	Minimal advice	No Pharmacological Adjunct	7 months (6-month CAR), SR	Low
Ferguson 2015 ⁽²⁵¹⁾	Australia	42.1	NS	51%	284	SMS and self-help booklet	Self-help booklet		Abstinence PP	Unclear
Haug 2013 ⁽²⁵²⁾	Switzerland	18.2	NS	52%	755	SMS (3months)	Online health screen		Abstinence at 6m (sr, pp)	Low
Naughton 2014 ⁽²⁵³⁾	UK	41	NS	53%	602	SMS (90 days), counselling, pharmacotherapy	Counselling, Pharmacotherapy - fairly intensive		Abstinence at 6m (sr, car)	Low

Whittaker 2011⁽²⁵⁴⁾	New Zealand	27	24% Maori	47%	226	SMS and video message (6 months)	General health video message		Abstinence at 6m (SR, car)	Low
---------------------------------------	-------------	----	-----------	-----	-----	----------------------------------	------------------------------	--	----------------------------	-----

Table 8.35 Included studies from systematic review: Telephone interventions

Study	Setting	Mean Age	Ethnicity	Female	Participants	Intervention	Control	Outcome	Risk of bias	
Abdullah 2005⁽²⁵⁵⁾	Hong Kong	NS; > 50% aged 36 - 45	NS	16%	903	Telephone intervention; smoker did NOT initiate contact	Self-help or minimal intervention	Abstinence at 6m (biochem, pp)	Unclear	
An 2006⁽²⁵⁶⁾	USA	57	NS	9%	821	Telephone intervention; smoker did NOT initiate contact	Self-help or minimal intervention	Abstinence at 12m (SR, CAR)	High	
Aveyard 2003⁽²²⁵⁾	UK	41	NS	54%	2471	Telephone intervention; smoker did NOT initiate contact	Self-help or minimal intervention	Abstinence at 12m (biochem, car)	Low	
Borland 2001⁽²⁵⁷⁾	Australia	NS; 37% aged 15 - 29, 26% aged 30 - 39	NS	52%	998	Additional proactive call(s)	Mailed material	Abstinence at 12m (SR, car)	High	
Borland 2003⁽²⁵⁸⁾	Australia	NS; modal age 30 - 49	NS	54%	1578	Additional proactive call(s)		Abstinence at 12m (SR, car)	High	
Borland 2008⁽²⁵⁹⁾	Australia	41	NS	55%	1039	Adjunct telephone intervention; smoker did NOT initiate contact	Brief intervention or counselling	Adjunct: pharmacotherapy to	Abstinence at 12m (SR, car)	High

								both groups		
Boyle 2007 ⁽²⁶⁰⁾	USA	47	NS	58%	1329	Adjunct telephone intervention to pharmacotherapy; smoker did NOT initiate contact	Pharmacotherapy		Abstinence at 12m (SR, PP)	High
Brown 1992 ⁽²⁶¹⁾	Australia	40	NS	62%	45	Adjunct telephone intervention; smoker did NOT initiate contact	Brief intervention or Counselling		Abstinence at 12m (SR, PP)	Unclear
Curry 1995 ⁽²⁶²⁾	USA	41	NS	52%	1137	Telephone intervention; smoker did NOT initiate contact	Self-help or minimal intervention		Abstinence at 12m (biochem, car)	Unclear
Ebbert 2007 ⁽²⁶³⁾	USA	NS	NS	NS	82	Adjunct telephone intervention; smoker did NOT initiate contact	Brief intervention or Counselling		Abstinence at 6m (SR, pp)	High
Ellerbeck 2009 ⁽²³²⁾	USA	47	NS	59%	750	Adjunct telephone intervention to pharmacotherapy; smoker did NOT initiate contact	Pharmacotherapy		Abstinence at 24m (biochem, pp)	Low
Emmons 2005 ⁽²⁶⁴⁾	USA	31	NS	47%	794	Telephone intervention; smoker did NOT initiate contact	Self-help or minimal intervention		Abstinence at 12m (SR, PP)	Unclear
Ferguson 2012 ⁽²⁶⁵⁾	UK: English QUITline	38	NS	55%	2591	Additional proactive call(s)	Standard phonecall	Adjunct: NRT provided to both group	Abstinence at 6m (biochem, car)	Low
Fiore 2004 ⁽²⁶⁶⁾	USA	40	NS	58%	961	Adjunct telephone intervention to pharmacotherapy;	Pharmacotherapy		Abstinence at 12m (biochem,	Unclear

						smoker did NOT initiate contact		car)	
Flöter 2009 ⁽²⁶⁷⁾	Germany	35.9	NS	NS	527	Adjunct telephone intervention; smoker did NOT initiate contact	Brief intervention or counselling	Abstinence at 6m (SR, pp)	High
Gilbert 2006 ⁽²⁶⁸⁾	UK: English QUITline	39	NS	66%	1457	Additional proactive call(s)		Abstinence at 12m (SR, car)	High
Girgis 2011 ⁽²⁶⁹⁾	Australia	29	NS	52%	407	GP-referral to telephone counselling	Usual care	Abstinence at 12m (SR, PP)	High
Graham 2011 ⁽²⁷⁰⁾	USA	35.9	NS	51%	2005	Telephone intervention; smoker did NOT initiate contact	Self-help or minimal intervention	Abstinence at 18m (SR, PP)	High
Halpin 2006 ⁽²⁷¹⁾	USA	67% age 40+	NS	66%	388	Coverage by HMO for TC and pharmacotherapy	Coverage by HMO for pharmacotherapy alone	Abstinence at 6m (SR, pp)	High
Hennrikus 2002 ⁽²⁷²⁾	USA	36	NS	50-64%	2402	Telephone counselling	Group therapy	Abstinence at 24m (biochem, car)	Unclear
Hollis 2007 ⁽²⁴³⁾	USA Oregon QUITline	41	NS	60%	4500	Additional proactive call(s)	Brief counselling	Abstinence at 12m (SR, car)	High
Joyce 2008 ⁽²⁷³⁾	USA	65+	NS	60%	7354	Offer of Quitline: Reactive or proactive counselling vs provider counselling		Abstinence at 12m (SR, pp)	High
Katz 2004 ⁽²⁷⁴⁾	USA	43/40	NS	56%	1141	Interactive TC	Information on guidelines, no specific advice	Abstinence at 6m (biochem, car)	High
Lando 1992 ⁽²⁷⁵⁾	USA	47	NS	50%	1827	Telephone intervention; smoker did NOT initiate	Self-help or minimal intervention	Abstinence at 18m (biochem,	High

						contact		car)	
Lando 1997⁽²⁷⁶⁾	USA	42	NS	56%	509	Adjunct telephone intervention to pharmacotherapy; smoker did NOT initiate contact	Pharmacotherapy	Abstinence at 12m (biochem, pp)	Unclear
Lichtenstein 2000⁽²⁷⁷⁾	USA	NS	NS	NS	1006	Telephone intervention; smoker did NOT initiate contact	Self-help or minimal intervention	Abstinence at 12m (SR, car)	High
Lichtenstein 2008⁽²⁷⁸⁾	USA	NS	NS	NS	1821	Telephone intervention; smoker did NOT initiate contact	Self-help or minimal intervention	Abstinence at 12m (SR, car)	High
Lipkus 1999⁽²⁷⁹⁾	USA	49% aged > 50	African American	52%	266	Telephone intervention; smoker did NOT initiate contact	Self-help or minimal intervention	Abstinence at 16m (SR, pp)	High
MacLeod 2003⁽²⁸⁰⁾	Australia	42	NS	51%	854	Adjunct telephone intervention to pharmacotherapy; smoker did NOT initiate contact	Pharmacotherapy	Abstinence at 6m (SR, CAR)	High
McBride 1999⁽²⁸¹⁾	USA	36	NS	100%	580	Telephone intervention; smoker did NOT initiate contact	Self-help or minimal intervention	Abstinence at 15m (biochem, pp)	Unclear
McClure 2005⁽²⁴⁵⁾	USA	33	NS	100%	275	Telephone intervention; smoker did NOT initiate contact	Self-help or minimal intervention	Abstinence at 12m (biochem, pp)	Unclear
McClure 2011⁽²⁸²⁾	USA	44.5	NS	67%	52	Telephone intervention; smoker did NOT initiate contact	Self-help or minimal intervention	Abstinence at 6m (SR, pp)	High

McFall 1993 ⁽²⁸³⁾	USA	NS; 23% age 18 - 30, 40% age 31 - 45, 30% 45 - 64	NS	70%	1745	Offer of Quitline: Quitline + Self Help	Nothing	Abstinence at 24m (SR, pp)	High
Metz 2007 ⁽²⁸⁴⁾	Germany	47	NS	41%	290	Adjunct Telephone intervention; smoker did NOT initiate contact	Brief intervention or counselling	Abstinence at 12m (SR, pp)	High
Miguez 2002 ⁽²⁸⁵⁾	Spain	35	NS	38%	200	Telephone intervention; smoker did NOT initiate contact	Self-help or minimal intervention	Abstinence at 12m (biochem, car)	Unclear
Miguez 2008 ⁽²⁸⁶⁾	Spain	37	NS	46%	228	Telephone intervention; smoker did NOT initiate contact	Self-help or minimal intervention	Abstinence at 12m (sr, car)	Unclear
Miller 1997 ⁽²⁸⁷⁾	USA	51	NS	49%	1942	Intensive intervention (TC and FTF)	Minimal intervention (TC + FTF)	Abstinence at 12m (biochem, car)	Low
Ockene 1991 ⁽⁴⁷⁾	USA	35	NS	57%	1223	Adjunct Telephone intervention; smoker did NOT initiate contact	Brief intervention or counselling	Abstinence at 6m (SR, pp)	High
Orleans 1991 ⁽²⁸⁸⁾	USA	44	NS	63%	2021	Telephone intervention; smoker did NOT initiate contact	Self-help or minimal intervention	Abstinence at 16m (biochem, car)	Unclear
Orleans 1998 ⁽²⁸⁹⁾	USA	NS; 62% in 20 - 39 age group	African American	64%	1422	2 different interventions during single call (tailored vs standard counselling)		Abstinence at 6m (SR, pp)	High

Osinubi 2003 ⁽²⁹⁰⁾	USA	52	NS	7%	58	Adjunct telephone intervention; smoker did NOT initiate contact	Brief intervention or counselling	Abstinence at 6m (SR, pp)	High
Ossip-Klein 1991 ⁽²⁹¹⁾	USA	43	NS	NS	1813	Offer of Quitline: Quitline + Self Help	Self help	Abstinence at 18m (biochem, car)	Unclear
Ossip-Klein 1997 ⁽²⁹²⁾	USA	NS	NS	61%	177	Telephone intervention; smoker did NOT initiate contact	Self-help or minimal intervention	Abstinence at 6m (SR, pp)	Unclear
Prochaska 1993 ⁽²⁹³⁾	USA	43	NS	62%	756	Telephone intervention; smoker did NOT initiate contact	Self-help or minimal intervention	Abstinence at 18m (sr, pp)	Unclear
Prochaska 2001 ⁽²⁹⁴⁾	USA	38	NS	56%	1447	Telephone intervention; smoker did NOT initiate contact	Self-help or minimal intervention	Abstinence at 18m (sr, car)	High
Rabius 2004 ⁽²⁹⁵⁾	USA	NS; ($\leq 25 / > 25$): 61%/67% F, av. age 22/44,	NS	NS	3522	Additional proactive call(s)		Abstinence at 6m (SR, car)	Unclear
Rabius 2007 ⁽²⁹⁵⁾	USA	43	NS	70%	6322	Additional proactive call(s)		Abstinence at 7m (SR, pp)	Unclear
Reid 1999 ⁽²⁹⁶⁾	Canada	38	NS	48%	396	Adjunct telephone intervention to pharmacotherapy; smoker did NOT initiate contact	Pharmacotherapy	Abstinence at 12m (SR, pp)	Unclear
Rimer	USA	61	NS	63%	1867	Telephone	Self-help or minimal	Abstinence	High

1994⁽²⁹⁷⁾						intervention; smoker did NOT initiate contact	intervention	at 12m (SR, pp)	
Sims 2013⁽²⁹⁸⁾	USA	21.3	NS	58%	410	Additional proactive call(s)		Abstinence at 6m (SR, pp)	Unclear
Smith 2004⁽²⁹⁹⁾	USA	42	NS	61%	632	Additional proactive call(s)		Abstinence at 12m (sr, car)	High
Solomon 2000⁽³⁰⁰⁾	USA	33	NS	100%	214	Adjunct telephone intervention to Pharmacotherapy; smoker did NOT initiate contact	Pharmacotherapy	Abstinence at 6m (biochem, pp)	High
Solomon 2005⁽³⁰¹⁾	USA	34	NS	100%	330	Adjunct telephone intervention to pharmacotherapy; smoker did NOT initiate contact	Pharmacotherapy	Abstinence at 6m (sr, pp)	High
Sood 2009⁽³⁰²⁾	USA	43	NS	62%	990	2 different interventions during single call (reactive counselling vs self-help materials)		Abstinence at 12m (SR, pp)	Low
Sorensen 2007a⁽³⁰³⁾	USA	40	NS	6%	231	Telephone intervention; smoker did NOT initiate contact	Self-help or minimal intervention	Abstinence at 6m (SR, pp)	High
Swan 2003⁽¹⁴⁶⁾	USA	45	NS	57%	1524	Free & clear proactive TC (4 brief calls), access to quitline & S-H materials	Zyban Advantage Program (ZAP) tailored S-H materials, single telephone call, Zuban support line	Abstinence at 12m (SR, pp)	High
Swan 2010⁽³⁰⁴⁾	USA	47.3	NS	67%	1202	Adjunct telephone intervention to pharmacotherapy; smoker did NOT initiate contact	Pharmacotherapy	Abstinence at 6m (SR, pp)	High

Thompson 1993⁽³⁰⁵⁾	USA	41	NS	59%	382	2 different interventions during single call (stage-based counselling vs general info)		Abstinence at 6m (SR, pp)	Unclear
Tzelepis 2011⁽³⁰⁶⁾	Australia	45	NS	50%	1562	Telephone intervention; smoker did NOT initiate contact	Self-help or minimal intervention	Abstinence at 13m (SR, car)	High
Velicer 2006⁽³⁰⁷⁾	USA	51	NS	23%	2054	Adjunct telephone intervention to pharmacotherapy; smoker did NOT initiate contact	Pharmacotherapy	Abstinence at 30m (SR, car)	High
Young 2008⁽³⁰⁸⁾	Australia	37	NS	53%	318	Telephone intervention; smoker did NOT initiate contact	Self-help or minimal intervention	Abstinence at 12m (SR, pp)	High
Zhu 1996⁽³⁰⁹⁾	USA (Quitline)	36	NS	57%	3030	Telephone contact & Self-help	Self-help	Abstinence at 13m (biochem, car)	High
Zhu 2002⁽³¹⁰⁾	USA (Quitline)	38	NS	56%	3282	Additional proactive call(s)		Abstinence at 13m (SR, car)	High
Zhu 2012⁽³¹¹⁾	USA (Quitline)	approx. 45% 25 - 44 and 45% 45 - 64	Chinese Vietnamese eKorean	10%	2278	Additional proactive call(s)		Abstinence at 7m (SR, car)	Low

Table 8.36 Updated review. Included studies: NRT versus placebo

Study	Setting	Mean Age	Female	Participants	Intervention Dose	Control Dose	Support	Outcome	Risk of Bias
Tuisku	Finland	21	52%	291	NRT: Nicotine	Placebo	All received 20 minute	6 month (PP,	High

2016a ⁽³¹²⁾					patch (10mg/16h) X 8/52	Patch X8/52	counselling based on MI X 22	SR)	
Ward 2013 ⁽³¹³⁾	Aleppo, Syria	40	22%	269	NRT: Nicotine patch (dose related to nicotine dependence) - 6 weeks	Placebo Patch	Individual behavioural counselling to both groups. 3x30 min individual counselling and 5x10min phone call	12 month (Prolonged, Biochem)	Low

Table 8.37 Updated review. Included studies: Bupropion plus Varenicline versus Varenicline

Study	Setting	Mean Age	Female	Participants	Intervention	Control	Adjunct	Outcome	Risk of Bias
Ebbert 2014 ⁽³¹⁴⁾	USA	42.2 (Intervention) 41.9 (Control)	45% (Intervention) 49% (Control)	506	Bupropion SR + Varenicline (12 weeks)	Placebo + Varenicline (12 weeks)	Brief advice to both groups	12 month (Prolonged, Biochem)	Low

Table 8.38 Updated review. Included studies: Varenicline plus NRT versus Varenicline

Study	Setting	Mean Age	Female	Participants	Intervention 1	Intervention 2	Adjunct	Outcome	Risk of Bias
Ramon 2014 ⁽³¹⁵⁾	Spain	45.1	42%	341	NRT (Nicotine Patch 21mg for 11weeks) + Varenicline (12 weeks)	Placebo Nicotine Patch + Varenicline (12 weeks)	Both groups received behavioural support 10-15min x 5	6 month (CAR, Biochem)	Low
Koegelenberg 2014 ⁽³¹⁶⁾	South Africa	46.3	15%	446	NRT (Nicotine Patch 15mg for 14 weeks) + Varenicline	Placebo Nicotine Patch + Varenicline		6 month (CAR, Biochem)	Low

Table 8.39 Updated review. Included studies: Varenicline versus NRT versus Bupropion

Study	Setting	Mean Age	Ethnicity	Female	Participants	Intervention 1	Intervention 2	Intervention 3	Outcome	Risk of Bias
Zincir 2013⁽³¹⁷⁾	Turkey	NS	NS	NS	251	NRT with patch or gum, or Combination NRT (Patch&gum) - 12 weeks	Varenicline (12 weeks)	Bupropion (12 weeks)	6 month (PP, Biochem)	High

Table 8.40 Updated review. Included studies: Varenicline versus NRT

Study	Setting	Mean Age	Ethnicity	Female	Participants	Intervention 1	Intervention 2	Intervention 3	Outcome	Risk of Bias
Tulloch 2016⁽³¹⁸⁾	Canada	48.61	NS	47%	737	NRT - 10 weeks (21mg daily)	NRT (35mg daily) + Nicotine gum or inhaler - 22 weeks	Varenicline (1mg BID) - 24 weeks	12 month (CAR, Biochem)	High (open-label)
Study	Setting	Mean Age	Ethnicity	Female	Participants	Intervention 1	Intervention 2		Outcome	Risk of Bias
Tuisku 2016b⁽³¹²⁾	Finland	22 (NRT), 21 (Varenicline)	NS	45% (NRT) 50% (Varenicline)	291	NRT: Nicotine patch (15mg/16h)	Varenicline		6 month (PP, SR)	High

Table 8.41 Updated review. Included studies: Telephone interventions

Study	Setting	Mean Age	Ethnicity	Female	Participants	Intervention	Control	Adjunct	Outcome	Risk of Bias
Killen 2008 ⁽³¹⁹⁾	USA	Telephone support: M 46.56, F 44.59; CBT: M 45.9, F 46.23	82% White	40%	301	CBT - on site - 30 mins X 4 sessions	Support telephone calls - 5 mins X 4 sessions	9 weeks of Bupropion and 8 weeks of NRT to all participants in Open-label 8 week period	12 months, CAR, Biochem	Low
Zwar 2014 ⁽³²⁰⁾	Australia	42.6 (Practice Nurse), 43.5 (QUITline), 42.1 (Control)	NS	55% (PN), 55% (QUITline), 53% (Control)	2390	Practice Nurse Intervention (face-to-face or by telephone)	Usual care (GP care: Brief advice)	Pharmacotherapy offered to all participants (NRT, bupropion or varenicline)	12 months (CAR 10 Months, SR)	Low
Zwar 2014	Australia	42.6 (Nurse), 43.5 (QUITline), 42.1 (Control)	NS	55% (PN), 55% (QUITline), 53% (Control)	2390	QUITline referral by GP (proactive call)	Usual care (GP care: Brief advice)	Pharmacotherapy offered to all participants (NRT, bupropion or varenicline)	12 months (CAR 10 Months, SR)	Low

Table 8.42 Updated review. Included studies: Internet interventions

Study	Setting	Mean Age	Female	Participants	Intervention	Control	Adjunct	Outcome	Risk of Bias
Borland 2013 ⁽²⁵⁰⁾	Australia	42.1	60%	3530	Internet intervention: personalised tailored internet-delivered (QuitCoach)	Minimal advice	No pharmacological adjunct	7 months (6-month CAR), SR	Low

Borland 2013	Australia	42.1	60%	3530	SMS intervention: interactive automated SMS program (onQ)	Minimal advice	No pharmacological adjunct	7 months (6-month CAR), SR	Low
Borland 2013	Australia	42.1	60%	3530	Combined internet and SMS intervention	Minimal advice	No pharmacological adjunct	7 months (6-month CAR), SR	Low
Stanczyk 2016⁽³²¹⁾	Holland	45.7	61%	2099	Enhanced internet (video messages)	Brief generic text advice	No pharmacological adjunct	12 months (CAR, Biochem)	Unclear
Stanczyk 2016	Holland	45.7	61%	2099	Basic internet (text messages)	Brief generic text advice	No pharmacological adjunct	12 months (CAR, Biochem)	Unclear

Table 8.43 Updated review. Included studies: Mobile phone interventions

Study	Setting	Mean Age	Female	Participants	Intervention	Control	Adjunct	Outcome	Risk of Bias
Borland 2013⁽²⁵⁰⁾	Australia	42.1	60%	3530	Internet intervention: personalised tailored internet-delivered (QuitCoach)	Minimal advice	No pharmacological adjunct	7 months (6-month CAR), SR	Low
Borland 2013	Australia	42.1 (Females)	60%	3530	SMS intervention: interactive automated SMS program (onQ)	Minimal advice	No pharmacological adjunct	7 months (6-month CAR), SR	Low
Borland 2013	Australia	42.1 (Females)	60%	3530	Combined Internet and SMS intervention	Minimal advice	No pharmacological adjunct	7 months (6-month CAR), SR	Low

Table 8.44 Updated review. Included studies: Motivational interviewing

Study	Setting	Mean Age	Ethnicity	Female	Participants	Intervention	Control	Adjunct	Outcome	Risk of Bias
Bock 2014 ⁽²³⁸⁾	USA	39.6	53% White	69%	846	Motivational interviewing (45 minutes trained professional & follow-up telephone calls) + brief physician advice + NRT	Usual care: brief physician advice + NRT	All received 8 weeks NRT	12 months (PP 7-day, Biochem)	Low

Table 8.45 Updated review. Included studies: Individual behavioural counselling

Study	Setting	Mean Age	Ethnicity	Female	Participants	Intervention	Control	Adjunct	Outcome	Risk of Bias
Killen 2008 ⁽³¹⁹⁾	USA	Telephone support: M 46.56, F 44.59; CBT: M 45.9, F 46.23	82% White	40%	301	CBT - on site - 30 mins X 4 sessions & Telephone Counselling	Support Telephone calls - 5 mins X 4 sessions	9 weeks of Bupropion and 8 weeks of NRT to all participants in open-label 8 week period	12 months (CAR, Biochem)	Low

Table 8.46 Updated review. Included studies: Nursing interventions


Study	Setting	Mean Age	Female	Participants	Intervention	Control	Adjunct	Outcome	Risk of Bias
Zwar 2014 ⁽³²⁰⁾	Australia	42.6 (Practice Nurse), 43.5 (QUITline), 42.1 (Control)	545% (PN), 57% (QUITline), 58% (Control)	2390	Practice Nurse Intervention (face-to-face or by telephone)	Usual care (GP care: brief advice)	Pharmacotherapy offered to all participants (NRT, bupropion or varenicline)	12 months (CAR 10 Months, SR)	Low
Zwar 2014	Australia	42.6 (Practice Nurse), 43.5	55% (PN), 57%	2390	QUITline referral by GP	Usual care (GP care: brief advice)	Pharmacotherapy offered to all	12 months	Low

		(QUITline), 42.1 (Control)	(QUITline), 58% (Control)		(proactive call)	brief advice)	participants (NRT, bupropion or varenicline)	(CAR 10 Months, SR)	
--	--	-------------------------------	------------------------------	--	---------------------	------------------	--	---------------------------	--

Table 8.47 Search strategy for studies involving the Allen Carr method

Database (Date of search)	Search string	Results
Pubmed (20/5/2016)	Allen AND Carr AND smoking	8
Embase (20/5/2016)	Allen AND Carr AND smoking	15
Cochrane Registry of Controlled Trials (20/5/2016)	Allen AND Carr AND smoking	0

Figure 8.1 Flowchart of included studies for the Allen Carr smoking cessation intervention



Appendix 9 Search details for review of smoking cessation interventions in pregnancy

Figure 9.1 Flow diagram of included studies from the updated search: Psychosocial and pharmacological interventions for smoking cessation in pregnancy




Table 9.1 Search string used in Medline (PubMed)

Pubmed 18/10/2016	Search strings	Results
Searches #1	((("Smoking Cessation"[Mesh]) OR ("Tobacco Use Cessation"[Mesh]) OR (Stop* smoking) OR (Quit* smoking)))	31604
#2	((("Pregnancy"[Mesh]) OR ("Pregnant Women"[Mesh]) OR ("Postpartum Period"[Mesh]) OR ("Postnatal Care"[Mesh]) OR (Pregnancy [tiab]) OR (Pregnant [tiab]) OR (Postpartum [tiab]) OR (Post-partum [tiab]) OR (Postnatal [tiab]) OR (Post-natal [tiab]) OR (Prenatal [tiab]) OR (Pre-natal [tiab])))	984290
#3	#1 AND #2	2264
#4	#3 AND Filters: <ul style="list-style-type: none"> • <u>From 2013/01/01 to 2016/10/18</u> • Clinical Trial; Randomized Controlled Trial; Review; Journal Article, Systematic Reviews; Meta-Analysis • Humans 	391

((((("Pregnancy"[Mesh]) OR ("Pregnant Women"[Mesh]) OR ("Postpartum Period"[Mesh]) OR ("Postnatal Care"[Mesh]) OR (Pregnancy [tiab]) OR (Pregnant [tiab]) OR (Postpartum [tiab]) OR (Post-partum [tiab]) OR (Postnatal [tiab]) OR (Post-natal [tiab]) OR (Prenatal [tiab]) OR (Pre-natal [tiab]))))) AND (((("Smoking Cessation"[Mesh]) OR ("Tobacco Use Cessation"[Mesh]) OR (Stop smoking) OR (Quit* smoking))))))*

Appendix 10 Included studies smoking cessation interventions in pregnancy

Table 10.1 Included studies: NRT and BCS versus placebo and BCS, n = 5

Study	Country	Ethnicity	Participants (n)	Gestational cut-off (weeks)	Intervention	Control	BCS	Outcome	Risk of bias
Berlin 2014 ⁽³²²⁾	France	European (96%)	402	≤ 20	Nicotine patch 10mg/16hr or 15mg/16hr for eight weeks. Dose adjusted to max of 30mg/hr	Placebo patch	Behavioural support at each visit from a midwife or doctor who had been specifically trained. At least 10mins of counselling/visit	Continuous abstinence until birth and birth weight	Low
Coleman 2012 ⁽³²³⁾	England	White British (97%)	1050	≤ 24	Nicotine patch 15mg/16h	Placebo patch	Behavioural support provided by research midwives. First face to face and three by telephone. Face to face if collected a second month's supply	Continuous abstinence until birth	Low
Kapur 2001 ⁽³²⁴⁾	Canada	NS	30	≤ 24	Nicotine patch 15mg/18 hr for eight weeks followed by 10mg/18hr for two weeks followed by 5mg/18hr for two weeks	Placebo patch	Four counselling sessions. One of the investigators maintained weekly telephone contact	Point prevalence abstinence after eight weeks of treatment	Low
Oncken 2008 ⁽²²⁹⁾	US	Hispanic	194	≤ 26	Nicotine gum 2mgs for six weeks followed by a six week taper period	Placebo gum	Two 35 min counselling sessions in English or Spanish delivered by a research assistant using a motivational interviewing approach	Seven day point prevalence abstinence after six weeks of gum use and at end of pregnancy	Low
Wisborg 2000 ⁽³²⁵⁾	Denmark	NS	250	< 22	Nicotine patch/16hr: 15mg/16hr for eight weeks followed by	Placebo	Four sessions of smoking cessation counselling with midwife independent	Seven day point prevalence abstinence at	Low

					10mg/16hr for three weeks		of routine antenatal care visits.	2 nd , 3 rd and 4 th prenatal visits	
--	--	--	--	--	---------------------------	--	-----------------------------------	---	--

BCS: behavioural cessation support

Table 10.2 Included studies: NRT patches and BCS versus BCS, n = 3

Study	Country	Ethnicity	Participants (n)	Gestational cut-off (weeks)	Intervention	Control	BCS	Outcome	Risk of bias
El-Mohandes 2013 ⁽³²⁶⁾	US	African-American Hispanic	52	≤ 30	Nicotine patch 7mg or 14mg or 21mg/24 hr – two dosing schedules for ten weeks based on baseline salivary cotinine levels	None	SCRIPT* programme – reinforcement and behavioural methods at four visits	Seven day point prevalence after six weeks and between 32 and 34 weeks' gestation	High
Hotham 2006 ⁽³²⁷⁾	Australia	NS	40	≤ 28	Nicotine patch 15mg/16h for 12 weeks	None	Counselled for approx 5 minutes at randomisation and for approx 2 minutes at subsequent visits by a researcher or midwives who had undergone training with QUIT	Self reported abstinence at end of treatment and within 48 hours of birth	High
Pollak 2007 ⁽³²⁸⁾	US	White 69%, Black 24%	181	≤ 25	Nicotine patch or gum (2mg) or lozenge. Patch 7mgs/16hr if less than 10cigs/day, 14mg/16hr if 10-14 14 cigs/day and 21mg/day if ≥ 15 cigs/day	None	Six one-on-one counselling sessions. Five face-to-face at prenatal visits and one via telephone. Contacts designed to be relapse-sensitive and to coincide with prenatal visits	Seven day point prevalence after seven weeks of treatment and between 32-34 weeks' gestation	High

*The SCRIPT (Smoking Cessation and Reduction in Pregnancy Treatment Program)

Table 10.3 Included studies: Bupropion and BCS versus BCS, n = 1

Study	Country	Ethnicity	Participants (n)	Gestational cut-off (weeks)	Intervention	Control	BCS	Outcome	Risk of bias
Stotts 2015 ⁽³²⁹⁾	US	African-American Hispanic	11	≤ 26	Bupropion 150mg/day for three days followed by 150mg bd for eight weeks	Placebo	Four, weekly, 15 minute smoking cessation counselling sessions delivered by a research nurse	Seven day point prevalence abstinence at end of treatment	Unclear

Table 10.4 Included studies: Counselling, n = 44

Study	Country	Ethnicity	Participants (n)	Gestational cut-off (weeks)	Intervention	Control	Comparison type	Outcome	Risk of bias
Baric 1976 ⁽³³⁰⁾	England	NS	110	< 20	One to one counselling from a senior medical student. Half were given a diary and a gift of a free smoking diary	Usual care	Counselling (single intervention) versus usual care	Self-reported abstinence 11 weeks after baseline visit	High
Cinciripini 2000 ⁽³³¹⁾	USA	NS	82	≤ 30	Control and posted a video with six 25-30 minute vignettes	Quit calendar and tip guide	Counselling (single intervention) versus less intensive Intervention	Biochemically validated point prevalence abstinence within 2-3 days of quit, 4-5 weeks after quit and 1 month (0-5) postpartum	High
Cook 1995 ⁽³³²⁾	USA	NS	150	< 24	Control and regular meetings with	Discussion of risks by a nutritionist	Counselling (multiple	Biochemically validated at	High

					smoking cessation counsellor + physician reinforcement at each visit. Feedback from urine cotinine	and resident physician at initial visit	intervention) versus less intensive intervention	term or birth, > 50% reduction in mean cotinine and mean birth weight	
Cummins 2016 ⁽³³³⁾	USA	NS	1173	< 27	Telephone counselling provided by veteran staff members of quitline and self-help materials	Self-help materials only	Counselling (multiple intervention) versus less intensive intervention	Self-reported abstinence 30-day abstinence in third trimester (about 29 weeks), 90 day abstinence at two months postpartum and 190 abstinence at six months postpartum	High
Dornelas 2006 ⁽³³⁴⁾	USA	66% Hispanic	107	< 30	One 90 min psychotherapy session by mental health therapist. Bimonthly phone calls	Usual care and booklet, chart prompt and audited chart	Counselling (single intervention) versus less intensive intervention	Biochemically validated seven day point prevalence abstinence and six months (6-11) postpartum	High
Dunkley 1997 ⁽³³⁵⁾	UK	NS	100	< 18	Brief contact (< 5mins) x eight visits with trained midwives	Usual care	Counselling (single intervention) versus usual care	Self-reported abstinence at 37 weeks and one month (0-5) postpartum	High
Eades 2012 ⁽³³⁶⁾	Australia	Aboriginal or Torres Straight Islander	263	≤ 20	Advised to quit "cold turkey" at 1st visit. Bring a support person to 2 nd visit. Offered NRT at 3 rd	Usual care	Counselling (tailored intervention) versus usual care	Biochemically validated point prevalence abstinence at 36 weeks	High

					visit. Follow up visits with health workers and midwives				
Ershoff 1989 ⁽³³⁷⁾	USA	64% white	242	< 18	Control and eight self-help booklets, taught behavioural strategies and asked to read first one, other mailed weekly.	Two page pamphlet, two minutes with a health educator and advised re free cessation programme	Counselling (single intervention) versus less intensive intervention	Biochemically validated abstinence at 34 weeks	High
Ershoff 1999 ⁽³³⁸⁾	USA	60% white	257	≤ 26	Control and 4-6 x 10-15 min phone counselling sessions by nurse educators trained in motivational interviewing	Received a 32 page self-help booklet	Counselling (single intervention) versus less intensive intervention	Biochemically validated cessation at 34 weeks and mean cigs/day	High
Gielen 1997 ⁽³³⁹⁾	USA	African American: 81%, 89%	391	≤ 28	Written information, a 15 minute 1:1 counselling session, educational materials, reinforcement and two letters of encouragement	Usual clinic + inpatient smoking cessation	Counselling (multiple intervention) versus usual care	Biochemically validated seven day point prevalence abstinence in hospital after delivery, six months (6-11) postpartum, > 50% reduction in cotinine from baseline to late pregnancy	High
Hajek 2001 ⁽³⁴⁰⁾	NS	NS	732	12	Midwives received two hours of training on CO monitor and stage of change advice and CO assessments. Participants received	Midwives received one hour of training to discuss the study and were asked to provide usual care and any usual	Counselling (tailored intervention) versus usual care	Biochemically validated point prevalence abstinence at birth and self-reported continuous	High

					written advice, motivational materials and offer of buddying to another pregnant smoker for support	pamphlets		abstinence at six months (6-11) postpartum	
Hartmann 1996 ⁽³⁴¹⁾	USA	White: 78%, 74%	219	≤ 36	Counselling by residents at each visit, Windsor's guide, contacted by smoking cessation counsellor and quitters sent encouraging postcard each week	Usual care	Counselling (multiple intervention) versus usual care	Biochemically validated abstinence at last prenatal visit, > 50% reduction in self-reported smoking and mean cigs/day	High
Haug 1994 ⁽³⁴²⁾	Norway	NS	322	≤ 12	GP advice < 15 minutes at first visit, written information, invited to consult GP after one, six, 12 and 18 months	Usual care	Counselling (multiple intervention) versus usual care	Self-reported smoking abstinence six months after study entry, biochemically validated 12 months after start (0-5 months postpartum), self-reported abstinence at 15 (6-11) and 18 (12-17) months postpartum	High
Hegaard 2003 ^(343, 344)	Netherlands	NS	647	≤ 22	30-40 minute consultation, written information, invitation to join smoking cessation programme based on CBT, all offered NRT (2mg gum or 15mg	Usual care (included advice in a 30 minute consultation)	Counselling (tailored intervention) versus usual care	Biochemically validated cessation at 37 weeks, mean birth weight, low birth weight and preterm	High

					patch/16 hours) for 11 weeks + encouragement at subsequent clinic visits			births	
Kendrick 1995 ⁽³⁴⁴⁾	USA	NS	1,885	≤ 32	Based on stages of change but differed by state. 1-5 minutes counselling or brief clinic based counselling or six minutes with brochures and written materials	Usual care (not otherwise specified by usual clinic staff)	Counselling (multiple intervention) versus usual care	Biochemically validated point prevalence abstinence at eight months' gestation	High
Lawrence 2003 ⁽³⁴⁵⁾	UK	NS	613	≤ 19	Midwives trained on theory of transtheoretical model (TTM). Assessed by midwife and given TTM based self-help manual and interactive computer program giving individualised advice	Usual care. Midwives asked to give a booklet to women	Counselling (multiple intervention) versus usual care	Biochemically validated point prevalence abstinence at 28-30 weeks, 10 days after birth and at 0-5 months postpartum	High
Lee 2015 ⁽³⁴⁶⁾	USA	56% Black 12% Hispanic	277	≤ 25	Cognitive behavioural therapy provided by a health educator- two prenatal and two postnatal	Best practice - brief advice based on the 5A's	Counselling (multiple intervention) versus usual care	Biochemically verified seven day point prevalence abstinence in late pregnancy and at one and five months postpartum	High
Lillington 1995 ⁽³⁴⁷⁾	USA	53% African American	34	NS	Assessment of motivation and intention. Bilingual health educators. 15 minute individual	Usual care. Printed information and group quit message	Counselling (multiple intervention) versus usual care	Self-reported smoking cessation at nine months' gestation and	High

					counselling, self-help guide, how to win prizes and booster postcard every month			six weeks (0-5 months) postpartum	
Loeb 1983 ⁽³⁴⁸⁾	USA	NS	963	NS	Letter of invitation, reminder letter, group information meeting, individual session with trained smoking counsellor, six 1.5 hour group sessions, once a week and subsequent optional groups, individual sessions + phone calls	Usual care	Counselling (tailored intervention) versus usual care	Self-reported smoking cessation in late pregnancy. Biochemically validated with cord blood thiocyanate	High
Mayer 1990 ⁽³⁴⁹⁾	USA	75% White	149	≤ 28	20 min 1:1 counselling and behavioural change manual focusing on contracting and self-monitoring (CBT)	Usual care which included printed information	Counselling (multiple intervention) versus usual care	Self-reported smoking cessation at nine months' gestation and approx. 4.7 weeks (0-5 months) postpartum	High
McBride 1999 ⁽³⁵⁰⁾	USA	88% White	501	< 20	Received a personalised letter, a relapse prevention kit, a booklet, three antenatal counselling phone calls, three additional counselling calls in first four months after birth and three newsletters	Self-help booklet and suggestions for quitting	Counselling (multiple intervention) versus less intensive intervention	Self-reported and biochemically validated 7 day point prevalence abstinence at 28 weeks. Abstinence at two (0-5), six (6-11) and 12 (12-17) months	High

								postpartum (combined baseline smokers & spontaneous quitters)	
McLeod 2004 ⁽³⁵¹⁾	New Zealand	NS	272	NS	Four armed. Intervention one: midwife training-smoking cessation and support. Intervention two: midwife - breastfeeding support for smokers. Intervention three: Interventions one and two combined. Control and intervention two compared with interventions one and three	Usual care and midwife training and support to implement education and support for breastfeeding for women who smoked	Counselling (single intervention) versus usual care	Biochemically validated cessation at 28 and 36 weeks and at six weeks and four months (0-5) postpartum. Breastfeeding outcomes also reported.	High
Messimer 1989 ⁽³⁵²⁾	USA	98% White	59	< 28	Control and use of American Lung Association materials, encouragement to send off for materials and slide tape presentation at first visit	Three counselling sessions with physicians, removal of ashtrays and staff asked not to smoke in front of patients	Counselling (multiple intervention) versus less intensive intervention	Self-reported smoking abstinence at 32-36 weeks and at first postpartum visit (0-5 months)	High
Moore 2002 ⁽³⁵³⁾	England	NS	1090	≤ 32	Midwives spent ≥ 5mins introducing a series of 5 self-help booklets based on stages of change theory. Given copy of 1 st booklet and remainder posted	Usual care	Counselling (single intervention) versus usual care	Self-reported 7 day point prevalence abstinence at 26 weeks (94% biochemically validated).	High

								Self-reported mean cigs/day in late pregnancy.	
Panjari 1999 ⁽³⁵⁴⁾	Australia	NS	1013	≤ 20	Control and four counselling sessions by a midwife specifically trained and employed to provide smoking cessation counselling using CBT. Video, interactive discussions and strong verbal messages. Followed by a 5-10min personalised counselling session	Usual care (advice and pamphlet distribution during a group info session)	Counselling (single intervention) versus less intensive intervention	Biochemically validated self-reported smoking cessation at 36 weeks, at six weeks (0-5 months) and at six months (6-11 months) postpartum. Mean birth weight, proportion < 2.5kg and preterm birth	High
Parker 2007 ⁽³⁵⁵⁾	USA	Majority White	736	≤ 26	Received quit kit, enrolled in quit and win monetary incentive lottery programme and up to three motivational interviewing (MI) phone calls	Received self-help materials (quit kit and video - SCRIPT)	Counselling (multiple intervention) versus less intensive intervention	Biochemically validated self-reported smoking cessation at 32 weeks, six weeks and six months postpartum (outcomes not reported)	High
Patten 2009 ⁽³⁵⁶⁾	USA	Alaskan natives	33	≤ 24	Self-help guide adapted from SCRIPT trials, 15-25 min face to face counselling based on 5A's, a video, 4 x 10-15 minute proactive interactive phone sessions	5A's component treatment, five minute face to face at 1st visit and culturally specific brochures	Counselling (multiple intervention) versus less intensive intervention	Biochemically validated tobacco use 60 days post randomisation (late pregnancy)	High

Pbert 2004 ⁽³⁵⁷⁾	USA	NS	434	< 32	Dissemination intervention consisted of provider training based on national CPGs, office management system and establishment of program boards. Based on MI and 4 "As" from the SCRIPT trial.	Usual care, in which no training or intervention occurred	Counselling (single intervention) versus usual care	Biochemically validated smoking cessation at one month postpartum (late pregnancy), at three (0-5) and at six (6-11) months postpartum. Mean cigs/day	High
Price 1991 ⁽³⁵⁸⁾	USA	70% White	141	≤ 28	Tailored educational videotape for 6.5 minutes, pamphlet on quitting, a second four minute video one month later and questions answered by health educator	Usual care	Counselling (single intervention) versus usual care	Biochemically validated smoking cessation two or three weeks prior to delivery. Mean cigs/day	High
Rigotti 2006 ⁽³⁵⁹⁾	USA	NS	421	≤ 26	Control and a series of telephone calls accompanied by additional mailed written materials. Each participant had a trained counsellor who offered up to 90 minutes of counselling in pregnancy and up to 15 minutes postpartum.	Usual care and mailed a validated smoking cessation booklet. Less than five minutes counselling at enrolment by a trained counsellor	Counselling (multiple intervention) versus less intensive intervention	Biochemically validated seven day point prevalence abstinence at 28 weeks to term, at three (0-5) months postpartum and > 50% reduction in cigs/day	High
Secker-Walker 1994 ⁽³⁶⁰⁾	USA	NS	513	< 25	Counselling from a trained health educator. Follow up at 2nd visit, 36 weeks and six weeks postpartum,	Usual advice about smoking provided by obstetrician or midwife	Counselling (multiple intervention) versus usual care	Self-reported smoking cessation at 36 weeks (75% biochemically	High

								validated). Self-reported cessation at 8-15 months (6-11), 16-24 months (18) and 25-54 months postpartum. Mean + low birth weight, PPROM, placenta praevia + placental abruption	
Secker-Walker 1997 ⁽³⁶¹⁾	USA	98% White	49	NS	Control and a 29 minute video of four women going through process of quitting during pregnancy. Based on social learning theory	Advice from obstetrician or nurse-midwife and a booklet on quitting	Counselling (single intervention) versus less intensive intervention	Biochemically validated cessation at 36 weeks	High
Secker-Walker 1998 ⁽³⁶²⁾	USA	NS	291	NS	Structured smoking cessation protocol provided by physicians trained in its use. 1st, 2nd, 3rd and 5th visits.	Received a baseline questionnaire, a booklet, physician advice, CO measured and a brief standardised health risk message from a research nurse	Counselling (multiple intervention) versus less intensive intervention	Biochemically validated seven day point prevalence abstinence at 36 weeks and one year postpartum. Mean birth weight, low birth weight, mean cigs/day at 36 weeks and one year postpartum. Preterm	High

								births.	
Stotts 2002 ⁽³⁶³⁾	USA	NS	269	28	Control and 20-30 minutes of MI telephone counselling, a personalised, stages of change based feedback letter and a final MI call 4-5 days after the feedback letter was sent	Provided with MI counselling (3-5 minutes) and a series of eight motivational self help books	Counselling (multiple intervention) versus less intensive intervention	Biochemically validated cessation at 34 weeks. Self-reported smoking cessation at six weeks and three and six months postpartum	High
Stotts 2004 ⁽³⁶⁴⁾	USA	NS	54	< 28	MI intervention over eight weeks consisting of one face to face MI session, three MI based telephone counselling calls and a personalised feedback letter providing assessment results	Usual care which included physicians or nurses acknowledging smoking and advising quitting	Counselling (multiple intervention) versus usual care	Biochemically validated cessation at post treatment assessment (late pregnancy)	High
Tappin 2000 ⁽³⁶⁵⁾	Scotland	NS	97	NS	Received 2-5 MI sessions (mean 2.6hrs) at home by a midwife with 3 weeks training in smoking cessation counselling	Usual care (received usual advice which should include information about smoking)	Counselling (single intervention) versus usual care	Biochemically validated cessation at greater than or equal to 27 weeks. Mean birth weight, preterm birth and stillbirths	High
Tappin 2005 ^(366, 367)	Scotland	NS	756	≤ 24	Control and offered 2-5 additional home visits of about 30 minutes from same study midwife	Midwives provided standard health promotion including information from a book given to all pregnant women in Scotland	Counselling (single intervention) versus usual care	Biochemically validated and self-reported quitting after the 36 week visit , mean birth weight, low and very	High

								low birth weight, perinatal deaths, admission to NICU, preterm delivery, stillbirths and neonatal deaths	
Thornton 1997 ⁽³⁶⁸⁾	Ireland	NS	367	NS	Control and structured 1:1 counselling by a trained facilitator. Partners invited to be involved in the programme. Received an information pack which included a self-help booklet and invited to join a support group. A CO monitor was available for the intervention group.	Routine prenatal advice on a range of health issues from midwives and obstetricians	Counselling (tailored intervention) versus usual care	Biochemically validated abstinence and relapse prevention at delivery and at three months postpartum, mean birth weight, low birth weight, preterm birth, perinatal deaths and admission to NICU	High
Tsoh 2010 ⁽³⁶⁷⁾	USA	NS	42	< 26	Tailored advice from "Video Doctor" designed to simulate an ideal discussion with a prenatal healthcare professional who provided non-judgmental counselling following principles of MI. At the end of each session two	Usual care	Counselling (multiple intervention) versus usual care	Self-reported 30 day abstinence after one month and two months. Mean reduction in cigs smoked/day and days smoked.	High

					documents printed automatically (a cueing sheet for providers and education worksheet for participants)				
Valbo 1996 ⁽³⁶⁹⁾	Norway	NS	130	18	Anaesthetist provided two 45 minute sessions at 2 week intervals of a protocol based script (Handbook of the American Society of Clinical Hypnosis). Two different tapes played after hypnosis established.	Usual care	Counselling (single intervention) versus usual care	Self-reported abstinence at birth, mean cigs/day at birth and self reported reduction in smoking	High
Walsh 1997 ⁽³⁷⁰⁾	Australia	NS	252	< 26	CBT. 2-3 minute standardised information from doctor, 14 minute video, 10 minute standardised information, counselling from midwife after video, self-help manual, 4 packets of confectionary gum, lottery chance for biochemically validated abstinence at next visit, social support, letters, 34 weeks, 5 minute counselling from doctor and 1-2minute risk advice from doctor	Women advised to stop smoking by doctors and midwives. Midwife provided a package which included (sticker, pamphlet and a two page cessation guide (not specifically tailored to pregnancy)	Counselling (tailored intervention) versus less intensive intervention	Biochemically validated abstinence at 34 weeks and at 6-12 weeks postpartum and preterm births	High
Windsor	USA	57% Black	206	< 32	Received a ten	Smoking cessation	Counselling	Biochemically	High

1985 ⁽³⁷¹⁾					minute standardised counselling session from a health educator, a self-help guide to quit smoking and a pamphlet	advice routinely given at prenatal visits: 2-3 minutes within a group prenatal education session at the first visit	(multiple intervention) versus usual care	validated point prevalence abstinence mid pregnancy, during last month of pregnancy or within 48hrs of birth. Self-reported reduction in late pregnancy	
Windsor 1993 ⁽³⁷²⁾	USA	52% Black	814	< 32	15 minute standardised cessation skills and risk counselling session based on CBT from trained female health education counsellor, a seven day self-directed cessation guide, clinic reinforcement, social support and monthly newsletter.	Two minute talk in a 30 minute group session. Given two pamphlets including details and contact of local quit programme	Counselling (multiple intervention) versus less intensive intervention	Biochemically validated point prevalence abstinence at four to eight weeks after first visit and at 32 weeks	High
Windsor 2011 ⁽³⁷³⁾	USA	NS	993	< 32	Received Assist procedures 4 through 8: 14 min video, written guide with 10 day self-help guide, a ≤ 10 min MI counselling session	Received four elements of the "5As" best practice guidelines	Counselling (multiple intervention) versus less intensive intervention	Biochemically validated point prevalence abstinence in late pregnancy (> 60 days after first visit and < 90 days postpartum)	High

Table 10.5 Included studies: Health education studies, n = 6

Study	Country	Ethnicity	Participants (n)	Gestational cut-off (weeks)	Intervention	Control	Outcome	Risk of bias
Burling 1991 ⁽³⁷⁴⁾	USA	52% White	139	any	Personal letter from Chief + pamphlet, CO test mentioned and simple guidelines for self-directed smoking cessation	Usual care	Biochemically validated point prevalence abstinence at 34 weeks	High
Herbec 2014 ⁽³⁷⁵⁾	UK	94% White	200	NS	Fully automated smoking cessation website targeted to pregnancy (tailored, personalised and structured) "MumsQuit"	information only website	Self reported continuous abstinence (4 week abstinence) 8 weeks post baseline	High
Hjalmarson 1991 ⁽³⁷⁶⁾	Sweden		653	< 12	Self-help manual on stopping smoking, based on Windsor 1985	Given information sheet by their doctor as included in the last pages of the self-help manual	Biochemically validated abstinence at 30-34 weeks and at eight weeks (0-5 months) postpartum. Mean birth weight, births < 36 weeks, low birth weight*, mean cigs/day at 30-34 weeks among baseline smokers	High
Lilley 1986 ⁽³⁷⁷⁾	England		145	< 28	Control + ten min advice from junior doctor based on booklet, a leaflet and copies of booklet for family, letter to GP, letter to woman and pre-planned home visit	Usual care with possible exposure to a concurrent television series on stopping smoking in pregnancy	Self reported abstinence 9-16 weeks after booking visit. Mean cigs/day	High
Naughton 2012 ⁽³⁷⁸⁾	England	100% White	198	< 21	Tailored self-help leaflet by post. Automated tailored text messages. Eighty sent over eleven weeks. Participants could request instant response supportive text 24 hours a day	Non-tailored self help leaflet and the same assessment texts as	Biochemically validated 7 day point prevalence abstinence at three month follow-up. Self-reported four week point prevalence	High

						intervention arm but no intervention texts	abstinence initiation and frequency of quit attempts and seven day point prevalence abstinence at three and seven weeks after enrolment	
Petersen 1992 ⁽³⁷⁹⁾	USA	Majority White	90	≤ 24	Brief counselling by a healthcare professional. Pregnancy specific self-help manual and an audiotape on safe exercise and relaxation posted to participants. Behavioural strategies for quitting and maintenance section for postpartum period.	Brief repeated counselling by a healthcare professional and information about community resources posted to participants	Self-reported abstinence at six months gestation and at eight weeks (0-5) postpartum	High

Table 10.6 Included studies: Financial incentives studies, n = 3

Study	Country	Ethnicity	Participants (n)	Gestation (weeks)	Intervention	Control	Outcome	Risk of bias
Harris 2015 ⁽³⁸⁰⁾	USA	88% White	17	≤ 12	A web based contingency management programme and financial incentives (\$100 cash)	A phone-delivered cessation counselling programme (Smoking Cessation for Healthy Births)	Biochemically validated continuous abstinence in late pregnancy. Abstinence was determined by self-reported smoking using a timeline follow-back calendar and urinary cotinine values.	High
Ondersma 2012 ⁽³⁸¹⁾	USA	90% Black	114	≤ 27	Incentives. Maximum of five episodes of reinforcement (in	Usual care	Biochemically validated seven day	High

					the form of retail gift cards worth \$50)		point prevalence abstinence at ten week follow-up	
Tappin 2015 ⁽³⁸²⁾	UK	NS	612	< 24	Routine care and the offer of up to £400 of shopping vouchers. £50 for attending a face to face appointment and setting a quit date, another £50 if at four weeks' post-quit date exhaled carbon monoxide confirmed quitting; a further £100 was provided for continued validated abstinence of exhaled carbon monoxide after 12 weeks; a final £200 voucher was provided for validated abstinence of exhaled carbon monoxide at 34-38 weeks' gestation.	Usual care (offer of a face to appointment), offer of free NRT x 10 weeks if set a quit date, 4 weekly support phone calls	Biochemically validated point prevalence abstinence at 34-38 weeks	High

Table 10.7 Included studies: Social support studies, n = 7

Study	Country	Ethnicity	Participants (n)	Gestation (weeks)	Intervention	Control	Outcome	Risk of bias
Albrecht 1998 ⁽³⁸³⁾	USA	63% African-American	84	≤ 28	Teen Fresh Start (TFS) plus peer support compared with TFS and control	30 minute education session with project nurse and provision of brochures	Biochemically validated point prevalence abstinence 4-6 wks after baseline	High
Albrecht 2006 ⁽³⁸⁴⁾	USA	NS	95	≤ 28	TFS plus a peer buddy	Usual care. Given educational materials and explanation at a 45-60 minute meeting, attendance incentive (lipstick or nail polish)	Biochemically validated point prevalence abstinence after eight weeks and at one year after the intervention (6-11 months)	High

Bullock 2009 ⁽³⁸⁵⁾	USA	95% White	66	< 24	Received eight booklets and nurse delivered social support	Pamphlet from AHA, research team member to call each month to arrange salivary sample, measure exposure, ask some questions for two more interviews	Biochemically validated point prevalence abstinence at 28-32 weeks and at six weeks (0-5 months) postpartum	High
Henrikus 2010 ⁽³⁸⁶⁾	USA	67% racial minority	82	≤ 28	Control and one in person visit and monthly phone sessions with peer supporters. Given a pregnancy scrapbook with pages that related to smoking cessation tasks.	One in person counselling session to increase motivation to quit and to provide information about community cessation resources	Biochemically validated smoking status just prior to EDD and at three (0-5) months postpartum	High
Malchodi 2003 ⁽³⁸⁷⁾	USA	Majority Black	142	< 20	Control and peer counselling (trained) from lay community outreach workers (phone or home visits)	Usual care which included a programme of AAA, provision of self-help materials, smoking cessation counselling as per protocol at each visit	Biochemically validated abstinence at 36 weeks, mean birth weight and proportion low birth weight	High
McBride 2004 ⁽³⁸⁸⁾	USA	77% White	180	≤ 20	Control and booklet, gift items, six counselling calls based on motivational interviewing (three in antenatal period and three in postnatal period) and partner assistance	Usual care, provider advice, mailed American Cancer Society self-help guide designed for pregnant women	Self-reported ppa at 28 weeks (baseline smokers and baseline quitters reported separately) combined continued abstinence at two (0-5), 6 (6-11) and 12 (12-17) months postpartum	High
Solomon 2000 ⁽³⁸⁹⁾	USA	Majority White	151	NS	Control + if intended to quit offered phone peer support by a healthcare professional. Phone peer support from a trained female ex-smoker - weekly, average duration ten minutes	Brief smoking cessation advice from trained healthcare professionals at each of three prenatal visits and provided with printed materials.	Biochemically validated seven day point prevalence abstinence at 28-34 weeks	High

Included studies: Feedback studies, n = 4


Study	Country	Ethnicity	Participants (n)	Gestation (weeks)	Intervention	Control	Outcome	Risk of bias
Bauman 1983 ⁽³⁹⁰⁾	USA	56% Black	79	≤ 32	Control and exhaled CO with feedback on result	A 135 word script which described the relationship between cigarette smoking and CO, and the harmful consequences of smoking during pregnancy was read.	Biochemically validated abstinence six weeks after intervention	High
Cope 2003 ⁽³⁹¹⁾	England	NS	270	NS	Urinary CO with feedback on result. Quit date set and given a leaflet. Information, feedback and encouragement protocol followed at every visit until 36 weeks	Urinary CO measured but without feedback	Biochemically validated point prevalence abstinence at 36 weeks, mean birth weight and preterm birth	High
Stotts 2009 ⁽³⁹²⁾	USA	NS	240	≤ 26	Three armed RCT. Best practice counselling, feedback on results of ultrasound and information about effect of cigarette smoke on the fetus and one 45-50min face to face session of motivational interviewing based counselling.	5As, 5Rs, 10-15 minute counselling (best practice counselling)	Biochemically validated abstinence at eight months gestation	High
Valbo 1994 ⁽³⁹³⁾	Norway	NS	111	18	Offered Windsor self-help manual, extra ultrasound at 32 weeks and two reminders	Information about harm at time of 18 week ultrasound. Given a pamphlet and encouraged to quit	Self-reported abstinence, self-reported reduction, mean cigs per day at delivery and stillbirths	High


Appendix 11 Studies smoking cessation interventions in pregnancy that were excluded from this review


	Exclude	Reason for Exclusion	Explanation
1	Belizan 1995	Intervention	Intervention included advice on nutrition, alcohol and drugs
2	Bullock 1995	Intervention	Intervention included advice on nutrition, alcohol and drugs
3	Byrd 1993	Outcome	Smoking status not reported by intervention group
4	Campbell 2006	Comparison	Assessed the differential effectiveness of two methods of disseminating a smoking cessation programme
5	Cinciripini 2010	Comparison	Counselling compared to an Alternative (Traditional health education or MI)
6	Donatelle 2000	Comparison	Comparison - incentives was part of a multiple intervention, compared to a less intensive intervention
7	Donovan 1977	Outcome	No data on smoking cessation (just mean cigs/day)
8	El-Mohandes 2011	Intervention	Smoking cessation part of multicomponent intervention
9	Graham 1992	Outcome	Unclear how many smokers were in each arm
10	Haddow 1991	Outcome	No data on smoking cessation
11	Haug 2004	Population	Opioid dependent only. In addition, actual figures not reported
12	Heil 2008	Comparison	Intervention group (contingent voucher) versus control (non-contingent voucher) - equally intensive alternative.
13	Hiett 2000	Outcome	Outcomes not reported
14	Hughes 2000	Outcome	Outcomes not reported
15	LeFevre 1995	Outcome	No specific smoking intervention provided. No data on cessation outcomes
16	Lowe 1997	Population	Spontaneous quitters within 3 months of first prenatal visit
17	Lowe 2002	Intervention	Dissemination trial. Smoking cessation rates not included as an aim of this study
18	Manfredi 1999	Population	Unable to separate pregnant women from women attending family planning and paediatric clinics
19	Moore 1998	Outcome	Not clear what proportion of outcomes related to smokers. Low birthweight and preterm births were study outcomes
20	Moore 2002	Outcome	Outcomes for smokers and recent quitters are combined in analysis
21	Olds 1986	Intervention	Multicomponent intervention (nutrition, alcohol, drugs, enhancement of informal support systems; high intensity)


22	Olds 2002	Outcome	Only mean reduction in cotinine reported
23	Polanska 2004	Outcome	Outcomes in late pregnancy not reported - 'shortly after delivery at home' (0 to 5 months postpartum)
24	Reading 1982	Comparison	Three arms based upon visualisation of ultrasound; no specific smoking cessation component. Also, control outcomes not reported
25	Strecher 2000	Population	Included women who had quit since becoming pregnant. In addition, no late pregnancy outcomes (only postpartum).
26	Tuten 2012	Population	Methadone users only
27	Vilches 2009	Outcome	Abstinence not reported: mean cigs/day reported in late pregnancy


Appendix 12 Cost-effectiveness plane showing the ICER for each strategy when all strategies that are more clinically effective are excluded (with e-cigarettes included as a treatment option)
































Appendix 13: References to included studies

1. Evins AE, Mays VK, Rigotti NA, Tisdale T, Cather C, Goff DC. A pilot trial of bupropion added to cognitive behavioral therapy for smoking cessation in schizophrenia. *Nicotine & tobacco research : official journal of the Society for Research on Nicotine and Tobacco*. 2001;3(4):397-403.
2. Evins AE, Cather C, Deckersbach T, Freudenreich O, Culhane MA, Olm-Shipman CM, et al. A double-blind placebo-controlled trial of bupropion sustained-release for smoking cessation in schizophrenia. *Journal of clinical psychopharmacology*. 2005;25(3):218-25.
3. Evins AE, Cather C, Culhane MA, Birnbaum A, Horowitz J, Hsieh E, et al. A 12-week double-blind, placebo-controlled study of bupropion sr added to high-dose dual nicotine replacement therapy for smoking cessation or reduction in schizophrenia. *Journal of clinical psychopharmacology*. 2007;27(4):380-6.
4. George TP, Vessicchio JC, Termine A, Bregartner TA, Feingold A, Rounsaville BJ, et al. A placebo controlled trial of bupropion for smoking cessation in schizophrenia. *Biological psychiatry*. 2002;52(1):53-61.
5. George TP, Vessicchio JC, Sacco KA, Weinberger AH, Dudas MM, Allen TM, et al. A placebo-controlled trial of bupropion combined with nicotine patch for smoking cessation in schizophrenia. *Biological psychiatry*. 2008;63(11):1092-6.
6. Williams JM, Anthenelli RM, Morris CD, Treadow J, Thompson JR, Yunis C, et al. A randomized, double-blind, placebo-controlled study evaluating the safety and efficacy of varenicline for smoking cessation in patients with schizophrenia or schizoaffective disorder. *The Journal of clinical psychiatry*. 2012;73(5):654-60.
7. Baker A, Richmond R, Haile M, Lewin TJ, Carr VJ, Taylor RL, et al. A randomized controlled trial of a smoking cessation intervention among people with a psychotic disorder. *The American journal of psychiatry*. 2006;163(11):1934-42.
8. Williams JM, Steinberg ML, Zimmermann MH, Gandhi KK, Stipelman B, Budsock PD, et al. Comparison of two intensities of tobacco dependence counseling in schizophrenia and schizoaffective disorder. *Journal of substance abuse treatment*. 2010;38(4):384-93.
9. George TP, Ziedonis DM, Feingold A, Pepper WT, Satterburg CA, Winkel J, et al. Nicotine transdermal patch and atypical antipsychotic medications for smoking cessation in schizophrenia. *The American journal of psychiatry*. 2000;157(11):1835-42.
10. Chengappa KN, Perkins KA, Brar JS, Schlicht PJ, Turkin SR, Hetrick ML, et al. Varenicline for smoking cessation in bipolar disorder: a randomized, double-blind, placebo-controlled study. *The Journal of clinical psychiatry*. 2014;75(7):765-72.
11. Ahluwalia JS, Okuyemi K, Nollen N, Choi WS, Kaur H, Pulvers K, et al. The effects of nicotine gum and counseling among African American light smokers: a 2 x 2 factorial design. *Addiction (Abingdon, England)*. 2006;101(6):883-91.

12. Areechon W, Punnotok J. Smoking cessation through the use of nicotine chewing gum: a double-blind trial in Thailand. *Clinical therapeutics*. 1988;10(2):183-6.
13. Blondal T. Controlled trial of nicotine polacrilex gum with supportive measures. *Archives of internal medicine*. 1989;149(8):1818-21.
14. Campbell IA, Lyons E, Prescott RJ. Stopping smoking. Do nicotine chewing-gum and postal encouragement add to doctors' advice. *The Practitioner*. 1987;231(1423):114-7.
15. Clavel F, Benhamou S, Company-Huertas A, Flamant R. Helping people to stop smoking: randomised comparison of groups being treated with acupuncture and nicotine gum with control group. *British medical journal (Clinical research ed)*. 1985;291(6508):1538-9.
16. Clavel-Chapelon F, Paoletti C, Benhamou S. A randomised 2 x 2 factorial design to evaluate different smoking cessation methods. *Revue d'epidemiologie et de sante publique*. 1992;40(3):187-90.
17. Cooper TV, Klesges RC, Debon MW, Zbikowski SM, Johnson KC, Clemens LH. A placebo controlled randomized trial of the effects of phenylpropanolamine and nicotine gum on cessation rates and postcessation weight gain in women. *Addictive behaviors*. 2005;30(1):61-75.
18. Fagerstrom KO. A comparison of psychological and pharmacological treatment in smoking cessation. *Journal of behavioral medicine*. 1982;5(3):343-51.
19. Fagerstrom KO. Effects of nicotine chewing gum and follow-up appointments in physician-based smoking cessation. *Preventive medicine*. 1984;13(5):517-27.
20. Fee WM, Stewart MJ. A controlled trial of nicotine chewing gum in a smoking withdrawal clinic. *The Practitioner*. 1982;226(1363):148-51.
21. Fortmann SP, Killen JD. Nicotine gum and self-help behavioral treatment for smoking relapse prevention: results from a trial using population-based recruitment. *Journal of consulting and clinical psychology*. 1995;63(3):460-8.
22. Quilez Garcia C, Hernando Arizaleta L, Rubio Diaz A, Granero Fernandez EJ, Vila Coll MA, Estruch Riba J. [Double-blind study of the efficacy of nicotine chewing gum for smoking cessation in the primary care setting]. *Atencion primaria*. 1989;6(10):719-26.
23. Garvey AJ, Kinnunen T, Nordstrom BL, Utman CH, Doherty K, Rosner B, et al. Effects of nicotine gum dose by level of nicotine dependence. *Nicotine & tobacco research : official journal of the Society for Research on Nicotine and Tobacco*. 2000;2(1):53-63.
24. Gilbert JR, Wilson DM, Best JA, Taylor DW, Lindsay EA, Singer J, et al. Smoking cessation in primary care. A randomized controlled trial of nicotine-bearing chewing gum. *The Journal of family practice*. 1989;28(1):49-55.

25. Gross J, Johnson J, Sigler L, Stitzer ML. Dose effects of nicotine gum. *Addictive behaviors*. 1995;20(3):371-81.
26. Hall SM, Tunstall C, Rugg D, Jones RT, Benowitz N. Nicotine gum and behavioral treatment in smoking cessation. *Journal of consulting and clinical psychology*. 1985;53(2):256-8.
27. Hall SM, Tunstall CD, Ginsberg D, Benowitz NL, Jones RT. Nicotine gum and behavioral treatment: a placebo controlled trial. *Journal of consulting and clinical psychology*. 1987;55(4):603-5.
28. Hall SM, Munoz RF, Reus VI, Sees KL, Duncan C, Humfleet GL, et al. Mood management and nicotine gum in smoking treatment: a therapeutic contact and placebo-controlled study. *Journal of consulting and clinical psychology*. 1996;64(5):1003-9.
29. Harackiewicz JM, Blair LW, Sansone C, Epstein JA, Stuchell RN. Nicotine gum and self-help manuals in smoking cessation: an evaluation in a medical context. *Addictive behaviors*. 1988;13(4):319-30.
30. Herrera N, Franco R, Herrera L, Partidas A, Rolando R, Fagerstrom KO. Nicotine gum, 2 and 4 mg, for nicotine dependence. A double-blind placebo-controlled trial within a behavior modification support program. *Chest*. 1995;108(2):447-51.
31. Hjalmarsen AI. Effect of nicotine chewing gum in smoking cessation. A randomized, placebo-controlled, double-blind study. *Jama*. 1984;252(20):2835-8.
32. Huber D. Combined and separate treatment effects of nicotine chewing gum and self-control method. *Pharmacopsychiatry*. 1988;21(6):461-2.
33. Hughes JR, Gust SW, Keenan RM, Fenwick JW, Healey ML. Nicotine vs placebo gum in general medical practice. *Jama*. 1989;261(9):1300-5.
34. Hughes JR, Gust SW, Keenan RM, Fenwick JW. Effect of dose on nicotine's reinforcing, withdrawal-suppression and self-reported effects. *The Journal of pharmacology and experimental therapeutics*. 1990;252(3):1175-83.
35. Jamrozik K, Fowler G, Vessey M, Wald N. Placebo controlled trial of nicotine chewing gum in general practice. *British medical journal (Clinical research ed)*. 1984;289(6448):794-7.
36. Jarvis MJ, Raw M, Russell MA, Feyerabend C. Randomised controlled trial of nicotine chewing-gum. *British medical journal (Clinical research ed)*. 1982;285(6341):537-40.
37. Jensen EJ, Schmidt E, Pedersen B, Dahl R. The effect of nicotine, silver acetate, and placebo chewing gum on the cessation of smoking. The influence of smoking type and nicotine dependence. *The International journal of the addictions*. 1991;26(11):1223-31.
38. Killen JD, Maccoby N, Taylor CB. Nicotine gum and self-regulation training in smoking relapse prevention. *Behavior Therapy*. 1984;15(3):234-48.

39. Killen JD, Fortmann SP, Newman B, Varady A. Evaluation of a treatment approach combining nicotine gum with self-guided behavioral treatments for smoking relapse prevention. *Journal of consulting and clinical psychology*. 1990;58(1):85-92.
40. Salvador Llivina T, Marin Tuya D, Gonzalez Quintana J, Iniesta Torres C, Castellvi Barrera E, Muriana Saez C, et al. [Treatment of smoking: efficacy of the use of nicotine chewing gum. Double-blind study]. *Medicina clinica*. 1988;90(16):646-50.
41. Malcolm RE, Sillett RW, Turner JA, Ball KP. The use of nicotine chewing gum as an aid to stopping smoking. *Psychopharmacology*. 1980;70(3):295-6.
42. McGovern PG, Lando HA. An assessment of nicotine gum as an adjunct to freedom from smoking cessation clinics. *Addictive behaviors*. 1992;17(2):137-47.
43. Nakamura M, Saito J, Oshima A, Miyamoto M, Matushita A, Endo S. Effect of nicotine chewing gun in smoking cessation classes. *The Global War. Proceedings of the 7th World Conference on Tobacco and Health; Perth, Western Australia Health Department of Western Australia*. 1990:665-7.
44. Nebot M, Cabezas C. Does nurse counseling or offer of nicotine gum improve the effectiveness of physician smoking-cessation advice? *Family practice research journal*. 1992;12(3):263-70.
45. Niaura R, Goldstein MG, Abrams DB. Matching high- and low-dependence smokers to self-help treatment with or without nicotine replacement. *Preventive medicine*. 1994;23(1):70-7.
46. Niaura R, Abrams DB, Shadel WG, Rohsenow DJ, Monti PM, Sirota AD. Cue exposure treatment for smoking relapse prevention: a controlled clinical trial. *Addiction (Abingdon, England)*. 1999;94(5):685-95.
47. Ockene JK, Kristeller J, Goldberg R, Amick TL, Pekow PS, Hosmer D, et al. Increasing the efficacy of physician-delivered smoking interventions: a randomized clinical trial. *Journal of general internal medicine*. 1991;6(1):1-8.
48. Page AR, Walters DJ, Schlegel RP, Best JA. Smoking cessation in family practice: the effects of advice and nicotine chewing gum prescription. *Addictive behaviors*. 1986;11(4):443-6.
49. Pirie PL, McBride CM, Hellerstedt W, Jeffery RW, Hatsukami D, Allen S, et al. Smoking cessation in women concerned about weight. *American journal of public health*. 1992;82(9):1238-43.
50. Puska P, Bjorkqvist S, Koskela K. Nicotine-containing chewing gum in smoking cessation: a double blind trial with half year follow-up. *Addictive behaviors*. 1979;4(2):141-6.
51. Richmond RL, Makinson RJ, Kehoe LA, Giugni AA, Webster IW. One-year evaluation of three smoking cessation interventions administered by general practitioners. *Addictive behaviors*. 1993;18(2):187-99.
52. Roto P, Ojala A, Sundman K, Jokinen K, Peltomaki R. Nicotine gum and withdrawal from smoking. *Suomen Laakarilehti*. 1987;36:3445-8.

53. Russell MA, Merriman R, Stapleton J, Taylor W. Effect of nicotine chewing gum as an adjunct to general practitioner's advice against smoking. *British medical journal (Clinical research ed)*. 1983;287(6407):1782-5.
54. Schneider NG, Jarvik ME, Forsythe AB, Read LL, Elliott ML, Schweiger A. Nicotine gum in smoking cessation: a placebo-controlled, double-blind trial. *Addictive behaviors*. 1983;8(3):253-61.
55. Segnan N, Ponti A, Battista RN, Senore C, Rosso S, Shapiro SH, et al. A randomized trial of smoking cessation interventions in general practice in Italy. *Cancer causes & control : CCC*. 1991;2(4):239-46.
56. Shiffman S, Ferguson SG, Strahs KR. Quitting by gradual smoking reduction using nicotine gum: a randomized controlled trial. *American journal of preventive medicine*. 2009;36(2):96-104.e1.
57. Tonnesen P, Fryd V, Hansen M, Helsted J, Gunnensen AB, Forchammer H, et al. Effect of nicotine chewing gum in combination with group counseling on the cessation of smoking. *N Engl J Med*. 1988;318(1):15-8.
58. Villa RS, Alvarez ABD, Hermida JRF. Effectiveness of a multicomponent program to quit smoking with and without nicotine chewing gum [Spanish] [Eficacia de un programa multicomponente para dejar de fumar con y sin chicle de nicotina]. *Psicologia Conductual*. 1999;7:107-18.
59. Zelman DC, Brandon TH, Jorenby DE, Baker TB. Measures of affect and nicotine dependence predict differential response to smoking cessation treatments. *Journal of consulting and clinical psychology*. 1992;60(6):943-52.
60. Abelin T, Buehler A, Muller P, Vesanen K, Imhof PR. Controlled trial of transdermal nicotine patch in tobacco withdrawal. *Lancet (London, England)*. 1989;1(8628):7-10.
61. Buchkremer G, Bents H, Horstmann M, Opitz K, Tolle R. Combination of behavioral smoking cessation with transdermal nicotine substitution. *Addictive behaviors*. 1989;14(2):229-38.
62. Tonnesen P, Paoletti P, Gustavsson G, Russell MA, Saracci R, Gulsvik A, et al. Higher dosage nicotine patches increase one-year smoking cessation rates: results from the European CEASE trial. Collaborative European Anti-Smoking Evaluation. *European Respiratory Society. The European respiratory journal*. 1999;13(2):238-46.
63. Daughton DM, Heatley SA, Prendergast JJ, Causey D, Knowles M, Rolf CN, et al. Effect of transdermal nicotine delivery as an adjunct to low-intervention smoking cessation therapy. A randomized, placebo-controlled, double-blind study. *Archives of internal medicine*. 1991;151(4):749-52.
64. Daughton D, Susman J, Sitorius M, Belenky S, Millatmal T, Nowak R, et al. Transdermal nicotine therapy and primary care. Importance of counseling, demographic, and participant selection factors on 1-year quit rates. The Nebraska Primary Practice Smoking Cessation Trial Group. *Archives of family medicine*. 1998;7(5):425-30.
65. Davidson M, Epstein M, Burt R, Schaefer C, Whitworth G, McDonald A. Efficacy and safety of an over-the-counter transdermal nicotine patch as an aid for smoking cessation. *Archives of family medicine*. 1998;7(6):569-74.

66. Ehksam RE, Buhler A, Muller P, Mauli D, Schumacher PM, Howald H, et al. [Weaning of young smokers using a transdermal nicotine patch]. *Schweizerische Rundschau fur Medizin Praxis = Revue suisse de medecine Praxis*. 1991;80(7):145-50.
67. Fiore MC, Kenford SL, Jorenby DE, Wetter DW, Smith SS, Baker TB. Two studies of the clinical effectiveness of the nicotine patch with different counseling treatments. *Chest*. 1994;105(2):524-33.
68. Glavas D, Rumboldt M, Rumboldt Z. Smoking cessation with nicotine replacement therapy among health care workers: randomized double-blind study. *Croatian medical journal*. 2003;44(2):219-24.
69. Glavas D, Rumboldt Z. [Smoking cessation using the transdermal nicotine system]. *Lijecnicki vjesnik*. 2003;125(1-2):8-12.
70. Hays JT, Croghan IT, Schroeder DR, Offord KP, Hurt RD, Wolter TD, et al. Over-the-counter nicotine patch therapy for smoking cessation: results from randomized, double-blind, placebo-controlled, and open label trials. *American journal of public health*. 1999;89(11):1701-7.
71. Hughes JR, Lesmes GR, Hatsukami DK, Richmond RL, Lichtenstein E, Jorenby DE, et al. Are higher doses of nicotine replacement more effective for smoking cessation? *Nicotine & tobacco research : official journal of the Society for Research on Nicotine and Tobacco*. 1999;1(2):169-74.
72. Hurt RD, Lauger GG, Offord KP, Kottke TE, Dale LC. Nicotine-replacement therapy with use of a transdermal nicotine patch--a randomized double-blind placebo-controlled trial. *Mayo Clinic proceedings*. 1990;65(12):1529-37.
73. Hurt RD, Dale LC, Fredrickson PA, Caldwell CC, Lee GA, Offord KP, et al. Nicotine patch therapy for smoking cessation combined with physician advice and nurse follow-up. One-year outcome and percentage of nicotine replacement. *Jama*. 1994;271(8):595-600.
74. Randomised trial of nicotine patches in general practice: results at one year. Imperial Cancer Research Fund General Practice Research Group. *BMJ (Clinical research ed)*. 1994;308(6942):1476-7.
75. Jorenby DE, Leischow SJ, Nides MA, Rennard SI, Johnston JA, Hughes AR, et al. A controlled trial of sustained-release bupropion, a nicotine patch, or both for smoking cessation. *N Engl J Med*. 1999;340(9):685-91.
76. Killen JD, Fortmann SP, Davis L, Varady A. Nicotine patch and self-help video for cigarette smoking cessation. *Journal of consulting and clinical psychology*. 1997;65(4):663-72.
77. Kornitzer M, Boutsen M, Dramaix M, Thijs J, Gustavsson G. Combined use of nicotine patch and gum in smoking cessation: a placebo-controlled clinical trial. *Preventive medicine*. 1995;24(1):41-7.
78. Oncken C, Cooney J, Feinn R, Lando H, Kranzler HR. Transdermal nicotine for smoking cessation in postmenopausal women. *Addictive behaviors*. 2007;32(2):296-309.

79. Paoletti P, Fornai E, Maggiorelli F, Puntoni R, Viegi G, Carrozzi L, et al. Importance of baseline cotinine plasma values in smoking cessation: results from a double-blind study with nicotine patch. *The European respiratory journal*. 1996;9(4):643-51.
80. Piper ME, Smith SS, Schlam TR, Fiore MC, Jorenby DE, Fraser D, et al. A randomized placebo-controlled clinical trial of 5 smoking cessation pharmacotherapies. *Archives of general psychiatry*. 2009;66(11):1253-62.
81. Richmond RL, Harris K, de Almeida Neto A. The transdermal nicotine patch: results of a randomised placebo-controlled trial. *The Medical journal of Australia*. 1994;161(2):130-5.
82. Sachs DP, Sawe U, Leischow SJ. Effectiveness of a 16-hour transdermal nicotine patch in a medical practice setting, without intensive group counseling. *Archives of internal medicine*. 1993;153(16):1881-90.
83. Sonderskov J, Olsen J, Sabroe S, Meillier L, Overvad K. Nicotine patches in smoking cessation: a randomized trial among over-the-counter customers in Denmark. *American journal of epidemiology*. 1997;145(4):309-18.
84. Stapleton JA, Russell MA, Feyerabend C, Wiseman SM, Gustavsson G, Sawe U, et al. Dose effects and predictors of outcome in a randomized trial of transdermal nicotine patches in general practice. *Addiction (Abingdon, England)*. 1995;90(1):31-42.
85. Johnston JA, Lineberry CG, Ascher JA, Davidson J, Khayrallah MA, Feighner JP, et al. A 102-center prospective study of seizure in association with bupropion. *The Journal of clinical psychiatry*. 1991;52(11):450-6.
86. Tonnesen P, Norregaard J, Simonsen K, Sawe U. A double-blind trial of a 16-hour transdermal nicotine patch in smoking cessation. *N Engl J Med*. 1991;325(5):311-5.
87. Westman EC, Levin ED, Rose JE. The nicotine patch in smoking cessation. A randomized trial with telephone counseling. *Archives of internal medicine*. 1993;153(16):1917-23.
88. Otero UB, Perez Cde A, Szklo M, Esteves GA, dePinho MM, Szklo AS, et al. [Randomized clinical trial: effectiveness of the cognitive-behavioral approach and the use of nicotine replacement transdermal patches for smoking cessation among adults in Rio de Janeiro, Brazil]. *Cad Saude Publica*. 2006;22(2):439-49.
89. Cinciripini PM, Cinciripini LG, Wallfisch A, Haque W, Van Vunakis H. Behavior therapy and the transdermal nicotine patch: effects on cessation outcome, affect, and coping. *Journal of consulting and clinical psychology*. 1996;64(2):314-23.
90. Prapavessis H, Cameron L, Baldi JC, Robinson S, Borrie K, Harper T, et al. The effects of exercise and nicotine replacement therapy on smoking rates in women. *Addictive behaviors*. 2007;32(7):1416-32.
91. Wong GY, Wolter TD, Croghan GA, Croghan IT, Offord KP, Hurt RD. A randomized trial of naltrexone for smoking cessation. *Addiction (Abingdon, England)*. 1999;94(8):1227-37.

92. Anthenelli RM, Benowitz NL, West R, St Aubin L, McRae T, Lawrence D, et al. Neuropsychiatric safety and efficacy of varenicline, bupropion, and nicotine patch in smokers with and without psychiatric disorders (EAGLES): a double-blind, randomised, placebo-controlled clinical trial. *Lancet (London, England)*. 2016;387(10037):2507-20.
93. Hjalmarson A, Nilsson F, Sjostrom L, Wiklund O. The nicotine inhaler in smoking cessation. *Archives of internal medicine*. 1997;157(15):1721-8.
94. Leischow SJ, Nilsson F, Franzon M, Hill A, Otte P, Merikle EP. Efficacy of the nicotine inhaler as an adjunct to smoking cessation. *American journal of health behavior*. 1996;20:364-71.
95. Schneider NG, Olmstead R, Nilsson F, Mody FV, Franzon M, Doan K. Efficacy of a nicotine inhaler in smoking cessation: a double-blind, placebo-controlled trial. *Addiction (Abingdon, England)*. 1996;91(9):1293-306.
96. Tonnesen P, Norregaard J, Mikkelsen K, Jorgensen S, Nilsson F. A double-blind trial of a nicotine inhaler for smoking cessation. *Jama*. 1993;269(10):1268-71.
97. Blondal T, Franzon M, Westin A. A double-blind randomized trial of nicotine nasal spray as an aid in smoking cessation. *The European respiratory journal*. 1997;10(7):1585-90.
98. Hjalmarson A, Franzon M, Westin A, Wiklund O. Effect of nicotine nasal spray on smoking cessation. A randomized, placebo-controlled, double-blind study. *Archives of internal medicine*. 1994;154(22):2567-72.
99. Schneider NG, Olmstead R, Mody FV, Doan K, Franzon M, Jarvik ME, et al. Efficacy of a nicotine nasal spray in smoking cessation: a placebo-controlled, double-blind trial. *Addiction (Abingdon, England)*. 1995;90(12):1671-82.
100. Sutherland G, Stapleton JA, Russell MA, Jarvis MJ, Hajek P, Belcher M, et al. Randomised controlled trial of nasal nicotine spray in smoking cessation. *Lancet (London, England)*. 1992;340(8815):324-9.
101. Dautzenberg B, Ruff F, Vaucher M, Maillon P, Jacob N, Kienzler JL. First demonstration of the good efficacy/safety ratio of Nicotinell® 1mg Lozenges (NL 1mg), a new form of nicotine substitution, by randomised clinical trial. *European Respiratory Journal*. 2001;18(suppl 33):12s.
102. Shiffman S, Dresler CM, Hajek P, Gilbert SJ, Targett DA, Strahs KR. Efficacy of a nicotine lozenge for smoking cessation. *Archives of internal medicine*. 2002;162(11):1267-76.
103. Glover ED, Glover PN, Franzon M, Sullivan CR, Cerullo CC, Howell RM, et al. A comparison of a nicotine sublingual tablet and placebo for smoking cessation. *Nicotine & tobacco research : official journal of the Society for Research on Nicotine and Tobacco*. 2002;4(4):441-50.
104. Wallstrom M, Nilsson F, Hirsch JM. A randomized, double-blind, placebo-controlled clinical evaluation of a nicotine sublingual tablet in smoking cessation. *Addiction (Abingdon, England)*. 2000;95(8):1161-71.

105. Tonnesen P, Lauri H, Perfekt R, Mann K, Batra A. Efficacy of a nicotine mouth spray in smoking cessation: a randomised, double-blind trial. *The European respiratory journal*. 2012;40(3):548-54.
106. Kralikova E, Kozak J, Rasmussen T, Cort N. The clinical benefits of NRT-supported smoking reduction. *Nicotine & Tobacco Research*. 2002;4:243.
107. Kralikova E, Kozak JT, Rasmussen T, Gustavsson G, Le Houezec J. Smoking cessation or reduction with nicotine replacement therapy: a placebo-controlled double blind trial with nicotine gum and inhaler. *BMC public health*. 2009;9:433.
108. Wittchen HU, Hoch E, Klotsche J, Muehlig S. Smoking cessation in primary care - a randomized controlled trial of bupropione, nicotine replacements, CBT and a minimal intervention. *International journal of methods in psychiatric research*. 2011;20(1):28-39.
109. Gariti P, Lynch K, Alterman A, Kampman K, Xie H, Varillo K. Comparing smoking treatment programs for lighter smokers with and without a history of heavier smoking. *Journal of substance abuse treatment*. 2009;37(3):247-55.
110. Smith SS, McCarthy DE, Japuntich SJ, Christiansen B, Piper ME, Jorenby DE, et al. Comparative effectiveness of 5 smoking cessation pharmacotherapies in primary care clinics. *Archives of internal medicine*. 2009;169(22):2148-55.
111. Puska P, Korhonen HJ, Vartiainen E, Urjanheimo EL, Gustavsson G, Westin A. Combined use of nicotine patch and gum compared with gum alone in smoking cessation - a clinical trial in North Karelia. *Tobacco control*. 1995;4(3):231-5.
112. Blondal T, Gudmundsson LJ, Olafsdottir I, Gustavsson G, Westin A. Nicotine nasal spray with nicotine patch for smoking cessation: randomised trial with six year follow up. *BMJ (Clinical research ed)*. 1999;318(7179):285-8.
113. Croghan GA, Sloan JA, Croghan IT, Novotny P, Hurt RD, DeKrey WL, et al. Comparison of nicotine patch alone versus nicotine nasal spray alone versus a combination for treating smokers: a minimal intervention, randomized multicenter trial in a nonspecialized setting. *Nicotine & tobacco research : official journal of the Society for Research on Nicotine and Tobacco*. 2003;5(2):181-7.
114. Bohadana A, Nilsson F, Rasmussen T, Martinet Y. Nicotine inhaler and nicotine patch as a combination therapy for smoking cessation: a randomized, double-blind, placebo-controlled trial. *Archives of internal medicine*. 2000;160(20):3128-34.
115. Lerman C, Kaufmann V, Rukstalis M, Patterson F, Perkins K, Audrain-McGovern J, et al. Individualizing nicotine replacement therapy for the treatment of tobacco dependence: a randomized trial. *Annals of internal medicine*. 2004;140(6):426-33.
116. Schnoll RA, Martinez E, Tatum KL, Glass M, Bernath A, Ferris D, et al. Nicotine patch vs. nicotine lozenge for smoking cessation: an effectiveness trial coordinated by the Community Clinical Oncology Program. *Drug and alcohol dependence*. 2010;107(2-3):237-43.

117. Bullen C, Howe C, Laugesen M, McRobbie H, Parag V, Williman J, et al. Electronic cigarettes for smoking cessation: a randomised controlled trial. *Lancet* (London, England) [Internet]. 2013; 382(9905):[1629-37 pp.]. Available from: <http://onlinelibrary.wiley.com/o/cochrane/clcentral/articles/021/CN-00876021/frame.html>.
118. Caponnetto P, Campagna D, Cibella F, Morjaria JB, Caruso M, Russo C, et al. Efficiency and Safety of an eElectronic cigAreTte (ECLAT) as Tobacco Cigarettes Substitute: A Prospective 12-Month Randomized Control Design Study. *PloS one* [Internet]. 2013; 8(6). Available from: <http://onlinelibrary.wiley.com/o/cochrane/clcentral/articles/653/CN-00904653/frame.html>.
119. Ahluwalia JS, Harris KJ, Catley D, Okuyemi KS, Mayo MS. Sustained-release bupropion for smoking cessation in African Americans: a randomized controlled trial. *Jama*. 2002;288(4):468-74.
120. Aubin HJ, Lebargy F, Berlin I, Bidaut-Mazel C, Chemali-Hudry J, Lagrue G. Efficacy of bupropion and predictors of successful outcome in a sample of French smokers: a randomized placebo-controlled trial. *Addiction* (Abingdon, England). 2004;99(9):1206-18.
121. Brown RA, Niaura R, Lloyd-Richardson EE, Strong DR, Kahler CW, Abrantes AM, et al. Bupropion and cognitive-behavioral treatment for depression in smoking cessation. *Nicotine & tobacco research : official journal of the Society for Research on Nicotine and Tobacco*. 2007;9(7):721-30.
122. Cinciripini PM, Robinson JD, Karam-Hage M, Minnix JA, Lam C, Versace F, et al. Effects of varenicline and bupropion sustained-release use plus intensive smoking cessation counseling on prolonged abstinence from smoking and on depression, negative affect, and other symptoms of nicotine withdrawal. *JAMA psychiatry*. 2013;70(5):522-33.
123. Collins BN, Wileyto EP, Patterson F, Rukstalis M, Audrain-McGovern J, Kaufmann V, et al. Gender differences in smoking cessation in a placebo-controlled trial of bupropion with behavioral counseling. *Nicotine & tobacco research : official journal of the Society for Research on Nicotine and Tobacco*. 2004;6(1):27-37.
124. Cox LS, Nollen NL, Mayo MS, Choi WS, Faseru B, Benowitz NL, et al. Bupropion for smoking cessation in African American light smokers: a randomized controlled trial. *Journal of the National Cancer Institute*. 2012;104(4):290-8.
125. Dalsgareth OJ, Hansen NC, Soes-Petersen U, Evald T, Hoegholm A, Barber J, et al. A multicenter, randomized, double-blind, placebo-controlled, 6-month trial of bupropion hydrochloride sustained-release tablets as an aid to smoking cessation in hospital employees. *Nicotine & tobacco research : official journal of the Society for Research on Nicotine and Tobacco*. 2004;6(1):55-61.
126. Fossati R, Apolone G, Negri E, Compagnoni A, La Vecchia C, Mangano S, et al. A double-blind, placebo-controlled, randomized trial of bupropion for smoking cessation in primary care. *Archives of internal medicine*. 2007;167(16):1791-7.

127. Gonzales D, Nides M, Ferry LH, Segall N, Herrero L, Modell J. Retreatment with bupropion SR: results from 12-month follow-up (RP-83). Rapid Communication Poster Abstracts. Society for Research on Nicotine and Tobacco 8th Annual Meeting, February 20-23 Savannah, Georgia. 2002.
128. Gonzales D, Rennard SI, Nides M, Oncken C, Azoulay S, Billing CB, et al. Varenicline, an alpha4beta2 nicotinic acetylcholine receptor partial agonist, vs sustained-release bupropion and placebo for smoking cessation: a randomized controlled trial. *Jama*. 2006;296(1):47-55.
129. Haggstrom FM, Chatkin JM, Sussenbach-Vaz E, Cesari DH, Fam CF, Fritscher CC. A controlled trial of nortriptyline, sustained-release bupropion and placebo for smoking cessation: preliminary results. *Pulmonary pharmacology & therapeutics*. 2006;19(3):205-9.
130. Hall SM, Humfleet GL, Reus VI, Munoz RF, Hartz DT, Maude-Griffin R. Psychological intervention and antidepressant treatment in smoking cessation. *Archives of general psychiatry*. 2002;59(10):930-6.
131. Holt S, Timu-Parata C, Ryder-Lewis S, Weatherall M, Beasley R. Efficacy of bupropion in the indigenous Maori population in New Zealand. *Thorax*. 2005;60(2):120-3.
132. Hurt RD, Sachs DP, Glover ED, Offord KP, Johnston JA, Dale LC, et al. A comparison of sustained-release bupropion and placebo for smoking cessation. *N Engl J Med*. 1997;337(17):1195-202.
133. Jorenby DE, Hays JT, Rigotti NA, Azoulay S, Watsky EJ, Williams KE, et al. Efficacy of varenicline, an alpha4beta2 nicotinic acetylcholine receptor partial agonist, vs placebo or sustained-release bupropion for smoking cessation: a randomized controlled trial. *Jama*. 2006;296(1):56-63.
134. Levine MD, Perkins KA, Kalarchian MA, Cheng Y, Houck PR, Slane JD, et al. Bupropion and cognitive behavioral therapy for weight-concerned women smokers. *Archives of internal medicine*. 2010;170(6):543-50.
135. McCarthy DE, Piasecki TM, Lawrence DL, Jorenby DE, Shiffman S, Fiore MC, et al. A randomized controlled clinical trial of bupropion SR and individual smoking cessation counseling. *Nicotine & tobacco research : official journal of the Society for Research on Nicotine and Tobacco*. 2008;10(4):717-29.
136. Nides M, Oncken C, Gonzales D, Rennard S, Watsky EJ, Anziano R, et al. Smoking cessation with varenicline, a selective alpha4beta2 nicotinic receptor partial agonist: results from a 7-week, randomized, placebo- and bupropion-controlled trial with 1-year follow-up. *Archives of internal medicine*. 2006;166(15):1561-8.
137. Piper ME, Federman EB, McCarthy DE, Bolt DM, Smith SS, Fiore MC, et al. Efficacy of bupropion alone and in combination with nicotine gum. *Nicotine & tobacco research : official journal of the Society for Research on Nicotine and Tobacco*. 2007;9(9):947-54.

138. Rovina N, Nikoloutsou I, Katsani G, Dima E, Fransis K, Roussos C, et al. Effectiveness of pharmacotherapy and behavioral interventions for smoking cessation in actual clinical practice. *Therapeutic advances in respiratory disease*. 2009;3(6):279-87.
139. Schmitz JM, Stotts AL, Mooney ME, Delaune KA, Moeller GF. Bupropion and cognitive-behavioral therapy for smoking cessation in women. *Nicotine & tobacco research : official journal of the Society for Research on Nicotine and Tobacco*. 2007;9(6):699-709.
140. Selby P, Brands B, Stepner N. Retreatment with ZYban SR: 52 week follow-up of a Canadian Multicentre trial (POS3-63). Society for Research on Nicotine and Tobacco 9th Annual Meeting, February 19-22, New Orleans. 2003.
141. GlaxoSmithKline Clinical Trials Register. Study No: SMK 20001. A Multi-Center, Double-Blind, Double-Dummy, Placebo-Controlled, Randomized, Parallel Group, Dose Response Evaluation of a New Chemical Entity (NCE) and ZYBAN (bupropion hydrochloride) Sustained Release (300mg/day) versus Placebo As Aids to Smoking Cessation. <http://www.gsk-clinicalstudyregister.com/files/pdf/812.pdf>
142. Tonnesen P, Tonstad S, Hjalmarson A, Leborgy F, Van Spiegel PI, Hider A, et al. A multicentre, randomized, double-blind, placebo-controlled, 1-year study of bupropion SR for smoking cessation. *Journal of internal medicine*. 2003;254(2):184-92.
143. Uyar M, Filiz A, Bayram N, Elbek O, Herken H, Topcu A, et al. A randomized trial of smoking cessation. Medication versus motivation. *Saudi medical journal*. 2007;28(6):922-6.
144. Zellweger JP, Boelcskei PL, Carrozzi L, Sepper R, Sweet R, Hider AZ. Bupropion SR vs placebo for smoking cessation in health care professionals. *American journal of health behavior*. 2005;29(3):240-9.
145. Stapleton J, West R, Hajek P, Wheeler J, Vangeli E, Abdi Z, et al. Randomized trial of nicotine replacement therapy (NRT), bupropion and NRT plus bupropion for smoking cessation: effectiveness in clinical practice. *Addiction (Abingdon, England)*. 2013;108(12):2193-201.
146. Swan GE, McAfee T, Curry SJ, Jack LM, Javitz H, Dacey S, et al. Effectiveness of bupropion sustained release for smoking cessation in a health care setting: a randomized trial. *Archives of internal medicine*. 2003;163(19):2337-44.
147. Bolliger CT, Issa JS, Posadas-Valay R, Safwat T, Abreu P, Correia EA, et al. Effects of varenicline in adult smokers: a multinational, 24-week, randomized, double-blind, placebo-controlled study. *Clinical therapeutics*. 2011;33(4):465-77.
148. Nakamura M, Oshima A, Fujimoto Y, Maruyama N, Ishibashi T, Reeves KR. Efficacy and tolerability of varenicline, an alpha4beta2 nicotinic acetylcholine receptor partial agonist, in a 12-week, randomized, placebo-controlled, dose-response study with 40-week follow-up for smoking cessation in Japanese smokers. *Clinical therapeutics*. 2007;29(6):1040-56.

149. Niaura R, Hays JT, Jorenby DE, Leone FT, Pappas JE, Reeves KR, et al. The efficacy and safety of varenicline for smoking cessation using a flexible dosing strategy in adult smokers: a randomized controlled trial. *Current medical research and opinion*. 2008;24(7):1931-41.
150. Oncken C, Gonzales D, Nides M, Rennard S, Watsky E, Billing CB, et al. Efficacy and safety of the novel selective nicotinic acetylcholine receptor partial agonist, varenicline, for smoking cessation. *Archives of internal medicine*. 2006;166(15):1571-7.
151. Rennard S, Hughes J, Cinciripini PM, Kralikova E, Raupach T, Arteaga C, et al. A randomized placebo-controlled trial of varenicline for smoking cessation allowing flexible quit dates. *Nicotine & tobacco research : official journal of the Society for Research on Nicotine and Tobacco*. 2012;14(3):343-50.
152. Tsai ST, Cho HJ, Cheng HS, Kim CH, Hsueh KC, Billing CB, Jr., et al. A randomized, placebo-controlled trial of varenicline, a selective alpha4beta2 nicotinic acetylcholine receptor partial agonist, as a new therapy for smoking cessation in Asian smokers. *Clinical therapeutics*. 2007;29(6):1027-39.
153. Wang C, Xiao D, Chan KP, Pothirat C, Garza D, Davies S. Varenicline for smoking cessation: a placebo-controlled, randomized study. *Respirology (Carlton, Vic)*. 2009;14(3):384-92.
154. de Dios MA, Anderson BJ, Stanton C, Audet DA, Stein M. Project Impact: a pharmacotherapy pilot trial investigating the abstinence and treatment adherence of Latino light smokers. *Journal of substance abuse treatment*. 2012;43(3):322-30.
155. Gonzales D, Hajek P, Pliamm L, Nackaerts K, Tseng LJ, McRae TD, et al. Retreatment with varenicline for smoking cessation in smokers who have previously taken varenicline: a randomized, placebo-controlled trial. *Clinical pharmacology and therapeutics*. 2014;96(3):390-6.
156. Heydari G, Talischi F, Tafti SF, Masjedi MR. Quitting smoking with varenicline: parallel, randomised efficacy trial in Iran. *The international journal of tuberculosis and lung disease : the official journal of the International Union against Tuberculosis and Lung Disease*. 2012;16(2):268-72.
157. Williams KE, Reeves KR, Billing CB, Jr., Pennington AM, Gong J. A double-blind study evaluating the long-term safety of varenicline for smoking cessation. *Current medical research and opinion*. 2007;23(4):793-801.
158. Ebbert JO, Hughes JR, West RJ, Rennard SI, Russ C, McRae TD, et al. Effect of varenicline on smoking cessation through smoking reduction: a randomized clinical trial. *Jama*. 2015;313(7):687-94.
159. Aubin HJ, Bobak A, Britton JR, Oncken C, Billing CB, Jr., Gong J, et al. Varenicline versus transdermal nicotine patch for smoking cessation: results from a randomised open-label trial. *Thorax*. 2008;63(8):717-24.
160. Baker TB, Piper ME, Stein JH, Smith SS, Bolt DM, Fraser DL, et al. Effects of Nicotine Patch vs Varenicline vs Combination Nicotine Replacement Therapy on Smoking Cessation at 26 Weeks: A Randomized Clinical Trial. *Jama*. 2016;315(4):371-9.

161. Tsukahara H, Noda K, Saku K. A randomized controlled open comparative trial of varenicline vs nicotine patch in adult smokers: efficacy, safety and withdrawal symptoms (the VN-SEESAW study). *Circulation journal : official journal of the Japanese Circulation Society*. 2010;74(4):771-8.
162. Scharfenberg G, Benndorf S, Kempe G. [Cytisine (Tabex) as a pharmaceutical aid in stopping smoking]. *Das Deutsche Gesundheitswesen*. 1971;26(10):463-5.
163. Vinnikov D, Brimkulov N, Burjubaeva A. A double blind, randomised, placebo-controlled trial of cytisine for smoking cessation in medium-dependent workers. *Journal of Smoking Cessation*. 2008;3(1):57-62.
164. West R, Zatonski W, Cedzynska M, Lewandowska D, Pazik J, Aveyard P, et al. Placebo-controlled trial of cytisine for smoking cessation. *N Engl J Med*. 2011;365(13):1193-200.
165. Walker N, Howe C, Glover M, McRobbie H, Barnes J, Nosa V, et al. Cytisine versus nicotine for smoking cessation. *N Engl J Med*. 2014;371(25):2353-62.
166. Cottraux JA, Harf R, Boissel JP, Schbath J, Bouvard M, Gillet J. Smoking cessation with behaviour therapy of acupuncture--a controlled study. *Behaviour research and therapy*. 1983;21(4):417-24.
167. Gillams J, Lewith GT, Machin D. Acupuncture and group therapy in stopping smoking. *The Practitioner*. 1984;228(1389):341-4.
168. He D, Berg JE, Hostmark AT. Effects of acupuncture on smoking cessation or reduction for motivated smokers. *Preventive medicine*. 1997;26(2):208-14.
169. Lamontagne Y, Annable L, Gagnon MA. Acupuncture for smokers: lack of long-term therapeutic effect in a controlled study. *Canadian Medical Association Journal*. 1980;5:787-90.
170. Leung JP. Smoking cessation by auricular acupuncture and behavioral therapy. *Psychologia*. 1991;34:177-87.
171. Martin GP, Waite PM. The efficacy of acupuncture as an aid to stopping smoking. *The New Zealand medical journal*. 1981;93(686):421-3.
172. Vandevenne A, Rempp M, Burghard G. Study of the specific contribution of acupuncture to tobacco detoxication [Etude de l'action spécifique de l'acupuncture dans la cure de sevrage tabagique]. *Sem Hôpital Paris*. 1985;61:2155-60.
173. Waite NR, Clough JB. A single-blind, placebo-controlled trial of a simple acupuncture treatment in the cessation of smoking. *The British journal of general practice : the journal of the Royal College of General Practitioners*. 1998;48(433):1487-90.
174. Wu TP, Chen FP, Liu JY, Lin MH, Hwang SJ. A randomized controlled clinical trial of auricular acupuncture in smoking cessation. *Journal of the Chinese Medical Association : JCMA*. 2007;70(8):331-8.

175. Bakkevig O, Steine S, von Hafenbradl K, Laerum E. Smoking cessation. A comparative, randomised study between management in general practice and the behavioural programme SmokEnders. *Scandinavian journal of primary health care*. 2000;18(4):247-51.
176. Batra A, Brömer A, Grüninger K, Schupp P, Buchkremer G. Behaviour therapy for smoking cessation in medical practices [Verhaltenstherapie Raucherentwöhnung in Arztpraxen]. *Verhaltensmodifikation und Verhaltensmedizin*. 1994;15(4):364-76.
177. Camarelles F, Asensio A, Jimenez-Ruiz C, Becerril B, Rodero D, Vidaller O. [Effectiveness of a group therapy intervention to quit smoking. Randomized clinical trial]. *Medicina clinica*. 2002;119(2):53-7.
178. Curry SJ, Marlatt GA, Gordon J, Baer JS. A comparison of alternative theoretical approaches to smoking cessation and relapse. *Health psychology : official journal of the Division of Health Psychology, American Psychological Association*. 1988;7(6):545-56.
179. Flay BR, Gruder CL, Warnecke RB, Jason LA, Peterson P. One year follow-up of the Chicago televised smoking cessation program. *American journal of public health*. 1989;79(10):1377-80.
180. Jason LA, Lesowitz T, Michaels M, Blitz C, Victors L, Dean L, et al. A worksite smoking cessation intervention involving the media and incentives. *American journal of community psychology*. 1989;17(6):785-99.
181. Jason LA, McMahon SD, Salina D, Hedeker D, Stockton M, Dunson K, et al. Assessing a smoking cessation intervention involving groups, incentives, and self-help manuals. *Behavior Therapy*. 1995;26:393-408.
182. Garcia MP, Becona E. Evaluation of the amount of therapist contact in a smoking cessation program. *The Spanish journal of psychology*. 2000;3(1):28-36.
183. Ginsberg D, Hall SM, Rosinski M. Partner support, psychological treatment, and nicotine gum in smoking treatment: an incremental study. *The International journal of the addictions*. 1992;27(5):503-14.
184. Glasgow RE, Schafer L, O'Neill HK. Self-help books and amount of therapist contact in smoking cessation programs. *Journal of consulting and clinical psychology*. 1981;49(5):659-67.
185. Gruder CL, Mermelstein RJ, Kirkendol S, Hedeker D, Wong SC, Schreckengost J, et al. Effects of social support and relapse prevention training as adjuncts to a televised smoking-cessation intervention. *Journal of consulting and clinical psychology*. 1993;61(1):113-20.
186. Hill R, Rigdon M, Johnson S. Behavioral smoking cessation treatment for older chronic smokers. *Behavior Therapy*. 1993;24:321-9.
187. Hilleman DE, Mohiuddin SM, Delcore MG, Lucas BD, Jr. Randomized, controlled trial of transdermal clonidine for smoking cessation. *The Annals of pharmacotherapy*. 1993;27(9):1025-8.

188. Hollis JF, Lichtenstein E, Vogt TM, Stevens VJ, Biglan A. Nurse-assisted counseling for smokers in primary care. *Annals of internal medicine*. 1993;118(7):521-5.
189. Jorenby DE, Smith SS, Fiore MC, Hurt RD, Offord KP, Croghan IT, et al. Varying nicotine patch dose and type of smoking cessation counseling. *Jama*. 1995;274(17):1347-52.
190. McDowell I, Mothersill K, Rosser W, Hartman R. A randomized trial of three approaches to smoking cessation. *Canadian family physician Medecin de famille canadien*. 1985;31:845-51.
191. Minthorn-Biggs MB. Smoking cessation using an interpersonal coping skills program. *Dissertation Abstracts International Section A: Humanities and Social Sciences*. 2000;60(8-A):2814.
192. Nevid JS, Javier RA. Preliminary investigation of a culturally specific smoking cessation intervention for Hispanic smokers. *Am J Health Promot*. 1997;11(3):198-207.
193. Omenn GS, Thompson B, Sexton M, Hessol N, Breitenstein B, Curry S, et al. A randomized comparison of Worksite-sponsored smoking cessation programs. *American journal of preventive medicine*. 1988;4(5):261-7.
194. Pederson LL, Baldwin N, Lefcoe NM. Utility of behavioral self-help manuals in a minimal-contact smoking cessation program. *The International journal of the addictions*. 1981;16(7):1233-9.
195. Pisinger C, Vestbo J, Borch-Johnsen K, Jorgensen T. Smoking cessation intervention in a large randomised population-based study. The Inter99 study. *Preventive medicine*. 2005;40(3):285-92.
196. Rabkin SW, Boyko E, Shane F, Kaufert J. A randomized trial comparing smoking cessation programs utilizing behaviour modification, health education or hypnosis. *Addictive behaviors*. 1984;9(2):157-73.
197. Romand R, Gourgou S, Sancho-Garnier H. A randomized trial assessing the Five-Day Plan for smoking cessation. *Addiction (Abingdon, England)*. 2005;100(10):1546-54.
198. Slovinec D'Angelo ME, Reid RD, Hotz S, Irvine J, Segal RJ, Blanchard CM, et al. Is stress management training a useful addition to physician advice and nicotine replacement therapy during smoking cessation in women? Results of a randomized trial. *Am J Health Promot*. 2005;20(2):127-34.
199. Zheng P, Guo F, Chen Y, Fu Y, Ye T, Fu H. A randomized controlled trial of group intervention based on social cognitive theory for smoking cessation in China. *Journal of epidemiology*. 2007;17(5):147-55.
200. Aleixandre i Marti E, Casanova Matutano MA, Mitjans Lafont J, Sanchez Monfort J, Sanmartin Almenar A. [Clinical trial of 2 tobacco use cessation interventions in primary care]. *Atencion primaria*. 1998;22(7):424-8.
201. Aveyard P, Brown K, Saunders C, Alexander A, Johnstone E, Munafo MR, et al. Weekly versus basic smoking cessation support in primary care: a randomised controlled trial. *Thorax*. 2007;62(10):898-903.

202. Bronson DL, Flynn BS, Solomon LJ, Vacek P, Secker-Walker RH. Smoking cessation counseling during periodic health examinations. *Archives of internal medicine*. 1989;149(7):1653-6.
203. Glasgow RE, Whitlock EP, Eakin EG, Lichtenstein E. A brief smoking cessation intervention for women in low-income planned parenthood clinics. *American journal of public health*. 2000;90(5):786-9.
204. Betson CL, Lam TH, Chung WH, Chung SF. A randomized controlled trial on smoking cessation in government out-patient clinics in Hong Kong [Abstract]. *Proceedings of the 10th World Conference on Tobacco or Health Beijing, China 1997*:113. 1997.
205. Butler CC, Rollnick S, Cohen D, Bachmann M, Russell I, Stott N. Motivational consulting versus brief advice for smokers in general practice: a randomized trial. *The British Journal of General Practice*. 1999;49(445):611-6.
206. Demers RY, Neale AV, Adams R, Trembath C, Herman SC. The impact of physicians' brief smoking cessation counseling: a MIRNET study. *The Journal of family practice*. 1990;31(6):625-9.
207. Haug K, Fugelli P, Aaro LE, Foss OP. Is smoking intervention in general practice more successful among pregnant than non-pregnant women? *Family practice*. 1994;11(2):111-6.
208. Higashi A, Ozasa K, Watanabe Y, Hayashi K, Aoike A, Kawai K, et al. [Efficacy of smoking cessation instruction for general smokers at an annual physical examination]. [*Nihon koshu eisei zasshi*] *Japanese journal of public health*. 1995;42(5):313-21.
209. Janz NK, Becker MH, Kirscht JP, Eraker SA, Billi JE, Woolliscroft JO. Evaluation of a minimal-contact smoking cessation intervention in an outpatient setting. *American journal of public health*. 1987;77(7):805-9.
210. Lang T, Nicaud V, Slama K, Hirsch A, Imbernon E, Goldberg M, et al. Smoking cessation at the workplace. Results of a randomised controlled intervention study. *Worksite physicians from the AIREL group. Journal of epidemiology and community health*. 2000;54(5):349-54.
211. Marshall A, Raw M. Nicotine chewing gum in general practice: effect of follow up appointments. *British medical journal (Clinical research ed)*. 1985;290(6479):1397-8.
212. Meyer C, Ulbricht S, Baumeister SE, Schumann A, Ruge J, Bischof G, et al. Proactive interventions for smoking cessation in general medical practice: a quasi-randomized controlled trial to examine the efficacy of computer-tailored letters and physician-delivered brief advice. *Addiction (Abingdon, England)*. 2008;103(2):294-304.
213. Morgan GD, Noll EL, Orleans CT, Rimer BK, Amfoh K, Bonney G. Reaching midlife and older smokers: tailored interventions for routine medical care. *Preventive medicine*. 1996;25(3):346-54.

214. Nebot-Adell M, Soler-Vila M, Martin-Cantera C, Birules-Pons M, Oller-Colom M, Sala-Carbonell E, et al. Effectiveness of the physician's advice to quit smoking: evaluation of the impact a year after the fact [Efectividad del consejo medico para dejar de fumar: evaluacion del impacto al ano de la intervencion]. *Revista Clinica Española*. 1989;184:210-5.
215. Pieterse ME, Seydel ER, DeVries H, Mudde AN, Kok GJ. Effectiveness of a minimal contact smoking cessation program for Dutch general practitioners: a randomized controlled trial. *Preventive medicine*. 2001;32(2):182-90.
216. Porter AM, McCullough DM. Counselling against cigarette smoking. A controlled study from a general practice. *The Practitioner*. 1972;209(253):686-9.
217. Richmond RL, Austin A, Webster IW. Three year evaluation of a programme by general practitioners to help patients to stop smoking. *British medical journal (Clinical research ed)*. 1986;292(6523):803-6.
218. Russell MA, Wilson C, Taylor C, Baker CD. Effect of general practitioners' advice against smoking. *British medical journal*. 1979;2(6184):231-5.
219. Severson HH, Andrews JA, Lichtenstein E, Wall M, Akers L. Reducing maternal smoking and relapse: long-term evaluation of a pediatric intervention. *Preventive medicine*. 1997;26(1):120-30.
220. Slama K, Redman S, Perkins J, Reid AL, Sanson-Fisher RW. The effectiveness of two smoking cessation programmes for use in general practice: a randomised clinical trial. *BMJ (Clinical research ed)*. 1990;300(6741):1707-9.
221. Thompson RS, Michnich ME, Friedlander L, Gilson B, Grothaus LC, Storer B. Effectiveness of smoking cessation interventions integrated into primary care practice. *Medical care*. 1988;26(1):62-76.
222. Unrod M, Smith M, Spring B, DePue J, Redd W, Winkel G. Randomized controlled trial of a computer-based, tailored intervention to increase smoking cessation counseling by primary care physicians. *Journal of general internal medicine*. 2007;22(4):478-84.
223. Wilson D, Wood G, Johnston N, Sicurella J. Randomized clinical trial of supportive follow-up for cigarette smokers in a family practice. *Canadian Medical Association journal*. 1982;126(2):127-9.
224. Wilson DH, Wakefield MA, Steven ID, Rohrsheim RA, Esterman AJ, Graham NM. "Sick of Smoking": evaluation of a targeted minimal smoking cessation intervention in general practice. *The Medical journal of Australia*. 1990;152(10):518-21.
225. Aveyard P, Griffin C, Lawrence T, Cheng KK. A controlled trial of an expert system and self-help manual intervention based on the stages of change versus standard self-help materials in smoking cessation. *Addiction (Abingdon, England)*. 2003;98(3):345-54.
226. Borrelli B, Novak S, Hecht J, Emmons K, Papandonatos G, Abrams D. Home health care nurses as a new channel for smoking cessation treatment: outcomes from project CARES (Community-nurse Assisted Research and Education on Smoking). *Preventive medicine*. 2005;41(5-6):815-21.

227. Lancaster T, Dobbie W, Vos K, Yudkin P, Murphy M, Fowler G. Randomized trial of nurse-assisted strategies for smoking cessation in primary care. *The British journal of general practice : the journal of the Royal College of General Practitioners*. 1999;49(440):191-4.
228. Sanz-Pozo B, de Miguel-Diez J, Anegon-Blanco M, Garcia-Carballo M, Gomez-Suarez E, Fernandez-Dominguez JF. [Effectiveness of a programme of intensive tobacco counselling by nursing professionals]. *Atencion primaria*. 2006;37(5):266-72.
229. Oncken C, Dornelas E, Greene J, Shankey H, Glasmann A, Feinn R, et al. Nicotine gum for pregnant smokers. *Obstetrics Gynecology*. 2008;112(4):859-67.
230. Brendryen H, Drozd F, Kraft P. A digital smoking cessation program delivered through internet and cell phone without nicotine replacement (happy ending): randomized controlled trial. *Journal of medical Internet research*. 2008;10(5):e51.
231. Brendryen H, Kraft P. Happy ending: a randomized controlled trial of a digital multi-media smoking cessation intervention. *Addiction (Abingdon, England)*. 2008;103(3):478-84; discussion 85-6.
232. Ellerbeck EF, Mahnken JD, Cupertino AP, Cox LS, Greiner KA, Mussulman LM, et al. Effect of varying levels of disease management on smoking cessation: a randomized trial. *Annals of internal medicine*. 2009;150(7):437-46.
233. Elfeddali I, Bolman C, Candel MJ, Wiers RW, de Vries H. Preventing smoking relapse via Web-based computer-tailored feedback: a randomized controlled trial. *Journal of medical Internet research*. 2012;14(4):e109.
234. McDonnell DD, Kazinets G, Lee HJ, Moskowitz JM. An internet-based smoking cessation program for Korean Americans: results from a randomized controlled trial. *Nicotine & tobacco research : official journal of the Society for Research on Nicotine and Tobacco*. 2011;13(5):336-43.
235. Smit ES, de Vries H, Hoving C. Effectiveness of a Web-based multiple tailored smoking cessation program: a randomized controlled trial among Dutch adult smokers. *Journal of medical Internet research*. 2012;14(3):e82.
236. Report on Nurse and Midwife Medicinal Product Prescribing Review of Existing Systems and Processes. Report prepared by: Nursing and Midwifery Board of Ireland and Office of the Nursing and Midwifery Services Director. Health Service Executive. 2015. Available at: <http://www.nmbi.ie/nmbi/media/NMBI/Nurse-Midwife-Prescribing-Review-of-Existing-Systems-Processes-2015.pdf?ext=.pdf>. Accessed 5/12/2016.
237. Bock BC, Becker BM, Niaura RS, Partridge R, Fava JL, Trask P. Smoking cessation among patients in an emergency chest pain observation unit: outcomes of the Chest Pain Smoking Study (CPSS). *Nicotine & tobacco research : official journal of the Society for Research on Nicotine and Tobacco*. 2008;10(10):1523-31.

238. Bock BC, Papandonatos GD, de Dios MA, Abrams DB, Azam MM, Fagan M, et al. Tobacco cessation among low-income smokers: motivational enhancement and nicotine patch treatment. *Nicotine & tobacco research : official journal of the Society for Research on Nicotine and Tobacco*. 2014;16(4):413-22.
239. Davis MF, Shapiro D, Windsor R, Whalen P, Rhode R, Miller HS, et al. Motivational interviewing versus prescriptive advice for smokers who are not ready to quit. *Patient education and counseling*. 2011;83(1):129-33.
240. de Azevedo RC, Mauro ML, Lima DD, Gaspar KC, da Silva VF, Botega NJ. General hospital admission as an opportunity for smoking-cessation strategies: a clinical trial in Brazil. *General hospital psychiatry*. 2010;32(6):599-606.
241. Dornelas EA, Sampson RA, Gray JF, Waters D, Thompson PD. A randomized controlled trial of smoking cessation counseling after myocardial infarction. *Preventive medicine*. 2000;30(4):261-8.
242. Hennrikus DJ, Lando HA, McCarty MC, Klevan D, Holtan N, Huebsch JA, et al. The TEAM project: the effectiveness of smoking cessation intervention with hospital patients. *Preventive medicine*. 2005;40(3):249-58.
243. Hollis JF, McAfee TA, Fellows JL, Zbikowski SM, Stark M, Riedlinger K. The effectiveness and cost effectiveness of telephone counselling and the nicotine patch in a state tobacco quitline. *Tobacco control*. 2007;16 Suppl 1:i53-9.
244. Lindqvist H, Forsberg LG, Forsberg L, Rosendahl I, Enebrink P, Helgason AR. Motivational interviewing in an ordinary clinical setting: a controlled clinical trial at the Swedish National Tobacco Quitline. *Addictive behaviors*. 2013;38(7):2321-4.
245. McClure JB, Westbrook E, Curry SJ, Wetter DW. Proactive, motivationally enhanced smoking cessation counseling among women with elevated cervical cancer risk. *Nicotine & tobacco research : official journal of the Society for Research on Nicotine and Tobacco*. 2005;7(6):881-9.
246. Rigotti NA, Arnsten JH, McKool KM, Wood-Reid KM, Pasternak RC, Singer DE. Efficacy of a smoking cessation program for hospital patients. *Archives of internal medicine*. 1997;157(22):2653-60.
247. Soria R, Legido A, Escolano C, Lopez Yeste A, Montoya J. A randomised controlled trial of motivational interviewing for smoking cessation. *The British journal of general practice : the journal of the Royal College of General Practitioners*. 2006;56(531):768-74.
248. Tevyaw TO, Colby SM, Tidey JW, Kahler CW, Rohsenow DJ, Barnett NP, et al. Contingency management and motivational enhancement: a randomized clinical trial for college student smokers. *Nicotine & tobacco research : official journal of the Society for Research on Nicotine and Tobacco*. 2009;11(6):739-49.
249. Harris KJ, Catley D, Good GE, Cronk NJ, Harrar S, Williams KB. Motivational interviewing for smoking cessation in college students: a group randomized controlled trial. *Preventive medicine*. 2010;51(5):387-93.
250. Borland R, Balmford J, Benda P. Population-level effects of automated smoking cessation help programs: a randomized controlled trial. *Addiction (Abingdon, England)*. 2013;108(3):618-28.

251. Ferguson SG, Walters JAE. The effect of mobile phone text messages on short and long term quitting in motivated smokers: a randomised controlled trial. *Proceedings of the Society for Research on Nicotine and Tobacco 21st Annual Meeting*. 2015;252:POS3-80.
252. Haug S, Schaub MP, Venzin V, Meyer C, John U. Efficacy of a text message-based smoking cessation intervention for young people: a cluster randomized controlled trial. *Journal of medical Internet research*. 2013;15(8):e171.
253. Naughton F, Jamison J, Boase S, Sloan M, Gilbert H, Prevost AT, et al. Randomized controlled trial to assess the short-term effectiveness of tailored web- and text-based facilitation of smoking cessation in primary care (iQuit in practice). *Addiction (Abingdon, England)*. 2014;109(7):1184-93.
254. Whittaker R, Dorey E, Bramley D, Bullen C, Denny S, Elley CR, et al. A theory-based video messaging mobile phone intervention for smoking cessation: randomized controlled trial. *Journal of medical Internet research*. 2011;13(1):e10.
255. Abdullah AS, Mak YW, Loke AY, Lam TH. Smoking cessation intervention in parents of young children: a randomised controlled trial. *Addiction (Abingdon, England)*. 2005;100(11):1731-40.
256. An LC, Zhu SH, Nelson DB, Arikian NJ, Nugent S, Partin MR, et al. Benefits of telephone care over primary care for smoking cessation: a randomized trial. *Archives of internal medicine*. 2006;166(5):536-42.
257. Borland R, Segan CJ, Livingston PM, Owen N. The effectiveness of callback counselling for smoking cessation: a randomized trial. *Addiction (Abingdon, England)*. 2001;96(6):881-9.
258. Borland R, Balmford J, Segan C, Livingston P, Owen N. The effectiveness of personalized smoking cessation strategies for callers to a Quitline service. *Addiction (Abingdon, England)*. 2003;98(6):837-46.
259. Borland R, Balmford J, Bishop N, Segan C, Piterman L, McKay-Brown L, et al. In-practice management versus quitline referral for enhancing smoking cessation in general practice: a cluster randomized trial. *Family practice*. 2008;25(5):382-9.
260. Boyle RG, Solberg LI, Asche SE, Maciosek MV, Boucher JL, Pronk NP. Proactive recruitment of health plan smokers into telephone counseling. *Nicotine & tobacco research : official journal of the Society for Research on Nicotine and Tobacco*. 2007;9(5):581-9.
261. Brown S, Hunt G, Owen N. The effect of adding telephone contact to self-instructional smoking-cessation materials. *Behavior Change*. 1992;9:216-22.
262. Curry SJ, McBride C, Grothaus LC, Louie D, Wagner EH. A randomized trial of self-help materials, personalized feedback, and telephone counseling with nonvolunteer smokers. *Journal of consulting and clinical psychology*. 1995;63(6):1005-14.
263. Ebbert JO, Carr AB, Patten CA, Morris RA, Schroeder DR. Tobacco use quitline enrollment through dental practices: a pilot study. *Journal of the American Dental Association (1939)*. 2007;138(5):595-601.

264. Emmons KM, Puleo E, Park E, Gritz ER, Butterfield RM, Weeks JC, et al. Peer-delivered smoking counseling for childhood cancer survivors increases rate of cessation: the partnership for health study. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*. 2005;23(27):6516-23.
265. Ferguson J, Docherty G, Bauld L, Lewis S, Lorgelly P, Boyd KA, et al. Effect of offering different levels of support and free nicotine replacement therapy via an English national telephone quitline: randomised controlled trial. *BMJ (Clinical research ed)*. 2012;344:e1696.
266. Fiore MC, McCarthy DE, Jackson TC, Zehner ME, Jorenby DE, Mielke M, et al. Integrating smoking cessation treatment into primary care: an effectiveness study. *Preventive medicine*. 2004;38(4):412-20.
267. Floter S, Kroger C. [Effectiveness of telephone aftercare following a smoking cessation program for women on inpatient rehabilitation]. *Deutsche medizinische Wochenschrift (1946)*. 2009;134(47):2382-7.
268. Gilbert H, Sutton S. Evaluating the effectiveness of proactive telephone counselling for smoking cessation in a randomized controlled trial. *Addiction (Abingdon, England)*. 2006;101(4):590-8.
269. Girgis S, Adily A, Velasco MJ, Zwar NA, Jalaludin BB, Ward JE. Feasibility, acceptability and impact of a telephone support service initiated in primary medical care to help Arabic smokers quit. *Australian journal of primary health*. 2011;17(3):274-81.
270. Graham AL, Cobb NK, Papandonatos GD, Moreno JL, Kang H, Tinkelman DG, et al. A randomized trial of Internet and telephone treatment for smoking cessation. *Archives of internal medicine*. 2011;171(1):46-53.
271. Halpin HA, McMenamin SB, Rideout J, Boyce-Smith G. The costs and effectiveness of different benefit designs for treating tobacco dependence: results from a randomized trial. *Inquiry : a journal of medical care organization, provision and financing*. 2006;43(1):54-65.
272. Hennrikus DJ, Jeffery RW, Lando HA, Murray DM, Brelje K, Davidann B, et al. The SUCCESS project: the effect of program format and incentives on participation and cessation in worksite smoking cessation programs. *American journal of public health*. 2002;92(2):274-9.
273. Joyce GF, Niaura R, Maglione M, Mongoven J, Larson-Rotter C, Coan J, et al. The effectiveness of covering smoking cessation services for medicare beneficiaries. *Health services research*. 2008;43(6):2106-23.
274. Katz DA, Muehlenbruch DR, Brown RL, Fiore MC, Baker TB. Effectiveness of implementing the agency for healthcare research and quality smoking cessation clinical practice guideline: a randomized, controlled trial. *Journal of the National Cancer Institute*. 2004;96(8):594-603.
275. Lando HA, Hellerstedt WL, Pirie PL, McGovern PG. Brief supportive telephone outreach as a recruitment and intervention strategy for smoking cessation. *American journal of public health*. 1992;82(1):41-6.

276. Lando HA, Rolnick S, Klevan D, Roski J, Cherney L, Lauger G. Telephone support as an adjunct to transdermal nicotine in smoking cessation. *American journal of public health*. 1997;87(10):1670-4.
277. Lichtenstein E, Andrews JA, Lee ME, Glasgow RE, Hampson SE. Using radon risk to motivate smoking reduction: evaluation of written materials and brief telephone counselling. *Tobacco control*. 2000;9(3):320-6.
278. Lichtenstein E, Boles SM, Lee ME, Hampson SE, Glasgow RE, Fellows J. Using radon risk to motivate smoking reduction II: randomized evaluation of brief telephone counseling and a targeted video. *Health education research*. 2008;23(2):191-201.
279. Lipkus IM, Lyna PR, Rimer BK. Using tailored interventions to enhance smoking cessation among African-Americans at a community health center. *Nicotine & tobacco research : official journal of the Society for Research on Nicotine and Tobacco*. 1999;1(1):77-85.
280. Macleod ZR, Charles MA, Arnaldi VC, Adams IM. Telephone counselling as an adjunct to nicotine patches in smoking cessation: a randomised controlled trial. *The Medical journal of Australia*. 2003;179(7):349-52.
281. McBride CM, Scholes D, Grothaus LC, Curry SJ, Ludman E, Albright J. Evaluation of a minimal self-help smoking cessation intervention following cervical cancer screening. *Preventive medicine*. 1999;29(2):133-8.
282. McClure JB, Catz SL, Ludman EJ, Richards J, Riggs K, Grothaus L. Feasibility and acceptability of a multiple risk factor intervention: the Step Up randomized pilot trial. *BMC public health*. 2011;11:167.
283. McFall SL, Michener A, Rubin D, Flay BR, Mermelstein RJ, Burton D, et al. The effects and use of maintenance newsletters in a smoking cessation intervention. *Addictive behaviors*. 1993;18(2):151-8.
284. Metz K, Floter S, Kroger C, Donath C, Piontek D, Gradl S. Telephone booster sessions for optimizing smoking cessation for patients in rehabilitation centers. *Nicotine & tobacco research : official journal of the Society for Research on Nicotine and Tobacco*. 2007;9(8):853-63.
285. Miguez MC, Vazquez FL, Becona E. Effectiveness of telephone contact as an adjunct to a self-help program for smoking cessation: a randomized controlled trial in Spanish smokers. *Addictive behaviors*. 2002;27(1):139-44.
286. Miguez MC, Becona E. Evaluating the effectiveness of a single telephone contact as an adjunct to a self-help intervention for smoking cessation in a randomized controlled trial. *Nicotine & tobacco research : official journal of the Society for Research on Nicotine and Tobacco*. 2008;10(1):129-35.
287. Miller NH, Smith PM, DeBusk RF, Sobel DS, Taylor CB. Smoking cessation in hospitalized patients. Results of a randomized trial. *Archives of internal medicine*. 1997;157(4):409-15.
288. Orleans CT, Schoenbach VJ, Wagner EH, Quade D, Salmon MA, Pearson DC, et al. Self-help quit smoking interventions: effects of self-help materials, social support instructions, and telephone counseling. *Journal of consulting and clinical psychology*. 1991;59(3):439-48.

289. Orleans CT, Boyd NR, Bingle R, Sutton C, Fairclough D, Heller D, et al. A self-help intervention for African American smokers: tailoring cancer information service counseling for a special population. *Preventive medicine*. 1998;27(5 Pt 2):S61-70.
290. Osinubi OY, Moline J, Rovner E, Sinha S, Perez-Lugo M, Demissie K, et al. A pilot study of telephone-based smoking cessation intervention in asbestos workers. *Journal of occupational and environmental medicine*. 2003;45(5):569-74.
291. Ossip-Klein DJ, Giovino GA, Megahed N, Black PM, Emont SL, Stiggins J, et al. Effects of a smoker's hotline: results of a 10-county self-help trial. *Journal of consulting and clinical psychology*. 1991;59(2):325-32.
292. Ossip-Klein DJ, Carosella AM, Krusch DA. Self-help interventions for older smokers. *Tobacco control*. 1997;6(3):188-93.
293. Prochaska JO, DiClemente CC, Velicer WF, Rossi JS. Standardized, individualized, interactive, and personalized self-help programs for smoking cessation. *Health psychology : official journal of the Division of Health Psychology, American Psychological Association*. 1993;12(5):399-405.
294. Prochaska JO, Velicer WF, Fava JL, Ruggiero L, Laforge RG, Rossi JS, et al. Counselor and stimulus control enhancements of a stage-matched expert system intervention for smokers in a managed care setting. *Preventive medicine*. 2001;32(1):23-32.
295. Rabius V, Pike KJ, Hunter J, Wiatrek D, McAlister AL. Effects of frequency and duration in telephone counselling for smoking cessation. *Tobacco control*. 2007;16 Suppl 1:i71-4.
296. Reid RD, Pipe A, Dafoe WA. Is telephone counselling a useful addition to physician advice and nicotine replacement therapy in helping patients to stop smoking? A randomized controlled trial. *CMAJ : Canadian Medical Association journal = journal de l'Association medicale canadienne*. 1999;160(11):1577-81.
297. Rimer BK, Orleans CT, Fleisher L, Cristinzio S, Resch N, Telepchak J, et al. Does tailoring matter? The impact of a tailored guide on ratings and short-term smoking-related outcomes for older smokers. *Health education research*. 1994;9(1):69-84.
298. Sims TH, McAfee T, Fraser DL, Baker TB, Fiore MC, Smith SS. Quitline cessation counseling for young adult smokers: a randomized clinical trial. *Nicotine & tobacco research : official journal of the Society for Research on Nicotine and Tobacco*. 2013;15(5):932-41.
299. Smith PM, Cameron R, McDonald PW, Kawash B, Madill C, Brown KS. Telephone counseling for population-based smoking cessation. *American journal of health behavior*. 2004;28(3):231-41.
300. Solomon L, Secker-Walke rR, Flynn B, Skelly J, EL C. Proactive telephone peer support to help pregnant women stop smoking. *Tobacco control*. 2000;9(Suppl 3):iii72-iii4.
301. Solomon LJ, Marcy TW, Howe KD, Skelly JM, Reinier K, Flynn BS. Does extended proactive telephone support increase smoking cessation among low-income women using nicotine patches? *Preventive medicine*. 2005;40(3):306-13.

302. Sood A, Andoh J, Verhulst S, Ganesh M, Edson B, Hopkins-Price P. "Real-world" effectiveness of reactive telephone counseling for smoking cessation: a randomized controlled trial. *Chest*. 2009;136(5):1229-36.
303. Sorensen G, Barbeau EM, Stoddard AM, Hunt MK, Goldman R, Smith A, et al. Tools for health: the efficacy of a tailored intervention targeted for construction laborers. *Cancer causes & control : CCC*. 2007;18(1):51-9.
304. Swan GE, McClure JB, Jack LM, Zbikowski SM, Javitz HS, Catz SL, et al. Behavioral counseling and varenicline treatment for smoking cessation. *American journal of preventive medicine*. 2010;38(5):482-90.
305. Thompson B, Kinne S, Lewis FM, Woolridge JA. Randomized telephone smoking-intervention trial initially directed at blue-collar workers. *Journal of the National Cancer Institute Monographs*. 1993(14):105-12.
306. Tzelepis F, Paul CL, Wiggers J, Walsh RA, Knight J, Duncan SL, et al. A randomised controlled trial of proactive telephone counselling on cold-called smokers' cessation rates. *Tobacco control*. 2011;20(1):40-6.
307. Velicer WF, Friedman RH, Fava JL, Gulliver SB, Keller S, Sun X, et al. Evaluating nicotine replacement therapy and stage-based therapies in a population-based effectiveness trial. *Journal of consulting and clinical psychology*. 2006;74(6):1162-72.
308. Young JM, Girgis S, Bruce TA, Hobbs M, Ward JE. Acceptability and effectiveness of opportunistic referral of smokers to telephone cessation advice from a nurse: a randomised trial in Australian general practice. *BMC family practice*. 2008;9:16.
309. Zhu SH, Stretch V, Balabanis M, Rosbrook B, Sadler G, Pierce JP. Telephone counseling for smoking cessation: effects of single-session and multiple-session interventions. *Journal of consulting and clinical psychology*. 1996;64(1):202-11.
310. Zhu SH, Anderson CM, Tedeschi GJ, Rosbrook B, Johnson CE, Byrd M, et al. Evidence of real-world effectiveness of a telephone quitline for smokers. *N Engl J Med*. 2002;347(14):1087-93.
311. Zhu SH, Cummins SE, Wong S, Gamst AC, Tedeschi GJ, Reyes-Nocon J. The effects of a multilingual telephone quitline for Asian smokers: a randomized controlled trial. *Journal of the National Cancer Institute*. 2012;104(4):299-310.
312. Tuisku A, Salmela M, Nieminen P, Toljamo T. Varenicline and Nicotine Patch Therapies in Young Adults Motivated to Quit Smoking: A Randomized, Placebo-controlled, Prospective Study. *Basic and Clinical Pharmacology and Toxicology*. 2016.
313. Ward KD, Asfar T, Al Ali R, Rastam S, Weg MW, Eissenberg T, et al. Randomized trial of the effectiveness of combined behavioral/pharmacological smoking cessation treatment in Syrian primary care clinics. *Addiction (Abingdon, England)*. 2013;108(2):394-403.
314. Ebbert JO, Hatsukami DK, Croghan IT, Schroeder DR, Allen SS, Hays JT, et al. Combination varenicline and bupropion SR for tobacco-dependence treatment in cigarette smokers: a randomized trial. *Jama*. 2014;311(2):155-63.
315. Ramon JM, Morchon S, Baena A, Masuet-Aumatell C. Combining varenicline and nicotine patches: a randomized controlled trial study in smoking cessation. *BMC medicine [Internet]*. 2014; 12:[172 p.]. Available from: <http://onlinelibrary.wiley.com/o/cochrane/clcentral/articles/789/CN-01066789/frame.html>

http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4198796/pdf/12916_2014_Article_172.pdf.

316. Koegelenberg CF, Noor F, Bateman ED, van Zyl-Smit RN, Bruning A, O'Brien JA, et al. Efficacy of varenicline combined with nicotine replacement therapy vs varenicline alone for smoking cessation: a randomized clinical trial. *Jama*. 2014;312(2):155-61.
317. Zincir SB, Zincir S, Kaymak E, Sunbul EA. Comparison of the effectiveness of varenicline, extended-release bupropion and nicotine replacement therapy on the success and the maintenance of a smoking cessation program. *Klinik Psikofarmakoloji Bulteni*. 2013;23(3):224-30.
318. Tulloch HE, Pipe AL, Els C, Clyde MJ, Reid RD. Flexible, dual-form nicotine replacement therapy or varenicline in comparison with nicotine patch for smoking cessation: A randomized controlled trial. *BMC Medicine*. 2016;14(1).
319. Killen JD, Fortmann SP, Schatzberg AF, Arredondo C, Murphy G, Hayward C, et al. Extended cognitive behavior therapy for cigarette smoking cessation. *Addiction (Abingdon, England)*. 2008;103(8):1381-90.
320. Zwar N, Richmond RL, Halcomb E, Furler J, Smith J, Hermiz O, et al. Supporting smoking cessation in Australian primary care: Results of the quit in general practice study. *Global Heart*. 2014;9(1):e232.
321. Stanczyk NE, de Vries H, Candel MJJM, Muris JWM, Bolman CAW. Effectiveness of video- versus text-based computer-tailored smoking cessation interventions among smokers after one year. *Preventive medicine*. 2016;82:42-50.
322. Berlin I, Grange G, Jacob N, Tanguy ML. Nicotine patches in pregnant smokers: randomised, placebo controlled, multicentre trial of efficacy. *BMJ (Clinical research ed)*. 2014;348:g1622.
323. Coleman T, Cooper S, Thornton JG, Grainge MJ, Watts K, Britton J, et al. A randomized trial of nicotine-replacement therapy patches in pregnancy. *New England Journal of Medicine*. 2012;366(9):808-18.
324. Kapur B, Hackman R, Selby P, Klein J, Koren G. Randomized, double-blind, placebo-controlled trial of nicotine replacement therapy in pregnancy. *Current Therapeutic Research*. 2001;62(4):274-8.
325. Wisborg K, Henriksen TB, Jespersen LB, Secher NJ. Nicotine patches for pregnant smokers: a randomized controlled study. *Obstetrics Gynecology*. 2000;96(6):967-71.
326. El-Mohandes AA, Windsor R, Tan S, Perry DC, Gantz MG, Kiely M. A randomized clinical trial of trans-dermal nicotine replacement in pregnant African-American smokers. *Maternal and child health journal [Internet]*. 2013; 17(5):[897-906 pp.]. Available from: <http://onlinelibrary.wiley.com/o/cochrane/clcentral/articles/976/CN-00959976/frame.html>.
327. Hotham ED, Gilbert AL, Atkinson ER. A randomised-controlled pilot study using nicotine patches with pregnant women. *Addictive behaviors*. 2006;31(4):641-8.
328. Pollak KI, Oncken CA, Lipkus IM, Lyna P, Swamy GK, Pletsch PK, et al. Nicotine replacement and behavioral therapy for smoking cessation in pregnancy. *American journal of preventive medicine*. 2007;33(4):297-305.

329. Stotts AL, Northrup TF, Cinciripini PM, Minnix JA, Blalock JA, Mullen PD, et al. Randomized, controlled pilot trial of bupropion for pregnant smokers: challenges and future directions. *American Journal of Perinatology*. 2015;32(4):351-6.
330. Baric L, MacArthur C, Sherwood M. A study of health education aspects of smoking in pregnancy. *International Journal of Health Education*. 1976;19((2 Suppl)):1-17.
331. Cinciripini P, McClure J, Wetter D, Perry J, Blalock J, Cinciripini L, et al. An evaluation of videotaped vignettes for smoking cessation and relapse prevention during pregnancy: The Very Important Pregnant Smokers (VIPS) Program. *Tobacco control*. 2000;9(3):iii61-iii3.
332. Cook C, Ward S, Myers S, Spinnato J. A prospective, randomized evaluation of intensified therapy for smoking reduction in pregnancy. *American journal of obstetrics and gynecology*. 1995;172(1):290.
333. Cummins SE, Tedeschi GJ, Anderson CM, Zhu SH. Telephone Intervention for Pregnant Smokers: A Randomized Controlled Trial. *American journal of preventive medicine*. 2016;51(3):318-26.
334. Dornelas E, Magnavita J, Beazoglou T, Fischer E, Oncken C, Lando H, et al. Efficacy and cost-effectiveness of a clinic-based counseling intervention tested in an ethnically diverse sample of pregnant smokers. *Patient education and counseling*. 2006;64(1-3):342-9.
335. Dunkley J. Training midwives to help pregnant women stop smoking. *Nursing Times* 1997;93(5):64-6.
336. Eades SJ, Sanson-Fisher RW, Panaretto K. An intensive smoking intervention for pregnant Aboriginal and Torres Strait Islander women: a randomised controlled trial. *The Medical journal of Australia*. 2012;198(1):23-4.
337. Ershoff D, Quinn V, Mullen P. A randomized trial of a serialized self-help smoking cessation program for pregnant women in an HMO. *American journal of public health*. 1989;79(2):182-7.
338. Ershoff DH, Quinn VP, Boyd NR, Stern J, Gregory M, Wirtschafter D. The Kaiser Permanente prenatal smoking-cessation trial: when more isn't better, what is enough? *American journal of preventive medicine*. 1999;17(3):161-8.
339. Gielen AC, Windsor R, Faden RR, O'Campo P, Repke J, Davis M. Evaluation of a smoking cessation intervention for pregnant women in an urban prenatal clinic. *Health education research*. 1997;12(2):247-54.
340. Hajek P, Stead L. Aversive smoking for smoking cessation. *Cochrane Database of Systematic Reviews*. 2001(3).
341. Hartmann K, Koch M, Pahel-Short L, Thorp J. A randomized controlled trial of smoking cessation intervention in pregnancy in an academic clinic. *Obstetrics & Gynecology* 1996;87:621-6.
342. Haug K, Fugelli P, Aaro L, Foss O. Is smoking intervention in general practice more successful among pregnant than non-pregnant women? . *Family practice*. 1994;11:111-6.
343. Hegaard HK, Kjaergaard H, Moller LF, Wachmann H, Ottesen B. Multimodal intervention raises smoking cessation rate during pregnancy. *Acta Obstet Gynecol Scand*. 2003;82(9):813-9.

344. Kendrick J, Metzger R, Sexton M, Spierto F, Floyd R, Gargiullo P, et al. Integrating smoking cessation into routine public prenatal care: the Smoking Cessation in Pregnancy project. *American journal of public health.* 1995;85:217-22.
345. Lawrence T, Aveyard P, Evans O, Cheng K. A cluster randomised controlled trial of smoking cessation in pregnant women comparing interventions based on the transtheoretical (stages of change) model to standard care. *Tobacco control.* 2003;12:168–77.
346. Lee M, Miller SM, Wen KY, Hui SK, Roussi P, Hernandez E. Cognitive-behavioral intervention to promote smoking cessation for pregnant and postpartum inner city women. *Journal of behavioral medicine.* 2015;38(6):932-43.
347. Lillington L, Royce J, Novak D, Ruvalcaba M, Chlebowski R. Evaluation of a smoking cessation program for pregnant minority women. *Cancer Pract.* 1995;3(3):157-63.
348. Loeb B, Waage G, Bailey J, editors. *Smoking intervention in pregnancy.* Fifth World Conference on Smoking and Health; 1983; Winnipeg, Canada.: Ottawa: Canadian Council on Smoking and Health.
349. Mayer JP, Hawkins B, Todd R. A randomized evaluation of smoking cessation interventions for pregnant women at a WIC clinic. *American journal of public health.* 1990;80(1):76-8.
350. McBride CM, Curry SJ, Lando HA, Pirie PL, Grothaus LC, Nelson JC. Prevention of relapse in women who quit smoking during pregnancy. *American journal of public health.* 1999;89(5):706-11.
351. McLeod D, Pullon S, Benn C, Cookson T, Dowell A, Viccars A, et al. Can support and education for smoking cessation and reduction be provided effectively by midwives within primary maternity care? *Midwifery.* 2004;20:37–50.
352. Messimer S, Henry R, Hickner J. A comparison of two antismoking interventions among pregnant women in eleven primary care practices. *Journal of Family Practice.* 1989;28(3):283–8.
353. Moore L, Campbell R, Whelan A, Mills N, Lupton P, Misselbrook E, et al. Self help smoking cessation in pregnancy: cluster randomised controlled trial. *British medical journal.* 2002;325:1383–6.
354. Panjari M, Bell R, Bishop S, Astbury J, Rice G, Doery J. A randomized controlled trial of a smoking cessation intervention during pregnancy. *Australian and New Zealand Journal of Obstetrics and Gynaecology.* 1999;39(3):312–7.
355. Parker DR, Windsor RA, Roberts MB, Hecht J, Hardy NV, Strolla LO, et al. Feasibility, cost, and cost-effectiveness of a telephone-based motivational intervention for underserved pregnant smokers. *Nicotine & tobacco research : official journal of the Society for Research on Nicotine and Tobacco.* 2007;9(10):1043-51.
356. Patten CA, Windsor RA, Renner CC, Enoch C, Hochreiter A, Nevak C, et al. Feasibility of a tobacco cessation intervention for pregnant Alaska Native women. *Nicotine & tobacco research : official journal of the Society for Research on Nicotine and Tobacco.* 2009;12(2):79-87.

357. Pbert L, Ockene J, Zapka J, Ma Y, Goins K, Oncken C, et al. A community health center smoking cessation intervention for pregnant and postpartum women. *American journal of preventive medicine*. 2004;26(5):377–85.
358. Price JH, Krol RA, Desmond SM, Losh DP, Roberts SM, Snyder FF. Comparison of three antismoking interventions among pregnant women in an urban setting: a randomized trial. *Psychological reports*. 1991;68(2):595-604.
359. Rigotti NA, Park ER, Regan S, Chang Y, Perry K, Loudin B, et al. Efficacy of telephone counseling for pregnant smokers: a randomized controlled trial. *Obstetrics and gynecology*. 2006;108(1):83-92.
360. Secker-Walker R, Mead P, Goodwin G, Lepage S, Skelly J, Flynn B, et al. Individualised smoking cessation counseling during prenatal and early postnatal care. *American journal of obstetrics and gynecology*. 1994;71:1347–55.
361. Secker-Walker RH, Solomon LJ, Geller BM, Flynn BS, Worden JK, Skelly JM, et al. Modeling smoking cessation: exploring the use of a videotape to help pregnant women quit smoking. *Women & health*. 1997;25(1):23-35.
362. Secker-Walker RH, Solomon LJ, Flynn BS, Skelly JM, Mead PB. Reducing smoking during pregnancy and postpartum: physician's advice supported by individual counseling. *Preventive medicine*. 1998;27(3):422-30.
363. Stotts AL, Diclemente CC, Dolan-Mullen P. One-to-one: a motivational intervention for resistant pregnant smokers. *Addictive behaviors*. 2002;27(2):275-92.
364. Stotts AL, DeLaune KA, Schmitz JM, Grabowski J. Impact of a motivational intervention on mechanisms of change in low-income pregnant smokers. *Addictive behaviors*. 2004;29(8):1649-57.
365. Tappin DM, Lumsden MA, McIntyre D, McKay C, Gilmour WH, Webber R, et al. A pilot study to establish a randomized trial methodology to test the efficacy of a behavioural intervention. *Health education research*. 2000;15(4):491-502.
366. Tappin DM, Lumsden MA, Gilmour WH, Crawford F, McIntyre D, Stone DH, et al. Randomised controlled trial of home based motivational interviewing by midwives to help pregnant smokers quit or cut down. *BMJ (Clinical research ed)*. 2005;331(7513):373-7.
367. Tsoh J, Kohn M, Gerbert B. Promoting smoking cessation in pregnancy with Video Doctor plus provider cueing: a randomized trial. . *Acta Obstetrica et Gynecologica Scandinavica*. 2010;89(4):515–23.
368. Thornton L. Smoking and pregnancy: feasibility and effectiveness of a smoking intervention programme among pregnant women. Dublin: Eastern Health Board., 1997.
369. Valbo A, Eide T. Smoking cessation in pregnancy: the effect of hypnosis in a randomized study. *Addictive behaviors*. 1996;21:29–35.
370. Walsh R, Melmeth A, Byrne J, Brinsmead M, Redman S. A smoking cessation program at a public antenatal clinic. *American journal of public health*. 1997;87:1201–4.

371. Windsor R, Spanos D, Samuelsson C, Bartlett E, Manzella B, Reese Y, et al. The effectiveness of smoking cessation methods for smokers in public health maternity clinics: a randomized trial. *American journal of public health*. 1985;75:1389–92.
372. Windsor R, Lowe J, Perkins L, Smith-Yoder D, Artz L, Crawford M, et al. Health education for pregnant smokers: its behavioral impact and cost benefit. *American journal of public health*. 1993;83:201–6.
373. Windsor R, Woodby L, Miller T, Hardin M. Effectiveness of Smoking Cessation and Reduction in Pregnancy Treatment (SCRIPT) methods in Medicaid-supported prenatal care: Trial III. *Health Education & Behavior*. 2011;38(4):412–22.
374. Burling T, Bigelow G, Robinson J, Mead A. Smoking during pregnancy: reduction via objective assessment and directive advice. *Behavior Therapy*. 1991;22:31-40.
375. Herbec A, Brown J, Tombor I, Michie S, West R. Pilot randomized controlled trial of an internet-based smoking cessation intervention for pregnant smokers ('MumsQuit'). *Drug and alcohol dependence*. 2014;140:130-6.
376. Hjalmarson A, Svanberg B, Hahn L. Stopping smoking in pregnancy: effect of a self-help manual in a controlled trial. *British Journal of Obstetrics and Gynaecology* 1991;98:260–4.
377. Lilley J, Forster D. A randomised controlled trial of individual counselling of smokers in pregnancy. *Public Health* 1986;100:309–15.
378. Naughton F PA, Gilbert H, Sutton S. Randomized controlled trial evaluation of a tailored leaflet and SMS text message self-help intervention for pregnant smokers (MiQuit). *Nicotine and Tobacco Research*. 2012;14(5):569–77.
379. Petersen L, Rosen A, Podedworny T, Kotch J, Handel J. Smoking reduction during pregnancy by a program of self-help and clinical support. *Obstetrics & Gynecology*. 1992;79:924–30.
380. Harris M, Reynolds B. A pilot study of home-based smoking cessation programs for rural, appalachian, pregnant smokers. *Journal of obstetric, gynecologic, and neonatal nursing : JOGNN*. 2015;44(2):236-45.
381. Ondersma S, Svikis D, Lam P, Connors-Burge V, Ledgerwood D, Hopper J. A randomized trial of computer-delivered brief intervention and low-intensity contingency management for smoking during pregnancy. *Nicotine & Tobacco Research* 2012;14(3):351–60.
382. Tappin D, Bauld L, Purves D, Boyd K, Sinclair L, MacAskill S, et al. Financial incentives for smoking cessation in pregnancy: randomised controlled trial. *BMJ (Clinical research ed)*. 2015;350:h134.
383. Albrecht S, Stone C, Payne L, Reynolds M. A preliminary study of the use of peer support in smoking cessation programs for pregnant adolescents. *Journal of the American Academy of Nurse Practitioners* 1998;10:119–25.
384. Albrecht S, Caruthers D, Patrick T, Reynolds M, Salamie D, Higgins L, et al. A randomised controlled trial of a smoking cessation intervention for pregnant adolescents. *Nursing Research* 2006;55(6):402–10.

385. Bullock L, Everett K, Mullen P, Geden E, Longo D, Madsen R, et al. A randomized controlled trial of nurses' individualized social support for poor rural pregnant smokers. *Maternal and Child Health Journal*. 2009;13(3):395–406.
386. Hennrikus D, Pirie P, Hellerstedt W, Lando H, Steele J, Dunn C. Increasing support for smoking cessation during pregnancy and postpartum: results of a randomized controlled pilot study. *Preventive medicine*. 2010;50(3):134–7.
387. Malchodi C, Oncken C, Dornelas E, Caramanica L, Gregonis E. The effects of peer counselling on smoking cessation and reduction. *Obstetrics and gynecology*. 2003;101(3):504–10.
388. McBride C, Baucom D, Peterson B, Pollak K, Palmer C, Westman E, et al. Prenatal and postpartum smoking abstinence: a partner assisted approach. *American journal of preventive medicine*. 2004;27(3):232–8.
389. Solomon LJ, Scharoun GM, Flynn BS, Secker-Walker RH, Sepinwall D. Free nicotine patches plus proactive telephone peer support to help low-income women stop smoking. *Preventive medicine*. 2000;31(1):68-74.
390. Bauman KE, Bryan ES, Dent CW, Koch GG. The influence of observing carbon monoxide level on cigarette smoking by public prenatal patients. *American journal of public health*. 1983;73(9):1089-91.
391. Cope GF, Nayyar P, Holder R. Feedback from a point-of-care test for nicotine intake to reduce smoking during pregnancy. *Annals of clinical biochemistry*. 2003;40(Pt 6):674-9.
392. Stotts AL, Groff JY, Velasquez MM, Benjamin-Garner R, Green C, Carbonari JP, et al. Ultrasound feedback and motivational interviewing targeting smoking cessation in the second and third trimesters of pregnancy. *Nicotine & tobacco research : official journal of the Society for Research on Nicotine and Tobacco*. 2009;11(8):961-8.
393. Valbo A, Nylander G. Smoking cessation in pregnancy. Intervention among heavy smokers. *Acta Obstet Gynecol Scand*. 1994;73(3):215-9.